



UvA-DARE (Digital Academic Repository)

First Ultraviolet Outburst Detected from ASASSN-18eh Strengthens Its Interpretation as a Cataclysmic Variable

Verberne, S.; Modiano, D.; Wijnands, R.

DOI

[10.3847/2515-5172/abd250](https://doi.org/10.3847/2515-5172/abd250)

Publication date

2020

Document Version

Submitted manuscript

Published in

Research Notes of the American Astronomical Society

License

CC BY-NC-ND

[Link to publication](#)

Citation for published version (APA):

Verberne, S., Modiano, D., & Wijnands, R. (2020). First Ultraviolet Outburst Detected from ASASSN-18eh Strengthens Its Interpretation as a Cataclysmic Variable. *Research Notes of the American Astronomical Society*, 4(12), [233]. <https://doi.org/10.3847/2515-5172/abd250>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

First ultraviolet outburst detected from ASASSN-18eh strengthens its interpretation as a cataclysmic variable

SILL VERBERNE,¹ DAVID MODIANO,¹ RUDY WIJNANDS,¹ AND ON BEHALF OF THE TUVO PROJECT¹

¹*Anton Pannekoek Institute for Astronomy, University of Amsterdam
Science Park 904, 1098 XH Amsterdam, Netherlands*

ABSTRACT

As part of the Transient UV Objects project, we have discovered a new outburst (at the beginning of October 2020) of the candidate cataclysmic variable (CV) ASASSN-18eh using the UV/Optical Telescope aboard the Neil Gehrels Swift Observatory. During the outburst its brightness increased by about 6 mag in UV compared to its brightness in the quiescent state. The properties of this outburst are consistent with it being a dwarf nova, strongly supporting the CV nature of ASASSN-18eh.

Keywords: novae, cataclysmic variables — stars: dwarf novae — ultraviolet: stars — white dwarfs

1. INTRODUCTION

Cataclysmic variables (CVs) consist of a white dwarf (WD) in close orbit, typically of a few hours (e.g. Knigge et al. 2011), with usually a red dwarf companion (e.g. Hillman et al. 2020). The companion overflows its Roche lobe and transfers material to the WD through an accretion disk. The accretion often is not stable but goes through episodic periods (lasting a few to a few tens of days) in which the accretion rate suddenly increases significantly causing bright outbursts in between periods of quiescence (so-called dwarf novae, or DNe, increase in V brightness by about $\sim 2 - 5$ mag). See Lasota (2001) for a detailed discussion about DNe and the physics involved.

The source ASASSN-18eh has been suggested as a potential CV after an outburst, reaching a V magnitude of 16.1, was observed in February 2018 (Shappee et al. 2014; ASAS-SN team 2020). As part of our Transient UV Objects (TUVO) project, we have discovered the first outburst in the ultraviolet (UV) of the same source in early October 2020.

2. OBSERVATIONS & ANALYSIS

The Neil Gehrels Swift Observatory (Gehrels et al. 2004) has been designed to provide rapid multi-wavelength follow-up observations of gamma-ray bursts using, among others, its UV/Optical Telescope (UVOT; Roming et al. 2005). The observatory offers an excellent opportunity to study transients in the UV given its many repeated pointings of the same field (owing to its high flexibility), and freely accessible daily data supply (Gehrels et al. 2004).

As part of the TUVO project (Wijnands et al. 2021), we make use of the TUVOpipe pipeline to search for transients in the data from UVOT (Modiano et al. 2021). During October 2020, a transient was detected by the pipeline at a position of 14 28 33.52, -46 11 26.5 (J2000; errors of $\lesssim 5$ arcsec; see Modiano et al. 2021), which is consistent with that of ASASSN-18eh, demonstrating that we detected another outburst of this source. This outburst was not reported by either the Zwicky Transient Facility (Bellm et al. 2019) or ASAS-SN (Shappee et al. 2014).

In total, UVOT has taken 36 exposures of ASASSN-18eh between December 2016 and October 2020. The observations are spread equally between the uw1, um2, and uw2 filters, which have central wavelengths of 2600, 2246, and 1928 Å respectively¹. In nine of the observations (obsIDs 00084507003, 00084507007, and 00084507013; all filters), the source position was right at the boundary of a readout steak caused by a bright star in the field-of-view (FoV) and in another one (obsID 00084507001; only uw1) the source position was located at the edge of the detector and only partially captured. For this reason these exposures have been omitted from the analysis. The resulting total exposure times per filter are ~ 975 , ~ 1284 , and ~ 1284 seconds for uw1 (8 exposures), um2 (9 exposures), and uw2 (9 exposures) respectively.

¹ <https://www.swift.ac.uk/analysis/uvot/filters.php>




Figure 1. Light curve of ASASSN-18eh showing the observed outburst in the uw1, um2, and uw2 filters. The 1σ error bars are plotted for the detections in outburst, along with the 1σ confidence intervals for the detections in quiescence obtained by stacking all the separate observations during which the source was not detected. The time between the last observation in quiescence and the first observation during the outburst is about 1.6 years.

A light curve of the source has been made with the following steps: firstly, all images were aligned per filter, since UVOT pointings can sometimes be offset by a few arcsec (Poole et al. 2008). As source extraction region we used a circular region with a radius of 5 arcsec, centered on the source position as determined by our pipeline. For the background extraction region we used a circular region with a radius of ~ 20 arcsec in an area close to the target, but without any visible sources. All images were then manually inspected for any potential artifacts such as, for example, the previously mentioned readout streaks close to the source. Specialized UVOT tools included with the HEASOFT software package² (Version 6.28) and the HEASARC’s calibration database (CALDB) system (UVOT version 20190101) were then used to analyse the data. Uvotsource³ was used to determine the magnitudes and fluxes of the source for each exposure. In individual exposures during quiescence the source was not detected. Therefore, all exposures during quiescence were stacked using uvotimsum⁴ in order to improve the detection limit, allowing us to detect the source in quiescence. The resulting stacked image for each filter was then again passed to uvotsource using the same source and background extraction regions.

3. RESULTS

Figure 1 shows the light curve of the source in the different filters. The first exposure that shows the source in outburst was taken at 21:21:36 UTC on 2 October 2020 and the last one at 00:28:48 UTC on 4 October 2020. This gives a lower limit for the outburst duration of just over a day. Unfortunately, no more observations were performed after 4 October 2020, since the field could not be observed anymore due to Solar constraints. This means that the source was still active at the time of our last observation, but further constraints are not possible.

Using the image stacking approach discussed above, we were able to detect the source in quiescence with an AB magnitude and flux density of:

² <https://heasarc.gsfc.nasa.gov/docs/software/heasoft/>

³ <https://www.swift.ac.uk/analysis/uvot/mag.php>

⁴ <https://www.swift.ac.uk/analysis/uvot/image.php>

uw1:	21.4 ± 0.3 mag	$4.3(\pm 1.2) \times 10^{-17}$ erg s ⁻¹ cm ⁻² Å ⁻¹	(4.0σ detection)
um2:	21.9 ± 0.4 mag	$3.7(\pm 1.2) \times 10^{-17}$ erg s ⁻¹ cm ⁻² Å ⁻¹	(3.1σ detection)
uw2:	22.2 ± 0.4 mag	$3.4(\pm 1.2) \times 10^{-17}$ erg s ⁻¹ cm ⁻² Å ⁻¹	(2.9σ detection)

This is in contrast with the brightest AB magnitudes and flux densities measured during outburst (i.e. observed during the first outburst observations) of:

uw1:	15.81 ± 0.06 mag	$8.32(\pm 0.46) \times 10^{-15}$ erg s ⁻¹ cm ⁻² Å ⁻¹
um2:	15.95 ± 0.11 mag	$9.24(\pm 0.72) \times 10^{-15}$ erg s ⁻¹ cm ⁻² Å ⁻¹
uw2:	15.89 ± 0.06 mag	$1.21(\pm 0.06) \times 10^{-14}$ erg s ⁻¹ cm ⁻² Å ⁻¹

Therefore, the outburst had an amplitude of at least ~ 6 mag across all filters. It is also noteworthy that all subsequent measurements during the outburst show a decreasing brightness, which indicates that the peak brightnesses were likely higher than the above quoted ones.

4. CONCLUSIONS

We have presented observations of the first detected UV outburst of the candidate CV ASASSN-18eh which was found during October 2020. The UV intensity during the outburst increased by at least ~ 6 mag with a minimum duration of just over a day. Unfortunately, more observations during the October 2020 outburst could not be obtained because of Solar constraints. The next time observations can be obtained is in late December 2020 by which time it is expected that the source has decayed into quiescence again. The maximum UV brightnesses we measured during the October 2020 outburst are slightly higher compared to the brightness in V measured by ASAS-SN during the outburst at the end of February 2018. This is consistent with DNe emitting a large fraction (if not most) of their energy in the UV (e.g. [Giovannelli 2008](#); [Parikh et al. 2019](#)), although we note that we might have missed the peak during the observations making stringent inferences difficult. The previous known outburst (and so far the only one) of the source was in February 2018, meaning that the recurrence time is not more than 2.7 years, although very likely it is shorter because additional outbursts were likely missed by surveying transient facilities. Typical recurrence times of DNe range from days to decades ([Belloni et al. 2016](#)), depending mostly on the mass ratio between the WD and its companion ([Patterson 2011](#)). These characteristics (i.e. outburst amplitude, duration, and recurrence time) are consistent with the outburst being a DN outburst of a CV, which would confirm the CV nature of the source.

Facilities: Swift (UVOT)

REFERENCES

- ASAS-SN team. 2020, ASAS-SN Transients, version Nov 11, 2020. <http://www.astronomy.ohio-state.edu/asassn/transients.html>
- Bellm, E. C., Kulkarni, S. R., Graham, M. J., et al. 2019, PASP, 131, 018002, doi: [10.1088/1538-3873/aaecbe](https://doi.org/10.1088/1538-3873/aaecbe)
- Belloni, D., Giersz, M., Askar, A., Leigh, N., & Hypki, A. 2016, MNRAS, 462, 2950, doi: [10.1093/mnras/stw1841](https://doi.org/10.1093/mnras/stw1841)
- Gehrels, N., Chincarini, G., Giommi, P., et al. 2004, ApJ, 611, 1005, doi: [10.1086/422091](https://doi.org/10.1086/422091)
- Giovannelli, F. 2008, Chinese Journal of Astronomy and Astrophysics Supplement, 8, 237
- Hillman, Y., Shara, M., Prialnik, D., & Kovetz, A. 2020, Nature Astronomy, 4, 886, doi: [10.1038/s41550-020-1062-y](https://doi.org/10.1038/s41550-020-1062-y)
- Knigge, C., Baraffe, I., & Patterson, J. 2011, The Astrophysical Journal Supplement Series, 194, 28, doi: [10.1088/0067-0049/194/2/28](https://doi.org/10.1088/0067-0049/194/2/28)
- Lasota, J.-P. 2001, NewAR, 45, 449, doi: [10.1016/S1387-6473\(01\)00112-9](https://doi.org/10.1016/S1387-6473(01)00112-9)

- Modiano, D., Parikh, A., Van Opijnen, J., et al. 2021, A&A in prep.
- Parikh, A. S., Hernández Santisteban, J. V., Wijnands, R., & Page, D. 2019, RMxAA, 55, 55
- Patterson, J. 2011, Monthly Notices of the Royal Astronomical Society, 411, 2695, doi: [10.1111/j.1365-2966.2010.17881.x](https://doi.org/10.1111/j.1365-2966.2010.17881.x)
- Poole, T. S., Breeveld, A. A., Page, M. J., et al. 2008, MNRAS, 383, 627, doi: [10.1111/j.1365-2966.2007.12563.x](https://doi.org/10.1111/j.1365-2966.2007.12563.x)
- Roming, P. W. A., Kennedy, T. E., Mason, K. O., et al. 2005, Space Science Reviews, 120, 95–142, doi: [10.1007/s11214-005-5095-4](https://doi.org/10.1007/s11214-005-5095-4)
- Shappee, B. J., Prieto, J. L., Grupe, D., et al. 2014, ApJ, 788, 48, doi: [10.1088/0004-637X/788/1/48](https://doi.org/10.1088/0004-637X/788/1/48)
- Wijnands, R., Modiano, D., Parikh, A., et al. 2021, A&A in prep.