



## UvA-DARE (Digital Academic Repository)

### Celebrating Haldane's 'Luttinger liquid theory'

Caux, J.-S.; Morais Smith, C.

**DOI**

[10.1088/1361-648X/aa6012](https://doi.org/10.1088/1361-648X/aa6012)

**Publication date**

2017

**Document Version**

Final published version

**Published in**

Journal of Physics Condensed Matter

**License**

Article 25fa Dutch Copyright Act

[Link to publication](#)

**Citation for published version (APA):**

Caux, J.-S., & Morais Smith, C. (2017). Celebrating Haldane's 'Luttinger liquid theory'. *Journal of Physics Condensed Matter*, 29(15), [151001]. <https://doi.org/10.1088/1361-648X/aa6012>

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.



## Viewpoint

# Celebrating Haldane's 'Luttinger liquid theory'

## Guest Editors

**Jean-Sébastien Caux**  
*Institute for Theoretical Physics Amsterdam and Delta Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands*  
 E-mail: [J.S.Caux@uva.nl](mailto:J.S.Caux@uva.nl)

**Cristiane Morais Smith**  
*Institute for Theoretical Physics, Center for Extreme Matter and Emergent Phenomena, Utrecht University, Leuvenlaan 4, 3584 CE Utrecht, The Netherlands*  
 E-mail: [C.deMoraisSmith@uu.nl](mailto:C.deMoraisSmith@uu.nl)

This Viewpoint celebrates Haldane's seminal (1981 *J. Phys. C: Solid State Phys.* **14** 2585) paper laying the foundations of the modern theory of Luttinger liquids in one-dimensional systems, and was published as part of a series of Viewpoints celebrating 50 of the most influential papers published in the Journal of Physics series, which is celebrating its 50th anniversary.

In a pioneering paper published in *J. Phys. C: Solid State Phys.* in 1981 [1], Duncan Haldane taught us that one-dimensional fermionic systems behave in a fundamentally different manner than those in the universality class described by Landau's Fermi liquid theory [2–4].

At the simplest level, one-dimensionality alters the kinematics of simple excitations around a Fermi sea (figure 1). The spectrum is formed by particle/hole excitations in the vicinity of the two Fermi points (figure 2), and their (multiple) Umklapp modes. In 1D, in contrast to higher dimensions, forbidden regions of the frequency  $\omega$ -momentum  $k$  plane exist. Low-energy modes around integer multiples of the Fermi momentum  $k_F$  are always present (figure 3), but there exist no low-energy excitations in the lobes between points  $2(j-1)k_F$  and  $2jk_F$  for any integer  $j$ . For energies (much) below the height of the lobes (so  $\omega \ll v_F k_F$  in which  $v_F \equiv \left. \frac{d\varepsilon(k)}{dk} \right|_{k_F}$  is the Fermi velocity), we can thus explicitly separate the excitations into different sectors labelled by an even integer  $J = 2j$ .


The lowest-energy state of sector  $J = 2j$  is the  $j$ th Umklapp state at momentum  $2jk_F$ . These states are 'persistent current' modes obtained by a Galilean transformation of the ground state, giving momentum  $2jk_F$  to the whole system, and carrying a quadratic (in  $j$ ) energy shift.

Above the persistent current Umklapp states, but for energies still low on the scale of  $v_F k_F$ , the spectrum of the theory is expected to be given by a linear in momentum boson (sound wave) spectrum (up to nonlinear corrections). Similarly, adding  $\tilde{N}$  particles shifts the Fermi momentum linearly (in  $\tilde{N}$ ), and the energy quadratically. We can thus immediately guess, following Haldane's reasoning in [1], that an effective theory for our fermions universally takes the form


$$H = E_{GS} + \sum_q v_S |q| b_q^\dagger b_q + \frac{2\pi}{L} (v_N \tilde{n}^2 + v_J j^2) \quad (1)$$

where  $E_{GS}$  is the ground-state energy and  $v_S$ ,  $v_N$  and  $v_J$  respectively denote sound, charge and current velocities. It was Haldane's great insight to realize that in the presence of interactions, the effective description based on equation (1) remains valid, albeit with renormalized parameters  $v_S$ ,  $v_N$  and  $v_J$  (constrained by  $v_S^2 = v_N v_J$ ). This theory, which is thus universally valid for gapless one-dimensional systems (be they based on underlying fermionic, bosonic or spin degrees of freedom), should be viewed as a 'theory of everything' in 1D, similarly to the status that Landau's Fermi liquid theory has achieved in higher dimensions. This is the point of view so eloquently put forward in Haldane's paper [1].


Many important contributions paved the way for this synthesis. The development of bosonization goes back almost to the very beginnings of quantum mechanics. In 1934, Bloch [5] used the fact that 1D fermions have the same type of low-energy excitations as a harmonic chain in his study of incoherent x-ray diffraction. Some years later, in 1950, Tomonaga [6] applied Bloch's sound wave method to interacting fermions in 1D. His main contribution was probably to realize that the physical density operator splits up into



**Figure 1.** Left: ground state for a generic system of noninteracting fermions in one dimension. The function  $\varepsilon(k)$  is the one-particle dispersion relation. The chemical potential  $\mu$  sets the value of the Fermi wavevector  $k_F$ . Filled single-particle states are represented by black dots, unfilled ones by open circles. Right: one particle-hole excitation continuum. The (a) and (b) labels refer to the location within the continuum of the particular single particle-hole excitations sketched in figure 2.



**Figure 2.** Simple examples of one particle-hole excitations. The (a) and (b) labels refer to the right panel of figure 1.



**Figure 3.** Multiple particle-hole continuum. The numerals indicate the minimal number of particle-hole excitations needed for a low-energy state to be found in this vicinity. The dashed line indicates the minimal energy parabola for multiple Umklapp states on an exaggerated scale (this is of order  $1/L$  for small numbers of Umklapps and is typically neglected).

left- and right-moving modes, obeying a bosonic Hamiltonian. He however missed the  $2k_F$  contribution to the density correlation function, and did not notice the anomalous decay of correlations. The fundamental paper of Luttinger [7] follows in 1963, in which (perhaps unaware of Tomonaga’s 1950 paper) he formulated his (part)namesake model. Using Toeplitz determinants, he found that the average occupation in the ground state behaves as a power-law with an interaction-dependent anomalous dimension. He thus realized the crucial fact that the Fermi surface discontinuity is destroyed by interactions in 1D. This work was however incorrect in its treatment of the commutation relations of the density operators. Later, Mattis and Lieb offered a correct treatment in their seminal paper [8].

The idea of bosonization, namely that bosons could be used to construct a complete set of states of a 1D fermionic system, appeared in 1965 in [9]. Subsequently, early computations of correlation functions appeared in [10] and [11]. In particular, in the first of

these, Theumann noticed the absence of single-particle poles in the Green's function. She thus correctly concluded that single-particle excitations are absent in such theories. In their famous 1974 paper [12], Dzyaloshinskii and Larkin recovered the absence of single-particle pole and of Fermi surface discontinuity, and offered an interpretation of Mattis and Lieb's solution starting from conventional diagrammatic perturbation theory.

An early version of the actual bosonization operator identity appeared in [13]. This was refined by Mattis in 1974 [14], rendering calculation of correlation functions straightforward. Similar results appeared in the work of Luther and Peschel [15]. The power-law form for correlations was also recovered from equations of motion techniques in [16]. Bosonization was then applied to spin chains and vertex models in [17]. The first precise field-theoretical bosonization formula (as an operator identity) including the (until that point neglected) particle number raising/lowering operators (Klein factors) is in general attributed to [18]. An early formulation of Luttinger liquid concepts appeared in 1975 in [19].

But the fact remains that it is Haldane, in a remarkable series of papers, who gave Luttinger liquid theory the form it has today. Starting in [20], he gave the first explicit construction of charge-raising operators (Klein factors). Subsequently, in [21, 22] and most notably in [1] he offered the complete and explicit construction of the bosonization operator identities, and cross-checked results with exactly solvable models. Most importantly, he proposed the concept of the Luttinger liquid (as he so defined it) as the proper replacement for the Fermi liquid in one dimension, and showed that many different types of systems of fermions, bosons and spins belong to this new universality class. This realization sparked much of the revolutionary advances achieved in low-dimensional quantum systems over the last 35 years.

## Acknowledgments

The authors would like to thank the Stichting voor Fundamenteel Onderzoek der Materie (FOM) and the Nederlandse Organisatie voor Wetenschappelijk onderzoek (NWO).

## References

- [1] Haldane F D M 1981 'Luttinger liquid theory' of one-dimensional quantum fluids. I. Properties of the Luttinger model and their extension to the general 1D interacting spinless Fermi gas *J. Phys. C: Solid State Phys.* **14** 2585
- [2] Landau L D 1957 Theory of Fermi-liquids *Sov. Phys.—JETP* **3** 920
- [3] Landau L D 1957 Oscillations in a Fermi-liquid *Sov. Phys.—JETP* **5** 101
- [4] Abrikosov A A, Gorkov L P and Dzyaloshinski I E 1963 *Methods of Quantum Field Theory in Statistical Physics* (New York: Dover)
- [5] Bloch F 1934 Inkohärente Röntgenstreuung und Dichteschwankungen eines entarteten Fermigas *Helv. Phys. Acta* **7** 385
- [6] Tomonaga S-I 1950 Remarks on Bloch's method of sound waves applied to many-Fermion problems *Prog. Theor. Phys.* **5** 544–69
- [7] Luttinger J M 1963 An exactly soluble model of a many-Fermion system *J. Math. Phys.* **4** 1154–62
- [8] Mattis D C and Lieb E H 1965 Exact solution of a many-Fermion system and its associated boson field *J. Math. Phys.* **6** 304–12
- [9] Overhauser A W 1965 Note on the band theory of magnetism *Physics* **1** 307
- [10] Theumann A 1967 Single-particle Green's function for a one-dimensional many-Fermion system *J. Math. Phys.* **8** 2460–7
- [11] Dover C B 1968 Properties of the Luttinger model *Ann. Phys., NY* **50** 500–33
- [12] Dzyaloshinskii I E and Larkin A I 1974 Correlation functions for a one-dimensional Fermi system with long-range interaction (Tomonaga model) *Sov. Phys.—JETP* **38** 1974  
Dzyaloshinskii I E and Larkin A I 1974 Russian original: *ZhETF Sov. Phys.—JETP* **65** 411
- [13] Schotte K D and Schotte U 1969 Tomonaga's model and the threshold singularity of x-ray spectra of metals *Phys. Rev.* **182** 479–82
- [14] Mattis D C 1974 New wave-operator identity applied to the study of persistent currents in 1D *J. Math. Phys.* **15** 609–12
- [15] Luther A and Peschel I 1974 Single-particle states, Kohn anomaly, and pairing fluctuations in one dimension *Phys. Rev. B* **9** 2911–9
- [16] Everts H U and Schulz H 1974 Application of conventional equation of motion methods to the Tomonaga model *Solid State Commun.* **15** 1413–16

- [17] Luther A and Peschel I 1974 Calculation of critical exponents in two dimensions from quantum field theory in one dimension *Phys. Rev. B* **12** 3908–17
- [18] Heidenreich R, Schroer B, Seiler R and Uhlenbrock D 1975 The sine-Gordon equation and the one-dimensional electron gas *Phys. Lett. A* **54** 119–22
- [19] Efetov K B and Larkin A I 1975 Correlation functions in one-dimensional systems with a strong interaction *Sov. Phys.—JETP* **42** 390
- [20] Haldane F D M 1979 Coupling between charge and spin degrees of freedom in the one-dimensional Fermi gas with backscattering *J. Phys. C: Solid State Phys.* **12** 4791
- [21] Haldane F D M 1980 General relation of correlation exponents and spectral properties of one-dimensional Fermi systems: application to the anisotropic  $S = \frac{1}{2}$  Heisenberg Chain *Phys. Rev. Lett.* **45** 1358–62
- [22] Haldane F D M 1981 Effective harmonic-fluid approach to low-energy properties of one-dimensional quantum fluids *Phys. Rev. Lett.* **47** 1840–3