UvA-DARE (Digital Academic Repository)

Improving risk of bias assessment and quality of reporting in clinical research

Bertizzolo, L.

Publication date

2021

Document Version

Final published version

[Link to publication](#)

Citation for published version (APA):

Bertizzolo, L. (2021). *Improving risk of bias assessment and quality of reporting in clinical research*. [Thesis, fully internal, Universiteit van Amsterdam, Université de Paris].

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

IMPROVING RISK OF BIAS ASSESSMENT AND QUALITY OF REPORTING IN CLINICAL RESEARCH

LORENZO BERTIZZOLO

Université de Paris
En cotutelle avec Universiteit van Amsterdam

Ecole doctorale Pierre Louis de Santé Publique ED 393
Centre de Recherche Épidémiologie et Statistique Sorbonne Paris Cité
INSERM UMR 1153, Équipe METHODS

**Improving risk of bias assessment and quality
of reporting in clinical research**

Lorenzo Bertizzolo

Thèse de doctorat en Épidémiologie clinique

Dirigée par Agnès Dechartres

Et par Patrick Bossuyt

Présentée et soutenue publiquement le 09 juin 2021

Devant un jury composé de :

Rapporteurs

Jörg Meerpohl
Emilie Sbidian

Dr. Prof.
PU-PH

University of Freiburg
Université Paris-Est Créteil

Examineurs

Matthieu Resche-Rigon
Marcel Dijkgraaf
Marc Besselink
Lotty Hooft

PU-PH
Dr. Prof.
Dr. Prof.
Dr.

Université de Paris
University of Amsterdam
University of Amsterdam
University of Utrecht

Directeurs de thèse

Agnès Dechartres
Patrick Bossuyt

PU-PH.
Dr. Prof.

Sorbonne Université
University of Amsterdam

Except where otherwise noted, this is work licensed under
<https://creativecommons.org/licenses/by-nc-nd/3.0/fr/>

Improving risk of bias assessment and quality of reporting in clinical research

ACADEMISCH PROEFSCHRIFT

ter verkrijging van de graad van doctor

aan de Universiteit van Amsterdam

op gezag van de Rector Magnificus

prof. dr. ir. K.I.J. Maex

ten overstaan van een door het College voor Promoties ingestelde commissie,

in het openbaar te verdedigen aan de Université de Paris in Parijs

op woensdag 9 juni, te 9.00 uur

door Lorenzo Bertizzolo

geboren te Trento

Promotiecommissie

Promotores: prof. dr. P.M.M. Bossuyt
prof. dr. A. Dechartres

AMC-UvA
Sorbonne Université

Overige leden: prof. dr. M. Resche-Rigon
prof. dr. J.J. Meerpohl
prof. dr. E. Sbidian
prof. dr. M.G.W. Dijkgraaf
prof. dr. M.G.H. Besselink
dr. L. Hooft

Université de Paris
Universität Freiburg
Université Paris-Est Créteil
AMC-UvA
AMC-UvA
UMC Utrecht

Faculteit der Geneeskunde

Dit proefschrift is tot stand gekomen in het kader van het Europese project 'MiRoR' GA No. 676207, met als doel het behalen van een gezamenlijk doctoraat. Het proefschrift is voorbereid in de Faculteit der Geneeskunde van de Universiteit van Amsterdam en in het Centre de Recherche Épidémiologie et Statistiques van de Université de Paris.

This thesis has been written within the framework of the European project 'MiRoR' GA No. 676207, with the purpose of obtaining a joint doctorate degree. The thesis was prepared in the Faculty of Medicine at the University of Amsterdam and in the Centre de Recherche Épidémiologie et Statistiques at the University of Paris.

Abstract

Research waste is a common problem in biomedical research. High risk of bias and poor quality of reporting are two main causes of waste, which can partly explain the lack of reproducibility of research findings.

The research reported in this thesis aimed to explore different questions related to risk of bias (RoB) assessment and inadequate reporting in clinical research. It consists of five different projects; two of them focused on sources of disagreements when assessing RoB for randomized controlled trials (RCTs) that were included in two different systematic reviews (SRs). A third project evaluated associations between RCT characteristics, including RoB items as well as other features, and reported treatment effects. The other projects explored the quality of reporting in observational studies, through an evaluation of the knowledge and adherence to reporting guidelines and an analysis of the use of causal language in abstracts.

In the first two projects, our results confirm the suboptimal agreement when assessing risk of bias using the RoB tool, with researchers agreeing 70% of the times. The most common reason of disagreements was a different interpretation of the same information and this was most frequently related to incomplete or unclear reporting in the study report. We also evaluated whether external factors related to the review, the authors, or the study itself may influence the risk of bias judgement in case of incomplete or unclear reporting in the study report. Review groups and authors that had completed a lower number of SRs assigned significantly more often a low risk of bias where others reported “unclear”. The study year of publication, compared to the publication date of other studies included in the review, was also associated with a different RoB judgment.

In the third project, evaluating 132 RCTs on testosterone therapy in men, we found no RCT characteristic significantly associated with reported treatment effects. Although trials rated at high overall RoB reported treatment effects that were 21% larger compared with trials rated at low RoB, the difference was not statistically significant.

Regarding observational research, we conducted a survey on the knowledge and use of STROBE, a reporting guideline dedicated to observational studies. Researchers who conducted an observational study were recruited and contacted through various channels (contact with editors, corresponding authors of relevant studies using an observational design, social media). Among 1015 participants, 62.6% had used STROBE in the past while 19% had never heard it.

The last study examined the use of causal language in 60 abstracts of observational studies published in *the BMJ* and found use of causal language in 48% of them, which outlines the need for more guidance.

The projects reported in this thesis explored different questions related to risk of bias and quality of reporting, across different methodologies and topics. The results suggest possible ways of improvement. Better guidance could improve agreement in the RoB assessment in case of incomplete information. Initiatives could be taken to improve adherence to reporting guidelines, as shown in the study on STROBE. Our work on use of causal language in observational research confirms that clarity and consistency on methodology of research is needed.

Keywords: clinical research, risk of bias, Cochrane, reporting, quality of research, waste of research, reporting guidelines

Résumé

Le gâchis de la recherche représente un problème majeur dans la recherche biomédicale. Un risque élevé de biais et une mauvaise qualité des rapports sont deux principales causes de gâchis, pouvant en partie expliquer le manque de reproductibilité des résultats de la recherche.

Cette thèse visait à explorer différentes questions liées à l'évaluation du risque de biais (RoB) et à des rapports incomplets ou inadéquats en recherche clinique. Elle se compose de cinq projets différents ; deux d'entre eux se concentrent sur l'exploration des raisons des désaccords lors de l'évaluation du RoB pour les mêmes essais contrôlés randomisés (ECR) inclus dans deux revues systématiques (RS) différentes. Un troisième projet évalue l'association entre certaines caractéristiques des ECR, y compris les items de l'outil pour évaluer le RoB (mais aussi l'année de publication, la taille de l'échantillon, l'enregistrement de l'essai, le statut mono ou multicentrique, la région, la source de financement et les conflits d'intérêts) et l'effet traitement. Les autres travaux explorent la qualité des rapports dans les études observationnelles, à travers une enquête sur la connaissance et l'utilisation des recommandations STROBE et l'utilisation d'un langage causal dans les résumés d'articles scientifiques.

Dans les deux premiers projets, nos résultats confirment la concordance sous-optimale lors de l'évaluation du risque de biais à l'aide de l'outil RoB, les chercheurs étant d'accord 70% du temps. La raison la plus fréquente de désaccord était une interprétation différente des mêmes informations rapportées dans l'article et ceci était le plus souvent lié à des informations incomplètes ou peu claires dans le rapport d'étude. Nous avons également évalué si des facteurs externes liés à la revue, aux auteurs ou à l'étude elle-même pouvaient influencer le risque de biais en cas d'informations incomplètes ou peu claires dans le rapport d'étude. Les groupes de revue et les auteurs qui avaient participé à un nombre inférieur de revues systématiques ont attribué plus souvent un faible risque de biais, alors que les autres attribuaient un risque de biais « incertain ». L'année de publication de l'étude, comparée à l'année de publication des autres études incluses dans la revue, était également associée à une évaluation différente.

Dans le troisième projet, à partir de 132 ECR évaluant un traitement par testostérone chez les hommes, nous n'avons trouvé aucune caractéristique associée de manière significative à l'effet traitement. Bien que les essais à RoB global élevé ou incertain aient rapporté un effet traitement plus important de l'ordre de 21% par rapport aux essais évalués à RoB global faible, la différence n'était pas statistiquement significative.

Concernant la recherche observationnelle, nous avons mené une enquête sur la connaissance et l'utilisation de STROBE. Les chercheurs ayant mené une étude observationnelle ont été recrutés et

contactés par différents canaux (contact avec les éditeurs, auteurs correspondants d'études observationnelles pertinentes, réseaux sociaux). Parmi 1015 participants, 62,6% avaient utilisé STROBE dans le passé tandis que 19% n'en avaient jamais entendu parler. La dernière étude a examiné l'utilisation du langage causal dans 60 résumés d'études observationnelles publiées dans le *BMJ* et a trouvé un langage causal dans 48% d'entre eux, ce qui souligne l'importance d'avoir des recommandations claires adaptées à cette problématique.

En conclusion, la thèse a exploré différentes questions liées au risque de biais et à la qualité des rapports à travers différentes méthodologies et suggère des pistes d'amélioration possibles. Des recommandations plus précises pourraient améliorer la reproductibilité de l'évaluation du risque de biais en cas d'informations incomplètes. Des initiatives similaires devraient être prises pour améliorer l'adhésion aux recommandations de reporting, comme indiqué dans l'étude sur STROBE. Enfin, le travail sur le langage causal dans les études observationnelles souligne l'importance d'utiliser un langage adapté.

Mots clés : recherche clinique, risque de biais, Cochrane, reporting, qualité de la recherche, gâchis de la recherche

Synthèse des travaux de thèse

1. Introduction et objectifs

La médecine fondée sur les preuves ou *evidence based medicine* (EBM) en anglais a introduit un nouveau paradigme dans la pratique de la médecine et dans la recherche biomédicale. L'EBM est définie comme « l'utilisation consciencieuse, explicite et judicieuse des meilleures données scientifiques actuelles dans la prise en charge personnalisée de chaque patient » [1]. Le principe de l'EBM est que toute décision médicale doit reposer sur les meilleurs éléments de preuve scientifiques issus de la recherche [2]. Bien que l'EBM intègre l'expérience du praticien et les préférences du patient, elle accorde une importance majeure à la littérature scientifique.

Malheureusement, la qualité de la littérature scientifique n'est pas optimale [3]. Il a été ainsi estimé que jusqu'à 85% de la recherche biomédicale était gâché [4, 5]. Ce gâchis survient à chaque étape du processus de recherche, de la planification à la présentation et diffusion des résultats de la recherche. Lors de la planification, les chercheurs peuvent formuler une mauvaise question de recherche ou planifier des méthodes inappropriées. Le gâchis peut ensuite se produire pendant la conduite de l'étude, en utilisant des méthodes susceptibles d'induire un biais dans les résultats. Enfin, il peut y avoir du gâchis en raison de rapports incomplets ou inadéquats des résultats, y compris en utilisant une terminologie inadaptée [6-9]. Dans cette thèse, nous avons cherché à nous concentrer sur deux causes principales de gâchis de la recherche responsable de la mauvaise qualité de la littérature scientifique : un risque de biais élevé et une mauvaise qualité des rapports.

Un biais en épidémiologie est défini comme une erreur systématique, entraînant une sous-estimation ou une surestimation systématique du véritable effet traitement [73]. L'identification et la mesure des biais sont difficiles. Paradoxalement, une étude avec de nombreuses faiblesses méthodologiques peut par hasard être non biaisée, alors que le contraire est également possible. Pour ces raisons, il est jugé plus approprié de se concentrer sur le risque de biais (RoB). Cela se traduit par l'évaluation des principales caractéristiques méthodologiques d'une étude pouvant conduire à des résultats trompeurs [10-13]. En 2005, la Cochrane a commencé à développer un outil pour évaluer le RoB dans les essais contrôlés randomisés (ECR) sur la base d'éléments de preuve issus d'études méta-épidémiologiques évaluant les caractéristiques méthodologiques associées à l'effet traitement [13-18]. L'outil RoB comprend les éléments suivants: Génération de la séquence de randomisation et assignation secrète, faisant référence au risque de biais de sélection; Insu des participants et du personnel, faisant référence au risque de biais de performance; Mise en insu de l'évaluation des critères de jugement, se référant au risque de biais de détection ; Données incomplètes, se référant au risque de biais d'attrition ; Rapport sélectif des critères de jugement ou

des résultats, se référant au risque de biais de présentation sélective des critères de jugement et autres biais. L'évaluation du risque de biais de chaque item se compose de deux parties : une brève description du support du jugement qui comprend généralement des citations textuelles des publications de l'étude évaluée et le jugement concernant le risque de biais. Le risque de biais peut être exprimé en 3 catégories : risque de biais faible, risque de biais élevé et risque de biais incertain s'il n'y avait pas suffisamment d'informations pour se prononcer [13] et la Cochrane fournit des recommandations pour évaluer le risque de biais pour chaque élément. L'outil RoB est rapidement devenu l'outil de référence pour évaluer le risque de biais dans les ECR et est aujourd'hui utilisé dans toutes les revues Cochrane et dans un nombre croissant de revues non Cochrane [19]. Cependant, plusieurs études ont montré une faible reproductibilité globale de l'évaluation réalisée avec l'outil RoB, en particulier pour certains éléments tels que les données incomplètes et les items se rapportant à l'insu [10, 20-25]. Aucune de ces études n'a exploré les raisons qui peuvent expliquer les discordances sur l'évaluation du risque de biais.

Un autre problème important affectant la littérature scientifique est lié à la mauvaise qualité des rapports des études scientifiques. Les rapports d'étude doivent être clairs, complets et facilement disponibles ; si les informations importantes d'une étude sont non ou mal rapportées, cela rend la réplication, l'évaluation critique et l'interprétation des résultats difficiles, voire impossibles. Un mauvais reporting limite donc l'évaluation du risque de biais avec des items généralement considérés comme à risque de biais incertain dans ce cas lors de l'utilisation de l'outil RoB. Avec des rapports incomplets ou inadéquats, les résultats ne peuvent pas non plus guider correctement la pratique clinique ou les politiques de santé, contribuant au problème global du gâchis dans la recherche [26-28]. De nombreuses études ont montré la mauvaise qualité des rapports pour différentes questions de recherche et différents types d'études [15, 29-35]. Pour lutter contre ce problème, la communauté scientifique a développé des recommandations structurées pour la rédaction d'articles scientifiques avec la création d'outils tels que des recommandations de reporting (en anglais *reporting guidelines*) (RG) [4, 36, 37]. Les premiers efforts structurés visant à améliorer la qualité des rapports portaient sur les ECR et remontent à 1993 avec la création de l'outil SORT [38, 39]. Simultanément et indépendamment, un groupe différent travaillait également sur des recommandations. Ainsi, en 1996, le groupe SORT, en collaboration avec le groupe de travail Asilomar, a publié le Consolidated Standards of Reporting Trials Statement (CONSORT) Statement [40-42] qui est considéré comme l'un des premiers RG structurés. Bien qu'initialement conçu pour les ECR, le CONSORT a inspiré de nombreuses extensions afin de fournir des recommandations spécifiques pour différents types d'étude ou d'interventions [43].

Les RG sont souvent organisés sous la forme d'une liste d'items, contenant des éléments essentiels à rapporter dans la publication. Ces listes d'items peuvent également être accompagnées d'un diagramme de flux, détaillant différentes informations, telles que le nombre de personnes initialement contactées, le nombre de personnes qui ont refusé de participer et le nombre d'abandons. Il est fréquemment associé un document « Élaboration et explication », fournissant une description détaillée de l'utilisation de la liste d'items et expliquant le rationnel pour rapporter chaque élément. L'un des RG les plus utilisés pour les études observationnelles est l'outil STROBE (STrengthening the Reporting of OBservational studies in Epidemiology), créé en 2007 [44]. STROBE comprend une liste de 22 items qui détaille les informations clés nécessaires pour rapporter les résultats d'une étude observationnelle. STROBE fournit également des détails et des conseils supplémentaires pour chaque élément de la liste d'items en donnant des exemples de bonnes manières de rapporter [45]. Grâce à ce document explicatif, STROBE a été utilisé par les chercheurs comme matériel de formation, à la fois pour l'utilisation de STROBE, mais aussi pour guider la méthodologie de la recherche [46]. Bien que STROBE soit largement utilisé par les chercheurs depuis plus de 10 ans, il existe peu d'informations sur les connaissances et la perception des chercheurs vis-à-vis de cet outil.

Une autre cause de rapport inadéquat est liée à l'utilisation d'une terminologie inappropriée pouvant induire le lecteur en erreur. On considère généralement que l'ECR est le seul type d'étude capable de produire une inférence causale [47, 48] car la randomisation permet une distribution comparable des facteurs de confusion entre les groupes expérimental et contrôle [49]. Cependant, il n'est pas toujours possible ou adapté de réaliser un ECR, par conséquent, malgré les limites des études observationnelles, les auteurs peuvent être tentés de tirer des conclusions causales dans des études observationnelles [50, 51].

La thèse vise à explorer différentes questions liées à l'évaluation du risque de biais et à des rapports incomplets ou inappropriés des articles scientifiques. Elle se compose de cinq projets différents présentés chacun dans un chapitre séparé. Les deux premiers projets se concentrent sur l'exploration des raisons de désaccords lors de l'évaluation du risque de biais pour les mêmes essais inclus dans deux revues systématiques (RS) Cochrane différentes. Un troisième projet évalue l'association entre certaines caractéristiques des ECR incluant les éléments de l'outil RoB (mais aussi l'année de publication, la taille de l'échantillon, l'enregistrement, le statut monocentrique ou multicentrique, la région, la source de financement et la déclaration des conflits d'intérêts) et l'effet traitement produit par la testostérone chez les hommes en utilisant une approche méta-épidémiologique. Les autres travaux explorent la qualité du reporting dans les études observationnelles, à travers une enquête sur les connaissances et l'utilisation de l'outil STROBE et l'utilisation d'un langage causal dans les

résumés d'études observationnelles. Le dernier chapitre est consacré à la discussion et à la conclusion de cette thèse.

2. Exploration des désaccords dans l'évaluation du risque de biais

2.1. Raisons de désaccords dans l'évaluation du risque de biais dans les essais inclus dans deux revues Cochrane

Les revues systématiques (RS) sont des synthèses systématiques de toutes les connaissances existantes concernant une question clinique pertinente suivant un processus rigoureux et reproductible. L'évaluation de la qualité méthodologique des études individuelles est une étape cruciale de la revue systématique, car si les études individuelles sont biaisées, la RS le sera également. Comme mentionné précédemment, la Cochrane a développé un outil pour évaluer le risque de biais dans les ECR (l'outil RoB) ; malgré son développement rigoureux et l'existence de recommandations sur la façon de l'utiliser, des études antérieures ont montré que la reproductibilité de l'outil n'était pas optimale. Les raisons pour lesquelles des désaccords se sont produits dans l'évaluation n'ont pas été étudiées en profondeur. Par exemple, on ne sait pas si ces désaccords sont liés à une interprétation différente ou à l'accès à des informations complémentaires en contactant les auteurs des études par exemple. Les deux premiers projets de la thèse se sont concentrés sur l'exploration des raisons de désaccord dans l'évaluation du risque de biais des ECR inclus dans deux revues Cochrane différentes. Le premier projet (Chapitre 2) visait à évaluer la fréquence des désaccords dans l'évaluation du risque de biais pour les ECR inclus dans deux revues Cochrane différents et à explorer les raisons de ces désaccords. Cette étude a été menée à l'aide d'un ensemble de données incluant toutes les revues Cochrane publiées entre mars 2011 et septembre 2014 dans lesquelles nous avons identifié les essais cliniques qui ont été évalués par l'outil RoB dans deux revues différentes. Ces essais ont été identifiés grâce à leur référence et nous avons vérifié manuellement qu'ils correspondaient bien à la même étude. L'évaluation RoB pour les cinq principaux items (génération de la séquence de randomisation, assignation secrète, mise en insu des participants et du personnel, mise en insu de l'évaluation des critères de jugement, données incomplètes) a été extraite pour chaque revue. Pour chaque item, nous avons comparé l'évaluation RoB effectuée dans chaque revue et calculé la proportion d'accord. L'analyse approfondie des désaccords était une analyse menée sur un échantillon aléatoire de 50% des essais. Deux évaluateurs ont indépendamment analysé les désaccords en comparant le support de chaque jugement avec le rapport d'étude en texte intégral pour évaluer si les désaccords étaient liés à une différence d'information (par exemple, contact avec l'auteur de l'étude ou accès au protocole) ou à

une différence d'interprétation (même support de jugement mais interprétation différente). Ils ont également identifié les principales raisons d'une interprétation différente.

Nous avons identifié 1604 ECR inclus dans deux revues différentes. La proportion d'accord entre les revues variait selon l'item de l'outil RoB. La proportion d'accord allait de 57% (770/1 348 essais) pour les données incomplètes à 81% pour la génération de la séquence de randomisation (1 193/1 466). L'analyse approfondie des désaccords a montré que la cause la plus fréquente de désaccord était une interprétation différente de la même information, allant de 65% (88/136) pour la génération de la séquence de randomisation à 90% (56/62) pour l'insu des participants et du personnel. Ces désaccords étaient généralement liés à des articles scientifiques incomplets ou insuffisamment clairs, ce qui semblait pousser les chercheurs à un jugement plus personnel et subjectif. Plus rarement, la raison du désaccord était l'accès à des informations supplémentaires telles que le protocole ou un contact direct avec les auteurs. Dans notre échantillon, l'utilisation d'informations complémentaires expliquait 32/136 (24%) des désaccords pour la génération de la séquence de randomisation et 38/205 (19%) pour l'assignation secrète.

Nos résultats confirment la reproductibilité imparfaite de l'évaluation du risque de biais retrouvé dans les études précédentes, les chercheurs étant d'accord 70% du temps sur le RoB. Nous avons exploré les raisons des désaccords et constaté que la raison la plus fréquente était liée à une interprétation différente d'une information peu claire ou incomplète dans l'article scientifique, ce qui semble pousser les chercheurs vers une interprétation plus personnelle. Nous avons identifié des situations fréquentes où l'information était incomplète dans les articles scientifiques, ce qui pourrait permettre de fournir des recommandations plus précises sur la façon d'évaluer le risque de biais dans ces situations.

2.2. Influence de facteurs extérieurs sur l'évaluation du risque de biais

Le deuxième projet (Chapitre 3) provient directement des résultats du premier. Nous avons montré que les chercheurs peuvent évaluer différemment les mêmes informations incomplètes. Nous nous sommes demandé si certains « facteurs contextuels » pouvaient influencer le risque de biais en cas d'information incomplète. Nous avons exploré des caractéristiques liées aux examinateurs, à la revue ou à l'étude elle-même, par rapport aux autres études incluses dans la revue. Notre idée était que les auteurs peuvent, par exemple, considérer les études qui sont de plus grande taille ou plus récentes différemment des études plus petites ou plus anciennes, ou que différents groupes de revue peuvent avoir des approches différentes pour traiter les informations incomplètes ou peu claires dans les études individuelles. Nous avons donc cherché à explorer si certains de ces facteurs sont

associés à un risque élevé ou faible de biais en cas d'information incomplète ou peu claire dans les rapports des études primaires pour lesquelles un risque de biais « incertain » aurait été attendu. Cette analyse était basée sur les 304 ECR pour lesquels nous avons considéré que le désaccord sur le risque de biais était lié à une interprétation différente d'une information incomplète ou peu claire dans l'article scientifique. Nous avons utilisé un plan expérimental de type cas-témoins pour répondre à cette question. Un cas a été défini comme un risque de biais élevé ou faible pour un ECR dans l'une des revues. Un témoin a été défini comme un risque de biais incertain pour le même ECR, pour le même item, dans l'autre revue. Les cas et les témoins ont été identifiés pour chaque item.

Nos résultats montrent que lorsque les auteurs de la revue étaient également auteurs de l'ECR évalué, ils étaient plus susceptibles d'évaluer un item à faible risque de biais qu'à risque incertain (OR 11,71 (IC à 95% 1,39-98,76)). Les essais plus anciens dans une revue étaient plus souvent considérés à faible risque de biais (OR 0,37 [0,15-0,96]). De plus, les groupes de revue et les auteurs qui avaient participé à un nombre inférieur de revues avaient tendance à attribuer plus souvent un risque de biais faible que les groupes de revue et auteurs plus expérimentés (OR 0,97 (IC à 95% 0,95 -0,99) pour les groupes de revue) et 0,97 (IC à 95% 0,95-0,998) pour les auteurs de la revue). Ceci pourrait suggérer que le niveau de formation peut différer selon les groupes de revue, ou que l'évaluation du risque de biais peut être influencée par l'expérience des évaluateurs. Nos résultats suggèrent que certains facteurs liés à la revue et à l'auteur peuvent jouer un rôle dans l'évaluation du risque de biais des ECR dans les revues systématiques. La connaissance de ces facteurs contextuels qui peuvent influencer sur les jugements de risque de biais peut aider les chercheurs à mener des évaluations plus reproductibles.

3. Association entre certaines caractéristiques méthodologiques et l'effet traitement : un exemple ciblé portant sur la testostérone chez l'homme

Nous avons mentionné précédemment que les items de l'outil RoB étaient des caractéristiques associées à l'effet traitement dans les études méta-épidémiologiques, qui ont été utilisées pour fournir des éléments de preuve empiriques pour le développement de l'outil.

Dans l'étude rapportée dans le Chapitre 4, nous avons cherché à explorer si d'autres facteurs étaient associés à l'effet traitement et nous nous sommes concentrés sur un sujet particulier, la thérapie par testostérone chez les hommes adultes.

Cette étude méta-épidémiologique a porté sur 132 essais inclus dans 19 SR et publiés entre 2008 et 2018. Aucune des caractéristiques étudiées, y compris l'année de publication, la taille de

l'échantillon, l'enregistrement de l'essai, le caractère mono ou multicentrique, la région, la source de financement, la déclaration des conflits d'intérêts étaient significativement associés à l'effet traitement pour cette question de recherche, probablement en raison du faible nombre d'essais examinés.

Les résultats, bien que non statistiquement significatifs, suggéraient que les essais avec un RoB élevé / incertain pour l'item insu des évaluateurs semblaient montrer des estimations d'effet plus importantes que les essais à RoB faible. De plus, les essais évalués à risque de biais global élevé ont rapporté des effets traitement plus importants que les essais avec un RoB global faible, mais ce résultat n'était pas non plus significatif (rapport des cotes : 0,79, intervalle de confiance à 95% : 0,60 à 1,03).

En conclusion, nous n'avons trouvé aucune preuve claire que l'une des caractéristiques des essais examinées était associée à l'effet traitement dans les essais randomisés évaluant la testostérone chez les hommes.

4. Qualité des rapports d'études observationnelles

4.1. Enquête sur la connaissance et l'utilisation de l'outil STROBE pour rapporter une étude observationnelle

L'une des approches pour améliorer la qualité des rapports d'études observationnelles consiste à utiliser un RG adapté au type d'étude. Les RG sont généralement organisés sous la forme d'une liste d'items, contenant les éléments essentiels devant être rapportés dans un article scientifique. Les auteurs ne connaissent pas toujours l'existence de RG pertinents et certains peuvent même être opposés à leur utilisation, car ils ont l'impression que les RG prennent du temps ou limitent leur liberté dans la rédaction de l'article. Les revues scientifiques n'exigent pas systématiquement l'utilisation des RG lors de la soumission d'un manuscrit. Le quatrième projet (Chapitre 5) a été conçu pour explorer la manière dont les chercheurs voient et interagissent avec le RG consacré aux études observationnelles, l'outil STROBE (STrengthening the Reporting of Observational Studies in Epidemiology) publié en 2007. Le but de l'étude était d'identifier les éléments facilitateurs ou les barrières et les motivations liés à l'utilisation de l'outil STROBE.

L'étude a été développée sous la forme d'une enquête en ligne qui a été envoyée aux auteurs d'études observationnelles. Le processus de recrutement des chercheurs pour l'enquête a utilisé différentes méthodes : invitation au sein du réseau professionnel, promotion de l'enquête via les réseaux sociaux, demande à 257 revues biomédicales d'inviter leurs auteurs à participer, contact direct avec les auteurs correspondants d'études observationnelles.

Un total de 1015 participants ont répondu à l'enquête. Parmi ceux-ci, 185 (18,2%) ont indiqué qu'ils n'avaient jamais entendu parler de l'outil STROBE ni ne l'avaient utilisé auparavant, 195 (19,2%) en avaient entendu parler mais ne l'avaient jamais utilisé et 635 (62,6%) l'avaient déjà utilisé. Lorsqu'on leur a demandé quels facteurs avaient ou auraient motivé l'utilisation de STROBE, le processus de soumission des revues et l'utilisation obligatoire des RG ont été les réponses les plus fréquentes suivies par la motivation personnelle. Environ la moitié (53%) des chercheurs qui connaissaient STROBE mais ne l'avaient pas utilisé ont déclaré que c'était parce que les revues ne le demandaient pas. La deuxième raison la plus fréquemment mentionnée était que leurs co-auteurs ne l'utilisaient pas (24,3%). Il est intéressant de noter que, bien que les revues aient été décrites comme le moyen le plus courant par lequel les participants ont initialement eu connaissance de STROBE, le fait que les revues ne l'exigent pas au moment de la soumission était la raison la plus fréquente pour laquelle les auteurs ne l'utilisaient pas. Les résultats de l'enquête ont également montré que les auteurs avaient bien perçu les avantages de l'utilisation d'un RG en termes d'amélioration de la qualité de leurs manuscrits et de chance d'être publié. Nos résultats suggèrent qu'une implémentation plus active de STROBE par les revues permettrait d'améliorer la sensibilisation des auteurs et son utilisation.

4.2. Utilisation d'allégations causales dans les résumés d'articles scientifiques décrivant une étude observationnelle

Dans l'étude rapportée dans le Chapitre 6, nous avons évalué si les chercheurs utilisaient un langage causal dans les résumés de publications d'études observationnelles et si cela était fait de manière cohérente dans le résumé et le manuscrit. Pour ce faire, nous avons recherché toutes les études observationnelles publiées dans le BMJ en 2018 et évalué leurs résumés pour rechercher un langage de type causal. Deux évaluateurs indépendants ont lu le résumé et évalué la cohérence des déclarations causales dans les différentes parties du résumé. Les résumés ont été classés comme « constamment causal », « inconstamment causal » ou « constamment non causal ». Nous avons considéré un résumé "inconstamment causal" si un langage causal était rapporté dans une section mais pas dans une autre. Des exemples de ce qui constitue un langage causal ont été recueillis. Nous avons en outre évalué le texte intégral publié pour évaluer les méthodes statistiques appliquées et rechercher toute autre allégation causale.

Pour les résumés classés comme « inconstamment causal », nous avons également évalué en double le texte intégral du manuscrit initialement soumis au BMJ pour évaluer si des changements étaient survenus au cours du processus d'examen par les pairs concernant l'utilisation du langage causal.

Sur les 151 articles de recherche publiés identifiés, 60 étaient éligibles et inclus dans cette étude. Parmi ceux-ci, 29 résumés (48%) ont été classés comme constamment causal. Douze autres résumés (20%) ont été jugés comme inconstamment causal, principalement parce que l'objectif énoncé évaluait une association, tandis que la conclusion présentait une conclusion causale (9/12) ou l'inverse (3/12). Enfin, il y avait des résumés qui ne rapportaient aucun langage causal, qui ont été considérés comme systématiquement non causaux ($n = 19$, 32%). Tous les articles classés comme « constamment causal » sur la base du résumé, utilisaient également un langage causal et contenaient des déclarations causales dans le texte intégral. C'était également le cas pour plus de la moitié (11/19) des résumés classés comme « inconstamment causal », où même si le résumé était formulé en termes d'association, les auteurs ont appliqué des méthodes causales, discuté le risque de confusion résiduelle, la plausibilité biologique ou une relation dose-réponse suggérant un lien causal.

Nos résultats suggèrent que des conseils supplémentaires pourraient aider les auteurs à identifier ce qui constitue une déclaration causale et comment la justifier et discuter les hypothèses sous-jacentes.

5. Discussion

Dans cette thèse, nous nous sommes concentrés sur deux causes importantes de gâchis dans la recherche : le risque de biais et le reporting incomplet ou inadéquat. Les deux premiers projets portaient sur l'évaluation du risque de biais dans les essais cliniques et sur la manière dont nous pouvons améliorer la reproductibilité de cette évaluation. Les autres projets se sont principalement concentrés sur la manière de rapporter les résultats de la recherche, en particulier sur les connaissances et l'utilisation du RG dédié aux études observationnelles par les chercheurs du domaine et l'utilisation d'un langage inapproprié ou ambigu concernant la causalité dans ces études. L'une des forces de cette thèse est la réalisation de différents projets en lien avec le risque de biais ou la qualité du reporting, différents types d'étude (schéma de type cas témoin, étude méta-épidémiologique, enquête en ligne, revue méthodologique) et l'implication d'un réseau pluridisciplinaire. En effet, nous avons travaillé en lien avec des membres de la Cochrane pour les projets en lien avec l'évaluation du risque de biais et avec des éditeurs de journaux notamment le BMJ pour les projets sur le reporting.

Les trois premiers projets se sont concentrés sur le risque de biais. Dans la première étude, nous avons évalué les discordances dans l'évaluation RoB dans une situation réelle en évaluant comment

les mêmes ECR ont été évalués pour le RoB dans deux revues Cochrane différentes. Notre étude était originale car les études précédentes sur la reproductibilité consistaient en une évaluation indépendante du RoB par deux ou plusieurs chercheurs de l'étude, ce qui n'était pas forcément représentatif de tous les auteurs Cochrane. Malgré cette différence de conception importante, nos résultats sont cohérents avec les études précédentes montrant que la reproductibilité du RoB était sous-optimale (la proportion d'accord était d'environ 70% globalement dans notre étude). Une autre originalité de notre étude a été l'exploration en profondeur des raisons potentielles de désaccords en évaluant le support du jugement et le texte intégral des études individuelles. Nous avons montré que l'interprétation différente d'une information incomplète ou peu claire dans le rapport d'étude était la principale cause de désaccord. La recherche d'informations complémentaires telles que le protocole ou un contact direct avec les auteurs était une autre cause de désaccord mais moins fréquente. Cette étude met en évidence l'importance d'améliorer la qualité des rapports pour les études individuelles. Dans la deuxième étude sur l'évaluation du risque de biais, tirée des résultats de la première étude, nous avons examiné la question de la reproductibilité sous un angle différent, en évaluant si certains facteurs contextuels pouvaient influencer l'évaluation RoB. Nous avons montré que certains facteurs peuvent avoir une influence, comme le fait d'avoir effectué plus de RS dans le passé ou d'être parmi les auteurs de l'étude évaluée. Nous pensons qu'il est important d'attirer l'attention des chercheurs sur l'existence de tels facteurs, qui ne font pas partie de l'outil RoB, car une meilleure connaissance de ceux-ci peut améliorer la reproductibilité de l'évaluation. À notre connaissance, aucune autre étude n'a évalué l'influence possible de tels facteurs contextuels.

L'étude méta-épidémiologique sur le traitement par la testostérone a donné des résultats intéressants. Dans ce sujet très restreint, nous n'avons trouvé aucune preuve que le RoB était associé à l'effet traitement. Cela était probablement dû à la petite taille de l'échantillon, mais aussi à une faible prévalence d'études à faible RoB dans notre échantillon.

Dans la dernière partie de la thèse, nous nous sommes concentrés sur le reporting à travers l'évaluation de la connaissance et de l'utilisation du RG dédié aux études observationnelles et l'utilisation d'un langage causal dans les études observationnelles. Concernant les RG, nos résultats suggèrent qu'une implémentation plus active par les revues est un élément clé pour améliorer leur utilisation. Les revues devraient recommander davantage l'utilisation des RG ou exiger leur utilisation avec une soumission de la liste d'items remplie avec le manuscrit.

Le dernier travail a évalué l'utilisation du langage causal dans les articles scientifiques rapportant une étude observationnelle, en se concentrant sur les allégations de causalité rapportées dans les résumés. Dans la conclusion, nous avons suggéré qu'un rôle plus actif des éditeurs et des pairs pourrait aider à éviter l'abus d'allégations causales dans les études observationnelles.

Néanmoins, les suggestions issues de ces travaux sur le reporting soulèvent d'importantes questions de faisabilité. Les améliorations suggérées prennent du temps et sont coûteuses et de nombreuses revues ne disposent pas des ressources nécessaires pour vérifier la soumission de la liste d'items du RG correspondant ou les aspects plus fins de sémantique liés au langage causal. Des données antérieures suggéraient que les revues ayant un facteur d'impact plus élevé étaient susceptibles d'avoir une meilleure qualité du reporting [54], ce qui peut être expliqué par davantage de ressources ou un engagement plus important dans l'amélioration de la qualité, mais aussi par le fait d'attirer généralement de « meilleures » recherches.

6. Implications

Le thème commun mis en évidence à travers les différentes études de ma thèse souligne l'importance de mener des recherches sur les méthodes pour améliorer la qualité de la littérature scientifique, au niveau de la planification, de la conduite et du reporting des études. Nous avons identifié plusieurs domaines d'intervention. Ces domaines concernent l'évaluation du risque de biais, la diffusion et l'acceptation des RG et la nécessité d'éviter les allégations causales inappropriées dans les études observationnelles.

Dans le premier projet sur l'évaluation du risque de biais, la présence d'informations incomplètes ou peu claires dans le rapport d'étude était la principale cause de désaccord dans l'évaluation du RoB. Nous avons donc souligné la nécessité d'améliorer les rapports, mais également la nécessité d'apporter des recommandations plus précises concernant l'évaluation des cas courants d'informations incomplètes. Un changement récent qui va dans cette direction est le nouvel outil lancé par la Cochrane en 2019 [55] et récemment mis à jour (novembre 2020) [56]. L'outil a été modifié pour adopter une approche pas à pas basée sur des « questions de signalisation » pour guider et faciliter l'évaluation. L'outil comprend également des algorithmes qui peuvent aider à améliorer la reproductibilité. Les recherches futures devront évaluer si cette mise à jour importante améliore la reproductibilité de l'évaluation du RoB.

Le deuxième projet sur l'évaluation du risque de biais a permis d'identifier des facteurs susceptibles d'influencer les évaluateurs du RoB. Les évaluateurs doivent être conscients que leur jugement, même s'il est méthodologiquement solide, peut être affecté ou biaisé par des facteurs ne faisant pas partie du processus. Ces facteurs peuvent ne pas être contrôlables par les évaluateurs, comme leur expérience dans l'évaluation du risque de biais, tandis que d'autres, comme le fait que l'évaluateur était aussi auteur de l'étude, sont plus évitables. Une suggestion, qui est également liée au premier

projet, est de se concentrer sur la formation des évaluateurs « moins expérimentés » à travers une approche plus pratique couvrant le problème que nous avons mis en évidence.

L'importance de la formation est également soulignée dans nos études sur le reporting. Dans notre projet sur un RG largement utilisé, les auteurs semblaient être conscients de son existence et de son importance, mais cette connaissance ne s'est pas nécessairement traduite par une mise en œuvre. De même, lorsque nous avons évalué l'utilisation de langage causal dans les études observationnelles, nous avons remarqué quelques utilisations abusives bien que nous soyons convaincus que les principes de causalité sont bien connus des auteurs.

Nos projets sur le risque de biais ont identifié des domaines d'intervention précis dans le domaine de la méthodologie de la recherche. Les projets de recherche sur le reporting, en revanche, mettent au premier plan les éditeurs de revues scientifiques. Les revues scientifiques ont été présentées comme des acteurs clés dans la promotion de l'utilisation des RG ; mais leur rôle devrait probablement s'élargir, même si nous sommes conscients que cela nécessite des ressources conséquentes que seules les grandes revues possèdent. Les revues dotées de ressources qui le permettent pourraient évoluer vers une implémentation plus active des RG, mais aussi avoir un rôle proactif en offrant des outils pour former et guider les auteurs dans l'utilisation des RG pertinents. De même, un effort rédactionnel important devrait être fait pour éviter l'utilisation d'allégations causales non étayées ; les rédacteurs en chef, axés sur la diffusion scientifique, devraient soutenir des initiatives visant à utiliser des formulations adaptées et à éviter celles qui sont trompeuses.

7. Conclusions

Ma thèse portait sur le thème du gâchis de la recherche en se concentrant sur différents moments du processus de recherche et différents types d'étude. Nous avons travaillé sur la méthodologie de la recherche et sur le reporting. Nos travaux ont impliqué des méthodologistes, des chercheurs d'études observationnelles et des éditeurs de revues. Nous avons identifié des aspects précis du processus d'évaluation du risque de biais qui peuvent être améliorés, tandis qu'une attention particulière doit être accordée aux facteurs extérieurs qui ne font pas partie de l'évaluation. En ce qui concerne le reporting, les revues doivent être conscientes de leur importance dans la mise en œuvre de ces recommandations pour améliorer la qualité du reporting et éviter les allégations causales inappropriées.

Summary

1. Introduction and objectives

Evidence based medicine (EBM) has introduced a new paradigm in both clinical practice and biomedical research. **(Chapter One)** EBM is defined as “the conscientious, explicit and judicious use of current best evidence in making decisions about the care of individual patients”[1]. The principle of EBM is that the foundation of any medical decisions should rely on the best scientific evidence from clinical research[2]. Although EBM integrates clinical experience and patient’s preference, it puts a major emphasis on the scientific literature. Unfortunately, the scientific literature quality is overall not optimal although there have been advances in methodology and in requirements by regulatory agencies[3]. It has been estimated that up to 85% of biomedical research could be wasted[4, 5]. Such waste can happen at every step of the research process from the planning to the reporting of the research. At the planning stage, researchers may formulate the wrong research question or use an inappropriate design. Waste can happen during the study conduct, using methods that are prone to induce bias in the results. Finally, there may be waste because of incomplete or inadequate reporting of results, including the use of wrong terminology[6-9]. In this thesis, we aimed to focus on two main causes of research waste: high risk of bias and poor quality of reporting.

Bias in epidemiology is defined as a systematic error, resulting in an underestimation or an overestimation of the true treatment effect [73]. The identification and measurement of bias is difficult. Paradoxically, a study with numerous flaws in its methodology may by chance be unbiased, while the contrary is also possible. For these reasons, it is considered more appropriate to focus on the risk of bias (RoB) to measure the likelihood of bias to happen. This translates in the assessment of key methodological features of a study, which could possibly lead to misleading results[10-13]. In 2005, Cochrane started to develop a tool to assess the RoB in Randomized Controlled Trials (RCTs) on the basis of empirical evidence from meta-epidemiological studies evaluating methodological characteristics associated with treatment effect estimates[13-18]. The RoB tool includes the following items: Random Sequence Generation and Allocation Concealment, referring to the risk of selection bias; Blinding of Participants and Personnel, referring to the risk of performance bias; Blinding of Outcome Assessment, referring to the risk of detection bias; Incomplete Outcome Data, referring to the risk of attrition bias; Selective Reporting, referring to the risk of reporting bias and Other bias. The RoB assessment of each item consists of two parts: a brief description of the support for judgement which usually includes verbatim quotes from the publications on the study evaluated and the judgment concerning risk of bias. The risk can be

expressed as low risk of bias, high risk of bias and unclear risk of bias if there were insufficient information to reach a judgment[13] and Cochrane provides recommendations for assessing the RoB for each item. The RoB tool has quickly become the reference tool for assessing RoB and is nowadays used in all Cochrane reviews and has also been increasingly common among non-Cochrane reviews [19]. However, several studies have found an overall low reproducibility of the assessment conducted through the RoB tool, especially for some items such as Incomplete Outcome Data and the blinding items [10, 20-25]. None of these studies explored the reasons that may explain such low agreement on RoB assessment.

Another important problem affecting the scientific literature is related to the poor quality of reporting in research studies. Study reporting should be clear, complete and easily available; if important information of a study is poorly or not reported this usually makes the replication, critical appraisal and interpretation of results difficult or impossible. Poor reporting therefore limits the RoB assessment with items generally evaluated as unclear in that case when using the RoB tool. With inadequate reporting, results also cannot properly inform clinical practice or health policies, contributing to the overall problem of waste in research[26-28]. Numerous studies have shown the poor quality of reporting across various topics and study types[15, 29-35]. To address this issue, the research community has developed structured guidance to scientific writing with the creation of tools such as reporting guidelines (RG) [4, 36, 37]. The first structured efforts to improve the quality of reporting were aimed toward RCTs and can be traced back to 1993 when the Standardized Reporting of Trials (SORT) Statement was created [38, 39]. Simultaneously and independently, a different group was also working on providing guidance for reporting. So, in 1996, the SORT group, in collaboration with the Asilomar Working Group, published the Consolidated Standards of Reporting Trials (CONSORT) Statement[40-42] which is considered one of the first structured RG. Although initially conceived for RCTs, CONSORT has inspired many offshoots, known as extensions, which provide specific guidance for different study designs, interventions, or data types[43].

RGs are often organised as checklists, containing essential items that should be reported in a publication. These checklists may also be accompanied by a flow diagram, detailing different information, such as how many people were initially contacted, how many declined to participate, and the number of dropouts. The reporting guideline is frequently associated with an “Elaboration and Explanation” document, providing a detailed description of the use of the checklist and giving supporting background for the tool. One of the most used RGs in the field of observational research is the STrengthening the Reporting of OBservational studies in Epidemiology (STROBE) Statement, created in 2007[44]. STROBE is a 22-item checklist that details the key information

needed when reporting the results of an observational study. STROBE also provides further details and guidance for each checklist item giving examples of good reporting[45]. Thank to this explanatory document, STROBE has been used by researchers as training material, both for the use of STROBE, but also to guide the methodology of research [46]. Although STROBE has been widely used by researches for more than 10 years, there is lack of evidence regarding the knowledge and the perception of researchers.

Another cause of inadequate reporting is related to the use of inappropriate terminology that may mislead the readers. It is generally considered that RCT is the only design able to draw causal inference[47, 48] because randomization allows comparable distribution of confounding factors between experimental and control groups[49]. RCTs however cannot be performed in every occasion, therefore, despite the limitations of observational research, authors may be tempted to draw causal conclusions in observational studies[50, 51]. To address the issue of correct reporting of non-causal results, researchers have proposed alternative ways to present observational research correctly. One suggestion is to discuss the subject in more details, including in the report an open disclosure regarding the differences between the different study designs and the implication of choosing one design over another [52, 53].

The thesis aims to explore different issues related to RoB assessment and inadequate reporting in clinical research. It consists of five different projects; two of them focus on exploration of reasons of disagreements when assessing RoB for the same trials included in two different systematic reviews (SRs). A third project evaluates the association between RCT characteristics including RoB items (but also year of publication, sample size, trial registration status, center status, region, funding source and conflict of interest) and the reported treatment effects of testosterone therapy in men using a meta-epidemiological approach. The other works explore the quality of reporting in observational studies, through the examination of the knowledge and adherence to reporting guidelines and the use of appropriate wording regarding causal claims in abstracts of observational studies.

2. Exploration of disagreements when assessing RoB in RCTs

2.1. Reasons of disagreements in risk of bias assessment for RCTs assessed in two different Cochrane reviews

Systematic reviews (SRs) are systematic synthesis of all the existing knowledge concerning a clinical relevant question following a rigorous and reproducible process. Assessing the methodological quality of individual studies is a crucial step when conducting a systematic review

because if the individual studies are biased, the SR will be also biased. As previously mentioned, Cochrane developed a tool to assess RoB in RCTs (the RoB tool); despite its rigorous development and the existence of guidelines on how to use it, previous studies showed that the reproducibility of assessments is often suboptimal. The reasons why disagreements in the assessment happened have not been explored in depth. For example, reviewers may base their judgment on additional information such as protocols or they may contact study authors, which may explain the low agreement observed. The first two projects in the thesis focused on exploring the reasons of disagreements in the RoB assessment of RCTs included in two different Cochrane reviews. The first project (**Chapter Two**) aimed to assess the frequency and explore the reasons for disagreements in RoB assessment for RCTs included in two different Cochrane SR. This study was conducted using a dataset of all Cochrane SRs published between March 2011 and September 2014 in which we identified clinical trials that were assessed for RoB in two different SRs. Trials included in two different reviews were identified through their primary reference and we manually checked that it corresponded to the same study. RoB assessment for five different items (Random Sequence Generation, Allocation Concealment, Blinding of Participants and Personnel, Blinding of Outcome Assessment, Incomplete Outcome Data) was extracted for each review. For each item, we compared RoB assessment made in each review and calculated the proportion of agreement. The in-depth analysis of disagreements was a second level analysis conducted on a random sample of 50% of eligible RCTs. Two reviewers independently analyzed the disagreements by comparing support for each judgement with information from the full-text study report to evaluate whether disagreements were related to a difference in information (e.g., contact with the study author) or a difference in interpretation (same support for judgement but different interpretation). They also identified main reasons for different interpretation.

We identified 1604 RCTs included in two different reviews. The agreement in the assessment varied between the different items of the RoB tool. The proportion of agreement ranged from 57% (770/1,348 trials) for incomplete outcome data to 81% for random sequence generation (1,193/1,466). The in-depth analysis of disagreements showed that the most common source of disagreement was a difference in interpretation of the same information, ranging from 65% (88/136) for random sequence generation to 90% (56/62) for blinding of participants and personnel. These disagreements were usually related to incomplete or unclear reporting in the study report, which seemed to push the researchers into more personal and subjective judgement. More rarely the reason of disagreement was access to different or additional information such as protocols or contact with authors. In our sample, the use of different information explained 32/136 (24%) of the disagreements for random sequence generation and 38/205 (19%) for allocation concealment.

Our results confirm the suboptimal agreement on the risk of bias assessment found in previous studies with researchers agreeing 70% of the times on RoB. We explored the reasons of disagreements and found that the most common reason of disagreement was related to a different interpretation of an unclear or incomplete information reported in scientific articles which seems to push the researchers into using much personal judgement. We identified common cases of incomplete information in scientific articles as this gives the opportunity to provide more detailed guidance on how to assess the risk of bias in these cases.

2.2. Influence of external factors on RoB assessment

The second project (**Chapter 3**) originated directly from the results of the first one. We showed that researchers, may evaluate differently the same incomplete information. We wondered whether some “contextual factors”, might influence the risk of bias judgements in case of incomplete information. We explored characteristics related to the reviewers, the review, or the study itself, when compared to the other studies included in the review. Our idea was that authors may, for example, consider studies that are larger or newer differently than smaller or older studies, or different review groups may have different approaches for handling incomplete or unclear information in individual studies. We therefore aimed to explore which of these factors, if any, are associated with a high or low risk of bias judgement in case of incomplete or unclear information in reports of primary studies for which an “unclear” risk of bias judgment would have been expected.

This analysis was based on the 304 RCTs for which we considered that disagreement on the risk of bias judgement was related to different interpretation of an incomplete or unclear information in the study report. We used a matched case-control design. A case was defined as a judgement of “high risk of bias” or “low risk of bias” for an RCT in one of the reviews. A matched control was defined as a judgement of “unclear risk of bias” for the same RCT, for the same item, in the second, different review. Cases and controls were identified for each item.

Our results showed that when review authors were also authors of the RCT assessed, they were more likely to assess an item at low risk of bias than unclear (OR 11.71 (95% CI 1.39-98.76). Older trials in a review were more often assigned a low risk of bias (OR 0.37 [0.15-0.96]). Also review groups and authors that had completed a lower number of reviews slightly more often assigned a low risk whereas others reported “unclear” (OR 0.97 (95% CI 0.95-0.99) for review groups) and 0.97 (95% CI 0.95-0.998) for review authors). This finding could suggest that the level of training may differ across author groups, or it may be influenced by the experience of the reviewers.

Our results suggest that some review and author factors may play a role in risk of bias judgements of RCTs in systematic reviews. The awareness of contextual characteristics that can affect risk of bias judgements may help researchers in conducting more reproducible assessments of risk of bias.

3. Association between methodological characteristics and treatment effect: a case example on testosterone therapy in men

We previously mentioned that the items of the RoB tool were characteristics associated with treatment effect in meta-epidemiological studies, which were used to provide empirical evidence for the development of the RoB tool.

In the study reported in **Chapter 4**, we aimed to explore whether other factors were associated with treatment effect and to focus on a particular topic, testosterone therapy in adult men.

This meta-epidemiological study involved 132 trials on testosterone therapy included in 19 different SRs and published between 2008 and 2018. None of the investigated design characteristics, including year of publication, sample size, trial registration status, centre status, region, funding, source, and conflict of interest were significantly associated with reported treatment effects of testosterone therapy in men, probably because of the low number of trials examined.

The results, although statistically not significant suggested that trials with high/unclear RoB due to blinding of outcome assessors seemed to show larger effect estimates than trials rated at low RoB. Moreover, trials rated at high overall risk of bias reported larger treatment effects than trials rated at low overall RoB but this result did not meet statistical significance (ratio of odds ratio: 0.79, 95% confidence interval: 0.60 to 1.03).

In conclusion, we found no clear evidence that any of the trial characteristics examined were associated with treatment effects in randomized trials of testosterone therapy in men.

4. Quality of reporting in observational studies

4.1. A survey on the use and knowledge of STROBE

In the fourth project (**Chapter 5**) we worked on issues regarding the reporting of observational studies. One of the approaches to improve the quality of reporting is to provide more structure and guidance to scientific writing using RGs. RG are commonly organized in the form of a checklist, containing essential items that need to be reported when reporting the study. Authors are often not aware of the existence of relevant RGs or they may even be against using one, considering that RG may take time or limit their freedom in writing a paper. Scientific journals do not systematically

require respecting RGs when submitting a manuscript. The fourth project was designed to explore how researchers view and interact with the reporting guideline dedicated to observational studies (e.g., cross-sectional, cohort, case-control), the STrengthening the Reporting of Observational studies in Epidemiology (STROBE) Statement published in 2007. The aim of the study was to identify the personal and environmental facilitators, barriers, and motivators to using the STROBE Statement.

The study was developed as an online survey which was sent to authors of observational studies. The process of recruiting researchers for the survey used different methods: authors among the professional network, promotion of the survey through social media, asking 257 biomedical journals to invite their authors to participate, direct contact of corresponding authors of observational studies.

A total of 1015 participants completed the survey. Of these, 185 (18.2%) indicated they had never heard of STROBE nor used it previously, 195 (19.2%) had heard of it but never used it, and 635 (62.6%) had used it. When asked about what factors would or have motivated use of STROBE, the journal submission process and mandatory use of reporting guidelines were the most frequent answers followed by self-motivation. Around half (53%) of researchers who were aware of STROBE but had not used it reported that this was because journals did not require it. The next most frequently reported reason was that their co-authors did not use it (24.3%). Interestingly although journals were described as the most typical medium by which participants originally became aware of STROBE, the fact that journals did not require to adhere to STROBE was the most common reason why authors did not use it.

The results of the survey showed that, despite time costs, authors perceived benefits of using an RG such as increasing the quality of their manuscripts and the chances of being published. Our results suggest that the endorsement of STROBE by journals would be important to improve authors' awareness and use of the guideline.

4.2. Use of causal claims in observational studies

In the study reported in **Chapter 6** we addressed the fact that researchers may tend to draw causal conclusions in publications when dealing with observational data. We explored whether and how statements in study reports of observational studies present confusing use of causal language. To do this we searched for all observational studies published in the BMJ in 2018 and evaluated their abstracts to identify causal language. Two independent reviewers read the abstract and evaluated the consistency of causal statements across the different parts of the abstract structure. The abstracts

were classified as “consistently causal”, “inconsistent” or “consistently non causal”. We considered an abstract “inconsistent” if for example causal claiming was made in one section but not in another. Examples of what constitutes causal language were collected. We further evaluated the full published text to highlight the statistical methods applied and any further causal claims. For the abstracts classified as inconsistent, we evaluated in duplicate the full-text of the published version and the manuscript initially submitted to the BMJ to highlight whether changes occurred during the peer review process regarding use of causal language.

From the 151 published research articles identified, 60 were eligible and included in this study. Among these, 29 studies (48%) were classified as using causal language. A further twelve (20%) abstracts were considered inconsistent mainly because the objective stated evaluating an association while the conclusion presented a causal finding (9/12) or the opposite (3/12). Finally, there were abstracts that did not report any causal language, which were considered as consistently not causal (n=19, 32%). All papers classified as ‘consistently causal’ based on the abstract, also used causal language and contained causal statements in the full text. This was additionally the case with more than half (11/19) of the abstracts classified as ‘consistently not causal’, where even though the abstract was carefully phrased in terms of association, the authors applied causal methods, discussed residual confounding, biological plausibility or a dose-response relationship suggesting a causal aim.

Our results suggest that further guidance could help authors in identifying what constitutes a causal statement and how is possible to justify or to discuss assumptions involved.

5. Discussion

In this thesis, we focused on two main causes of waste in research: risk of bias and incomplete or inadequate reporting. The core projects dealt with the assessment of risk of bias in clinical trials and how we may improve reproducibility of this assessment. The other projects mainly focused on reporting especially attitudes of researchers regarding reporting guidelines and use of inappropriate or ambiguous language in observational studies.

Dealing with different moments and aspects of the research process allowed the thesis to be multidisciplinary. The aspect of multidisciplinary was held both in the people involvement and in the restitution of results to them. Our first project involved research methodologists and provided them areas of attention for the assessment of RoB; the other projects directly involved researchers

of observational studies and journal editors and provided them hints to better understand practical issues in the reporting.

The first three projects focused on risk of bias. In the first study we assessed the agreement in the RoB assessment in a real-life situation by evaluated how the same RCTs were assessed for RoB in two different Cochrane reviews. Our study was innovative because previous studies on reproducibility consisted in an independent evaluation of the RoB by two or more researchers from the study, who may not be representative of all Cochrane reviewers. Despite this important difference in design, our results are consistent with previous studies showing that agreement between reviewers was suboptimal (overall agreement was about 70% in our study). Another originality of our study was the in-depth exploration of potential reasons of disagreements by evaluating the support for judgement and the full text of individual studies if necessary. We showed that different interpretation of an incomplete or unclear information in the study report was the main cause of disagreement. Search for additional information such as the protocol or contact with authors was another cause of disagreement but less frequent. This study highlights the importance of improving the quality of reporting in individual studies although this was not the aim of our study. In the second study on RoB assessment, steamed from the results of the first study, we looked at the issue of the agreement from a different angle, evaluating whether some contextual factors may influence RoB assessment. We showed that some factors may have an influence, such as the fact of having conducted more SRs in the past or being among the authors of the study under scrutiny. We think that it is important to draw attention to the existence of such factors, not part of the RoB tool, because better knowledge of them may improve reproducibility of the assessment. To our knowledge, no other study evaluated the possible influence of such contextual factors.

The meta-epidemiological study on the testosterone therapy replacement, had interesting results. In that very narrow topic, we found no evidence that RoB was associated with treatment effect estimates. This was probably due to the small sample size, but also a low prevalence of studies at low RoB in our sample.

In the last part of the thesis, we focused on reporting through the assessment of the knowledge about reporting guidelines and the use of inappropriate language in observational studies.

Concerning reporting guidelines, our results suggest that a more active endorsement by journals is a key element to improve the use of reporting guidelines. Journals could consider recommending the use of RG or require their use with a jointed submission of the filled checklist with the main manuscript.

The final work of the thesis evaluated the use of inadequate use of causal language when reporting an observational study, focusing on the improper claims of causality reported in abstracts. In the

conclusion we suggested that a more active role of editors and reviewers could help avoiding the abuse of causal claims in observational studies.

Nevertheless, the suggestions coming from these works on reporting raise important feasibility issues. The improvements suggested are time consuming and expensive and many journals do not have the resources to verify submission of the corresponding reporting guideline checklist or check the finer aspects of semantics relative to causal language. Previous evidence suggested that journals with higher impact factor may have a better overall quality of reporting[54], which may be due to having more resources or being more engaged in quality improvement but also generally attract “better” research.

6. Implications

The common theme highlighted across the different studies of my thesis is the importance of conducting research on methods to improve the quality of research methodology, at the level of planning, conducting and reporting. We identified several areas of intervention. These areas concern some aspects and passage of RoB assessment, the dissemination and acceptance of RGs and the need for research on how to avoid improper causal claiming while reporting observational research.

In the first project on RoB assessment, dealing with incomplete or unclear information in the study report was the main cause of disagreement in RoB assessment. We therefore underlined the need to improve the reporting but also the need for better guidance regarding the assessment of common cases of incomplete information. One recent change that can help in this direction is the new tool launched by Cochrane in 2019[55] and recently updated (November 2020)[56]. The tool changes some aspects of the RoB assessment procedure and has adopted a step by step approach based on “signalling questions” to guide and facilitate the assessment. The tool also includes algorithms that could help improving the reproducibility. Future research should evaluate whether this important update improves the reproducibility of RoB assessment.

The second project on RoB hinted the existence of factors that may influence reviewers in the RoB assessment. The project showed how reviewers should be aware that their judgement, even if methodologically sound, could be affected or biased by factors not part of the process. Such factors may not be controllable by the reviewers such as their experience in conducting reviews, while others, such as the fact the reviewer also authored the study under scrutiny, are more avoidable. A

suggestion, that also links with the first project, is to focus on the training of “less experienced” reviewers through a more practical approach covering the common issue we highlighted. The importance of guidance and formation is also affirmed in our studies on reporting. In our project on a widely used reporting guideline, authors seemed to be aware of its existence and importance, but this knowledge did not necessarily translate into action. Similarly, when looking for causal claims in observational research, we notice frequent slips although we are confident that the principles of causality are well known by authors. Our projects on RoB identified precise areas of intervention in the field of research methodology. The projects on reporting, on the other hand, put at the centre of the stage editors of scientific journals. Scientific journals were shown as key actors in promoting the use of RGs; but their role should probably expand, even if we realize this will require additional resources and only the larger journals would be able to act such changes. Journals with additional resources could move toward demanding the use of RGs, but also have a more proactive role offering tools to form and guide authors in the use of relevant RGs. Similarly, a strong editorial effort should be made to avoid the use of unbacked causal claims; editors, focusing on science dissemination, should support and host initiatives toward the use of the right wording in science.

7. Conclusions

My thesis dealt with the topic of research waste focusing on different moments in the research process and different figures. We worked on methodology of research conduct and reporting, our work involved methodologists, researchers of observational studies and journal editors. We identified precise aspects of the RoB assessment process that are passible of improvement, while attention to factors that are not part of the assessment should also be given. Regarding reporting, journals should be aware of their importance in the implementation of reporting guidelines and for guidance in the avoidance of improper causal claims.

Samenvatting

1. Inleiding en Doel

Evidence based medicine (EBM) heeft een nieuw paradigma geïntroduceerd, zowel in de klinische praktijk als in het biomedisch onderzoek. EBM wordt gedefinieerd als "het gewetensvolle, expliciete en oordeelkundige gebruik van de beste, beschikbare onderbouwing bij het nemen van beslissingen over de zorg voor individuele patiënten"[1] (**Hoofdstuk 1**). Het principe van EBM is dat de basis van alle medische beslissingen moet berusten op de resultaten uit het beste klinisch-wetenschappelijk onderzoek[2].

Hoewel EBM klinische ervaring en de voorkeur van de patiënt integreert, legt het een grote nadruk op de wetenschappelijke literatuur. Helaas is de kwaliteit van de wetenschappelijke literatuur over het algemeen niet optimaal, hoewel er vooruitgang is geboekt in de methodologie en in de eisen van regelgevende instanties[3]. Er wordt geschat dat tot 85% van het biomedisch onderzoek verspild wordt[4, 5]. Dergelijke verspilling kan zich voordoen in elke stap van het onderzoeksproces, van de planning tot de rapportering van het onderzoek. In de planningsfase kunnen onderzoekers de verkeerde onderzoeksvraag formuleren of een ongeschikte opzet gebruiken. Verspilling kan zich voordoen tijdens de uitvoering van het onderzoek, door het gebruik van methoden die een vertekend beeld van de resultaten kunnen geven. Tenslotte kan er verspilling optreden door onvolledige of inadequate rapportage van resultaten, inclusief het gebruik van verkeerde terminologie[6-9]. In dit proefschrift hebben wij ons gericht op twee hoofdoorzaken van verspilling in onderzoek: een hoog risico op vertekening en slechte kwaliteit van de rapportage.

Bias wordt in de epidemiologie gedefinieerd als een systematische vertekening, resulterend in een onderschatting of overschatting van het ware behandelingseffect [73]. Het identificeren en schatten van bias is moeilijk. Paradoxaal genoeg kan een studie met talrijke methodologische gebreken toevallig vrij van vertekening zijn, terwijl ook het tegendeel mogelijk is. Om deze redenen wordt het zinvoller geacht zich te concentreren op het risico op vertekening (risk of bias - RoB) om de kans op vertekening te beoordelen. Dit vertaalt zich in de beoordeling van belangrijke methodologische kenmerken van een studie, die mogelijk tot misleidende resultaten kunnen leiden[10-13].

In 2005 is Cochrane begonnen met het ontwikkelen van een instrument om de RoB in Randomized Controlled Trials (RCT's) te beoordelen op basis van empirisch bewijs uit meta-epidemiologische studies, die methodologische kenmerken evalueren samenhangend met schattingen van het behandelingseffect[13-18]. De RoB-tool omvat de volgende onderdelen: Random Sequence Generation and Allocation Concealment, verwijzend naar het risico van selectiebias; Blinding van

deelnemers en personeel, verwijzend naar het risico van performance bias; Blinding van Outcome Assessment, verwijzend naar het risico van detectiebias; Incomplete Outcome Data, verwijzend naar het risico van attrition bias; Selective Reporting, verwijzend naar het risico van reporting bias en Other bias. De RoB-beoordeling van elk item bestaat uit twee delen: een korte beschrijving van de ondersteuning van het oordeel, die gewoonlijk letterlijke citaten uit de publicaties over de beoordeelde studie omvat, en het oordeel over het risico van vertekening. Het risico kan worden uitgedrukt als een laag risico op vertekening, hoog risico op vertekening en onduidelijk risico op vertekening, indien er onvoldoende informatie was om tot een oordeel te komen[13]. Cochrane geeft aanbevelingen voor de beoordeling van de RoB voor elk item.

De RoB-tool is snel uitgegroeid tot het referentie-instrument voor de beoordeling van de RoB en wordt tegenwoordig gebruikt in alle Cochrane-reviews; het instrument wordt ook steeds vaker gebruikt in niet-Cochrane-reviews [19]. Verschillende studies hebben echter een lage reproduceerbaarheid van de beoordeling met de RoB-tool gevonden, vooral voor sommige items zoals Incomplete Outcome Data en de blinding items [10, 20-25]. Geen van deze studies onderzocht de redenen die een dergelijke lage overeenstemming over de RoB beoordeling kunnen verklaren.

Een ander belangrijk probleem in de wetenschappelijke literatuur heeft te maken met de slechte kwaliteit van de rapportage van onderzoek. De rapportage van studies moet duidelijk, volledig en gemakkelijk beschikbaar zijn. Indien belangrijke informatie van een studie slecht of niet wordt gerapporteerd, maakt dit gewoonlijk de replicatie, kritische beoordeling en interpretatie van de resultaten moeilijk of onmogelijk. Een gebrekkige rapportage beperkt dan ook de RoB-beoordeling; in dat geval worden de items bij gebruik van het RoB-instrument doorgaans als onduidelijk beoordeeld. Door een ontoereikende rapportage kunnen de resultaten ook geen goede informatie verschaffen voor de klinische praktijk of het gezondheidsbeleid, wat bijdraagt tot het algemene probleem van verspilling in onderzoek[26-28]. Talrijke studies hebben de slechte kwaliteit van de rapportage over verschillende onderwerpen en studietypen aangetoond[15, 29-35]. Om dit probleem aan te pakken, heeft de onderzoeksgemeenschap gestructureerde richtlijnen voor het schrijven van wetenschappelijke artikelen ontwikkeld (*reporting guidelines* - RG) [4, 36, 37].

De eerste gestructureerde inspanningen om de kwaliteit van de rapportage te verbeteren waren gericht op RCT's en kunnen worden teruggevoerd tot 1993, toen de Standardized Reporting of Trials (SORT) Statement werd opgesteld [38, 39]. Gelijktijdig en onafhankelijk daarvan, werkte een andere groep ook aan het verschaffen van richtlijnen voor rapportage. Zo publiceerde de SORT groep, in samenwerking met de Asilomar Werkgroep, in 1996 de Consolidated Standards of Reporting Trials (CONSORT) Verklaring [40-42] die wordt beschouwd als één van de eerste

gestructureerde RG. Hoewel oorspronkelijk ontworpen voor RCT's, heeft CONSORT vele varianten geïnspireerd, bekend als extensies, die specifieke richtlijnen geven voor verschillende studie ontwerpen, interventies, of datatypes[43].

RG's zijn vaak georganiseerd als checklists, die essentiële items bevatten die in een publicatie moeten worden gerapporteerd. Deze checklists kunnen ook vergezeld gaan van een stroomdiagram, waarin verschillende gegevens worden gedetailleerd, zoals hoeveel mensen aanvankelijk werden gecontacteerd, hoeveel mensen afzagen van deelname, en het aantal uitvallers. De rapportagerichtlijn gaat vaak vergezeld van een document "Uitwerking en toelichting", waarin het gebruik van de checklist gedetailleerd wordt beschreven en ondersteunende achtergrondinformatie voor het hulpmiddel wordt gegeven.

Een van de meest gebruikte RG's op het gebied van observationeel onderzoek is de STrengthening the Reporting of OBservational studies in Epidemiology (STROBE) Statement, opgesteld in 2007[44]. STROBE is een 22-item checklist die de belangrijkste informatie beschrijft die nodig is bij het rapporteren van de resultaten van een observationele studie. STROBE geeft ook verdere details en richtlijnen voor elk item van de checklist met voorbeelden van goede rapportage[45]. Dankzij dit verklarend document is STROBE door onderzoekers gebruikt als trainingsmateriaal, zowel voor het gebruik van STROBE, maar ook om de methodologie van onderzoek te begeleiden [46]. Hoewel STROBE al meer dan 10 jaar op grote schaal door onderzoekers wordt gebruikt, is er een gebrek aan bewijs met betrekking tot de kennis en de perceptie van onderzoekers.

Een andere oorzaak van inadequate rapportage houdt verband met het gebruik van ongeschikte terminologie, die de lezers kan misleiden. Algemeen wordt aangenomen dat RCT's het enige ontwerp zijn dat in staat is causale conclusies te trekken[47, 48] omdat randomisatie een vergelijkbare verdeling van versturende factoren tussen experimentele en controlegroepen mogelijk maakt[49]. RCT's kunnen echter niet in alle gevallen worden uitgevoerd, daarom kunnen auteurs, ondanks de beperkingen van observationeel onderzoek, in de verleiding komen om causale conclusies te trekken in observationele studies[50, 51]. Om het probleem van de correcte rapportering van niet-causale resultaten aan te pakken, hebben onderzoekers alternatieve manieren voorgesteld om observationeel onderzoek correct te presenteren. Eén suggestie is om het onderwerp meer in detail te bespreken, door in het rapport een open uiteenzetting op te nemen over de verschillen tussen de verschillende studieontwerpen en de implicatie van het kiezen van het ene ontwerp boven het andere [52, 53].

Het in dit proefschrift gerapporteerde onderzoek had tot doel verschillende kwesties te onderzoeken die verband houden met de beoordeling van RoB en inadequate rapportage in klinisch onderzoek. Het bestaat uit vijf verschillende projecten; twee daarvan richten zich op de verkenning van de

redenen van onenigheid bij de beoordeling van RoB voor dezelfde trials die zijn opgenomen in twee verschillende systematische reviews (SRs). Een derde project evalueert het verband tussen RCT-kenmerken met inbegrip van RoB-items (maar ook jaar van publicatie, steekproefgrootte, proefregistratiestatus, centrumstatus, regio, financieringsbron en belangenconflict) en de gerapporteerde behandelingseffecten van testosterontherapie bij mannen met behulp van een meta-epidemiologische benadering. De andere projecten onderzoeken de kwaliteit van de rapportage in observationele studies, door de kennis en de naleving van de rapportagerichtlijnen en het gebruik van passende bewoordingen met betrekking tot causale beweringen in samenvattingen van observationele studies te onderzoeken.

2. Verkenning van discrepanties bij de beoordeling van Risk of Bias in RCT's

2.1. Redenen voor discrepanties bij de beoordeling van risico op vertekening voor RCT's die in twee verschillende Cochrane-reviews zijn beoordeeld

Systematisch literatuuronderzoek (*Systematische reviews* - SR) bieden een systematisch samengestelde synthese van bestaande kennis over een klinisch relevante vraagstelling, volgens een rigoreus en reproduceerbaar proces. Het beoordelen van de methodologische kwaliteit van individuele studies is een cruciale stap bij het uitvoeren van een systematische review, want als de individuele studies vertekend zijn, zal de SR ook vertekend zijn. Zoals reeds eerder vermeld, heeft Cochrane een instrument ontwikkeld om de reproduceerbaarheid van RCT's te beoordelen (het RoB-instrument); ondanks de zorgvuldige ontwikkeling van dit instrument en het bestaan van richtsnoeren voor het gebruik ervan, is uit eerdere studies gebleken dat de reproduceerbaarheid van de beoordelingen vaak niet optimaal is. De redenen waarom er discrepanties bij de beoordeling zijn, zijn niet grondig onderzocht. Beoordelaars kunnen hun oordeel bijvoorbeeld baseren op aanvullende informatie zoals protocollen of zij kunnen contact opnemen met studieauteurs, wat de geconstateerde lage overeenstemming kan verklaren.

De eerste twee projecten in dit proefschrift waren gericht op het onderzoeken van de redenen van discrepanties in de RoB-beoordeling van RCT's opgenomen in twee verschillende Cochrane reviews. Het eerste project (**Hoofdstuk 2**) had tot doel de frequentie van en de redenen voor discrepanties in de RoB beoordeling voor RCT's opgenomen in twee verschillende Cochrane SR te onderzoeken. Deze studie werd uitgevoerd met behulp van een dataset van alle Cochrane SR's gepubliceerd tussen maart 2011 en september 2014 waarin we klinische trials identificeerden die werden beoordeeld op RoB in twee verschillende SR's. Trials opgenomen in twee verschillende

reviews werden geïdentificeerd via hun primaire referentie; we controleerden handmatig of deze overeenkwam met dezelfde studie.

De RoB-beoordeling voor vijf verschillende items (Random Sequence Generation, Allocation Concealment, Blinding of Participants and Personnel, Blinding of Outcome Assessment, Incomplete Outcome Data) werd geëxtraheerd voor elke review. Voor elk item vergeleken we de RoB-beoordeling in elke review en berekenden we de mate van overeenstemming. De grondige evaluatie van de verschillen van mening was een analyse van het tweede niveau, uitgevoerd op een willekeurige steekproef van 50% van de in aanmerking komende RCT's. Twee beoordelaars analyseerden onafhankelijk van elkaar de discrepanties, door de ondersteuning voor elk oordeel te vergelijken met informatie uit het full-text onderzoeksverslag, om na te gaan of de discrepanties verband hielden met een verschil in informatie (bijv. contact met de auteur) of met een verschil in interpretatie (dezelfde ondersteuning voor het oordeel, maar een andere interpretatie). Zij identificeerden ook de belangrijkste redenen voor verschillende interpretatie.

Wij identificeerden 1604 RCT's die in twee verschillende reviews waren opgenomen. De overeenstemming in de beoordeling varieerde tussen de verschillende items van het RoB-instrument. De mate van overeenstemming varieerde van 57% (770/1.348 trials) voor onvolledige uitkomstgegevens tot 81% voor het genereren van willekeurige reeksen (1.193/1.466). De diepgaande analyse van de discrepanties toonde aan dat de meest voorkomende bron van onenigheid een verschil in interpretatie van dezelfde informatie was, variërend van 65% (88/136) voor random sequence generation tot 90% (56/62) voor blinding van deelnemers en personeel. Deze discrepanties hielden meestal verband met onvolledige of onduidelijke rapportage in het onderzoeksverslag, waardoor de onderzoekers leken over te gaan tot een meer persoonlijk en subjectief oordeel. In zeldzamere gevallen was de reden van onenigheid de toegang tot andere of aanvullende informatie, zoals protocollen of contact met de auteurs. In onze steekproef verklaarde het gebruik van verschillende informatie 32/136 (24%) van de discrepanties voor random sequence generation en 38/205 (19%) voor allocation concealment.

Onze resultaten bevestigen de suboptimale overeenstemming over de beoordeling van het risico van vertekening die in eerdere studies werd gevonden, waarbij onderzoekers het in 70% van de gevallen eens waren over RoB. We onderzochten de redenen van discrepanties en ontdekten dat de meest voorkomende reden van verschil van mening te maken had met een verschillende interpretatie van onduidelijke of onvolledige informatie zoals die in wetenschappelijke artikels was terug te vinden, wat de onderzoekers ertoe lijkt aan te zetten om veel persoonlijke oordelen te gebruiken. We hebben veel voorkomende gevallen van onvolledige informatie in wetenschappelijke artikelen

geïdentificeerd omdat dit de mogelijkheid biedt om meer gedetailleerde richtlijnen op te stellen over hoe het risico op vertekening in deze gevallen moet worden beoordeeld.

2.2. Invloed van externe factoren op de RoB-beoordeling

Het tweede project (gerapporteerd in **Hoofdstuk 3**) vloeide rechtstreeks voort uit de resultaten van het eerste. We toonden aan dat onderzoekers dezelfde onvolledige informatie verschillend kunnen beoordelen. We vroegen ons af of sommige "contextuele factoren", de beoordeling van het risico op vertekening zouden kunnen beïnvloeden in geval van onvolledige informatie. We onderzochten kenmerken die verband hielden met de beoordelaars, de beoordeling, of de studie zelf, in vergelijking met de andere studies die in de beoordeling waren opgenomen. Onze hypothese was dat auteurs bijvoorbeeld studies die groter of nieuwer zijn anders kunnen beoordelen dan kleinere of oudere studies, of dat verschillende beoordelingsgroepen verschillende benaderingen kunnen hebben voor het omgaan met onvolledige of onduidelijke informatie in individuele studies. Daarom wilden wij nagaan welke van deze factoren eventueel geassocieerd zijn met een hoge of lage beoordeling van het risico op vertekening in geval van onvolledige of onduidelijke informatie in verslagen van primaire studies waarvoor een "onduidelijk" oordeel over het risico op vertekening zou zijn verwacht.

Deze analyse was gebaseerd op de 304 RCT's waarvoor wij aannamen dat onenigheid over de beoordeling van het risico op vertekening verband hield met een verschillende interpretatie van onvolledige of onduidelijke informatie in het onderzoeksverslag. Wij gebruikten een gematched case-control design. Een geval werd gedefinieerd als een beoordeling van "hoog risico op vertekening" of "laag risico op vertekening" voor een RCT in een van de reviews. Een gemaachte controle werd gedefinieerd als een beoordeling van "onduidelijk risico op vertekening" voor dezelfde RCT, voor hetzelfde item, in de tweede, andere review. Gevallen en controles werden voor elk item geïdentificeerd.

Onze resultaten toonden aan dat wanneer auteurs van het review ook auteur waren van de beoordeelde RCT, zij vaker een item beoordeelden met een laag risico op bias dan als onduidelijk (OR 11,71 (95% CI 1,39-98,76)). Oudere trials in een review kregen vaker een laag risico op vertekening (OR 0,37 [0,15-0,96]). Ook reviewgroepen en auteurs die een lager aantal reviews hadden voltooid, kenden iets vaker een laag risico toe, terwijl anderen "onduidelijk" rapporteerden (OR 0,97 (95% CI 0,95-0,99) voor reviewgroepen) en 0,97 (95% CI 0,95-0,998) voor reviewauteurs). Deze bevinding zou kunnen suggereren dat het opleidingsniveau verschilt tussen de auteursgroepen, of dat het beïnvloed wordt door de ervaring van de reviewers.

Onze resultaten suggereren dat sommige beoordelings- en auteursfactoren een rol kunnen spelen bij de beoordeling van het risico op vertekening van RCT's in systematische reviews. De bewustwording van contextuele kenmerken die de beoordeling van het risico op vertekening kunnen beïnvloeden, kan onderzoekers helpen bij het uitvoeren van meer reproduceerbare beoordelingen van het risico op vertekening.

3. Associatie tussen methodologische kenmerken en behandel-effect: een voorbeeld over testosterontherapie bij mannen

We beschreven eerder dat de items van het RoB-instrument kenmerken waren die geassocieerd waren met behandel-effect in meta-epidemiologische studies, die werden gebruikt om empirisch bewijs te leveren voor de ontwikkeling van het RoB-instrument.

In de studie die wordt beschreven in **Hoofdstuk 4** wilden we onderzoeken of andere factoren geassocieerd waren met behandel-effect en wilden we ons richten op een specifiek onderwerp, testosterontherapie bij volwassen mannen.

Deze meta-epidemiologische studie omvatte 132 trials over testosterontherapie, opgenomen in 19 verschillende SR's en gepubliceerd tussen 2008 en 2018. Geen van de onderzochte designkenmerken, waaronder jaar van publicatie, steekproefgrootte, proefregistratiestatus, centrumstatus, regio, financiering, bron, en belangenverstrengeling, waren significant geassocieerd met de gerapporteerde behandelingseffecten van testosterontherapie bij mannen, waarschijnlijk vanwege het lage aantal onderzochte trials.

Deze resultaten, hoewel statistisch niet significant, suggereerden dat trials met een hoge/onduidelijke RoB als gevolg van blinding van de uitkomstbeoordelaars grotere effectschattingen laten zien dan trials met een lage RoB. Bovendien rapporteerden trials met een hoog geheel risico op vertekening grotere behandel-effecten dan trials met een laag geheel risico op vertekening, maar dit resultaat was niet statistisch significant (odds ratio: 0,79, 95% betrouwbaarheidsinterval: 0,60 tot 1,03).

Concluderend vonden we geen duidelijk bewijs dat een van de onderzochte trialkarakteristieken geassocieerd was met behandelingseffecten in gerandomiseerde trials van testosterontherapie bij mannen.

4. Kwaliteit van de rapportage in observationele studies

4.1. Een enquête over het gebruik en de kennis van STROBE

In het vierde project, gerapporteerd in **Hoofdstuk 5**, hebben we gewerkt aan kwesties betreffende de rapportage van observationele studies. Een van de benaderingen om de kwaliteit van de rapportage te verbeteren is het bieden van meer structuur en begeleiding bij het schrijven van wetenschappelijke artikelen met behulp van *reporting guidelines* (RG). RG's worden gewoonlijk gepresenteerd in de vorm van een checklist, met essentiële items die gerapporteerd moeten worden bij de rapportage van de studie. Auteurs zijn vaak niet op de hoogte van het bestaan van relevante RG's of zijn zelfs tegen het gebruik ervan, omdat RG's tijd kunnen kosten of hun vrijheid bij het schrijven van een artikel kunnen beperken. Wetenschappelijke tijdschriften schrijven niet systematisch voor dat RG's moeten worden gerespecteerd bij het indienen van een manuscript. Het vierde project was bedoeld om na te gaan hoe onderzoekers aankijken tegen en omgaan met de rapporteringsrichtlijn voor observationele studies (bv. cross-sectioneel, cohort, case-control), de STrengthening the Reporting of Observational studies in Epidemiology (STROBE)-verklaring die in 2007 werd gepubliceerd. Het doel van de studie was om de persoonlijke en omgevingsfacilitatoren, barrières en motivatoren voor het gebruik van STROBE te identificeren. De studie werd ontwikkeld als een online enquête die naar auteurs van observationele studies werd gestuurd. Voor het rekruteren van onderzoekers voor de enquête werden verschillende methoden gebruikt: auteurs uit het professionele netwerk, promotie van de enquête via sociale media, 257 biomedische tijdschriften vragen hun auteurs uit te nodigen om deel te nemen, direct contact met corresponderende auteurs van observationele studies.

In totaal hebben 1.015 deelnemers de enquête ingevuld. Hiervan hadden 185 (18,2%) nog nooit van STROBE gehoord of het eerder gebruikt, 195 (19,2%) hadden er wel van gehoord maar het nooit gebruikt, en 635 (62,6%) hadden het gebruikt. Op de vraag welke factoren het gebruik van STROBE zouden of hebben bevorderd, waren de indieningsprocedure bij tijdschriften en het verplichte gebruik van rapportagerichtlijnen de meest frequente antwoorden, gevolgd door zelfmotivatie. Ongeveer de helft (53%) van de onderzoekers die op de hoogte waren van STROBE maar er geen gebruik van hadden gemaakt, meldden dat dit zo was omdat de tijdschriften het niet verplicht stelden. De volgende reden die het vaakst werd vermeld, was dat hun co-auteurs het niet gebruikten (24,3%). Interessant is dat, hoewel tijdschriften werden beschreven als het meest typische medium waardoor deelnemers aanvankelijk bewust werden van STROBE, het feit dat tijdschriften niet eisten dat zij zich ook daadwerkelijk aan STROBE hielden de meest voorkomende reden was waarom auteurs het instrument niet gebruikten.

De resultaten van de enquête toonden aan dat, ondanks de tijdskosten, auteurs voordelen zagen in het gebruik van een RG, zoals het verhogen van de kwaliteit van hun manuscripten en de kans om gepubliceerd te worden. Onze resultaten suggereren dat de bekrachtiging van STROBE door tijdschriften belangrijk zou zijn om de bekendheid en het gebruik van de richtlijn door auteurs te verbeteren.

4.2. Gebruik van causale beweringen in observationele studies

In de studie waarvan in **Hoofdstuk 6** verslag wordt uitgebracht hebben we bestudeerd of onderzoekers de neiging hebben om causale conclusies te trekken in publicaties over observationele gegevens. We onderzochten of en hoe uitspraken in onderzoeksrapporten van observationele studies verwarrend gebruik maken van causale uitdrukkingen. Om dit te doen zochten we naar alle observationele studies gepubliceerd in BMJ in 2018 en evalueerden we hun abstracts om causaal taalgebruik te identificeren.

Twee onafhankelijke beoordelaars lazen het abstract en evalueerden de consistentie van causale verklaringen doorheen de verschillende onderdelen van het abstract. De abstracts werden geclassificeerd als "consistent causaal", "inconsistent" of "consistent niet causaal". We beschouwden een samenvatting als "inconsistent" als er bijvoorbeeld in één onderdeel wel een causale bewering werd gedaan, maar in een ander onderdeel niet. We verzamelden voorbeelden van wat causaal taalgebruik is. We evalueerden verder de volledige gepubliceerde tekst om de toegepaste statistische methoden en eventuele verdere causale beweringen te benadrukken. Voor de samenvattingen die als inconsistent werden geclassificeerd, evalueerden we in tweevoud de volledige tekst van de gepubliceerde versie en het manuscript dat aanvankelijk bij BMJ was ingediend om te benadrukken of er tijdens het peer review proces veranderingen waren opgetreden met betrekking tot het gebruik van causaal taalgebruik.

Van de 151 gepubliceerde artikelen die werden geïdentificeerd, kwamen er 60 in aanmerking en werden opgenomen in deze studie. Daarvan werden 29 studies (48%) geclassificeerd als studies met causaal taalgebruik. Twaalf andere samenvattingen (20%) werden als inconsistent beschouwd, voornamelijk omdat in de doelstelling een associatie werd geëvalueerd terwijl in de conclusie een causale bevinding werd gepresenteerd (9/12) of het tegenovergestelde (3/12). Ten slotte waren er samenvattingen waarin geen enkel causaal verband werd vermeld, die werden beschouwd als consistent niet causaal (n=19, 32%). Alle papers die op basis van het abstract als 'consistent causaal' werden geclassificeerd, gebruikten ook causaal taalgebruik en bevatten causale verklaringen in de volledige tekst. Dit was ook het geval bij meer dan de helft (11/19) van de samenvattingen die als

'consistent niet causaal' werden geclassificeerd, waarbij de auteurs, hoewel de samenvatting zorgvuldig was geformuleerd in termen van associaties, causale methoden toepasten, residuele confounding bespraken, biologische plausibiliteit of een dosis-respons relatie die een causaal doel suggereerde.

Onze resultaten suggereren dat verdere begeleiding auteurs zou kunnen helpen bij het identificeren van wat een causale verklaring is en hoe het mogelijk is om de betrokken veronderstellingen te rechtvaardigen of te bespreken.

5. Discussie

In dit proefschrift hebben we ons geconcentreerd op twee belangrijke oorzaken van verspilling in onderzoek: risico op vertekening en onvolledige of inadequate rapportage. De sleutelprojecten bestudeerden de beoordeling van het risico op vertekening in klinisch onderzoek en hoe we de reproduceerbaarheid van deze beoordeling kunnen verbeteren. De andere projecten richtten zich vooral op verslaglegging, met name de houding van onderzoekers ten opzichte van verslagleggingsrichtlijnen en het gebruik van ongepast of dubbelzinnig taalgebruik in observationele studies.

Door verschillende momenten en aspecten van het onderzoeksproces te behandelen, kon dit proefschrift multidisciplinair zijn. Het multidisciplinaire aspect was zowel aanwezig in de betrokkenheid van de mensen als in de teruggave van de resultaten aan hen. Ons eerste project betrof methodologen en gaf hen aandachtspunten voor de beoordeling van RoB; de andere projecten betroffen rechtstreeks onderzoekers van observationele studies en tijdschriftredacteurs en gaven hen hints om praktische kwesties in de verslaglegging beter te begrijpen.

De eerste drie projecten waren gericht op het risico van vertekening. In de eerste studie hebben we de overeenstemming in de RoB beoordeling in een real-life situatie beoordeeld door te evalueren hoe dezelfde RCT's werden beoordeeld op RoB in twee verschillende Cochrane reviews. Onze studie zou als vernieuwend kunnen worden beschouwd, omdat eerdere studies naar reproduceerbaarheid bestonden uit een onafhankelijke beoordeling van de RoB door twee of meer onderzoekers uit de studie, die mogelijk niet representatief zijn voor alle Cochrane reviewers.

Ondanks dit belangrijke verschil in opzet zijn onze resultaten consistent met eerdere studies die aantonen dat de overeenstemming tussen reviewers suboptimaal was (de totale overeenstemming was ongeveer 70% in onze studie). Een andere origineel punt van onze studie was de diepgaande exploratie van mogelijke redenen van meningsverschillen, door het evalueren van de ondersteuning

van het oordeel en de volledige tekst van individuele studies, indien nodig. Wij toonden aan dat een verschil in interpretatie van onvolledige of onduidelijke informatie in het onderzoeksverslag de belangrijkste oorzaak van discrepanties was. Zoeken naar aanvullende informatie zoals het protocol of contact met de auteurs was een andere oorzaak van onenigheid, maar minder frequent. Deze studie benadrukt het belang van de verbetering van de kwaliteit van de rapportage in individuele studies, hoewel dit niet het doel van onze studie was.

In de tweede studie over RoB-beoordeling, geïnspireerd door de resultaten van de eerste studie, bekeken we de kwestie van de overeenstemming vanuit een andere invalshoek, door te evalueren of sommige contextuele factoren de RoB-beoordeling kunnen beïnvloeden. Wij toonden aan dat sommige factoren een invloed kunnen hebben, zoals het feit dat men in het verleden meer SR's heeft uitgevoerd of dat men tot de auteurs van de onderzochte studie behoort. Wij denken dat het belangrijk is de aandacht te vestigen op het bestaan van dergelijke factoren, die geen deel uitmaken van het RoB-instrument, omdat een betere kennis ervan de reproduceerbaarheid van de beoordeling kan verbeteren. Voor zover wij weten, heeft geen enkele andere studie de mogelijke invloed van dergelijke contextuele factoren geëvalueerd.

De meta-epidemiologische studie over de vervanging van testosterontherapie had o.i. interessante resultaten. We vonden geen bewijs dat RoB geassocieerd was met de effectschattingen van de behandeling. Dit was waarschijnlijk te wijten aan de kleine steekproefgrootte, maar ook een lage prevalentie van studies met een lage RoB in onze steekproef.

In het laatste deel van het proefschrift richtten we ons op rapportage door de kennis over rapportagerichtlijnen en het gebruik van ongepast taalgebruik in observationele studies te beoordelen. Wat de rapportagerichtlijnen betreft, suggereren onze resultaten dat een actievere ondersteuning door tijdschriften een sleutelement is om het gebruik van rapportagerichtlijnen te verbeteren. Tijdschriften zouden kunnen overwegen het gebruik van de RG aan te bevelen of het gebruik ervan verplicht te stellen door de ingevulde checklist samen met het hoofdmanuscript in te dienen.

Het laatste deel in het proefschrift brengt verslag uit van een evaluatie van het gebruik van inadequaat causaal taalgebruik bij de rapportage van een observationele studie, waarbij de nadruk lag op onjuiste beweringen over causaliteit zoals die in samenvattingen werden gerapporteerd. In de conclusie stelden we voor dat een actievere rol van redacteurs en reviewers het misbruik van causale beweringen in observationele studies zou kunnen helpen vermijden. Toch werpen deze suggesties ook vragen op over de haalbaarheid. De voorgestelde verbeteringen zijn vaak tijdrovend en duur en veel tijdschriften beschikken niet over de middelen om na te gaan of de overeenkomstige checklist uit de rapportagerichtlijn wordt nageleefd, of om de fijnere aspecten van de semantiek met

betrekking tot causaal taalgebruik te controleren. In het verleden is gebleken dat tijdschriften met een hogere impactfactor een betere algemene kwaliteit van de verslaggeving hebben[54], wat te wijten kan zijn aan het feit dat ze over meer middelen beschikken of meer betrokken zijn bij kwaliteitsverbetering, maar over het algemeen ook "beter" onderzoek aantrekken.

6. Implicaties

Een gemeenschappelijke thema voor de verschillende studies in dit proefschrift was en is het belang van onderzoek naar methoden om de kwaliteit van onderzoeksmethodologie te verbeteren, op het niveau van planning, uitvoering en rapportage. We hebben verschillende gebieden voor verbeteringen geïdentificeerd. Deze gebieden betreffen enkele aspecten en het verloop van de RoB-beoordeling, de verspreiding en acceptatie van richtlijnen voor de rapportage en de noodzaak van onderzoek naar hoe onjuiste causale beweringen kunnen worden vermeden bij de rapportage van observationeel onderzoek.

In het eerste project over RoB-beoordeling was het omgaan met onvolledige of onduidelijke informatie in het onderzoeksverslag de belangrijkste oorzaak van discrepanties bij de RoB-beoordeling. Daarom onderstreepten wij de noodzaak om de rapportage in de primaire studies te verbeteren, maar ook de behoefte aan betere begeleiding bij de beoordeling van veel voorkomende gevallen van onvolledige informatie. Een recente verandering kan helpen: de nieuwe tool die Cochrane in 2019 heeft gelanceerd[55] en die onlangs is bijgewerkt (november 2020)[56]. De tool verandert enkele aspecten van de RoB-beoordelingsprocedure en heeft een stapsgewijze aanpak aangenomen op basis van "signaleringsvragen" om de beoordeling te ondersteunen en te vergemakkelijken. Het instrument bevat ook algoritmen die de reproduceerbaarheid zouden kunnen helpen verbeteren. Toekomstig onderzoek moet uitwijzen of deze belangrijke update de reproduceerbaarheid van de RoB-beoordeling verbetert.

Het tweede project over RoB hintte op het bestaan van factoren die beoordelaars kunnen beïnvloeden bij de RoB-beoordeling. Het project toonde aan hoe beoordelaars zich ervan bewust moeten zijn dat hun oordeel, ook al is het methodologisch verantwoord, kan worden beïnvloed of vertekend door factoren die geen deel uitmaken van het proces. Dergelijke factoren zijn wellicht niet beheersbaar voor de beoordelaars, zoals hun ervaring met het uitvoeren van beoordelingen, terwijl andere, zoals het feit dat de beoordelaar ook auteur is van de onderzochte studie, beter vermijdbaar zijn. Een suggestie, die ook verband houdt met het eerste project, is zich te richten op

de opleiding van "minder ervaren" beoordelaars door middel van een meer praktische aanpak van de gemeenschappelijke kwestie die wij hebben belicht.

Het belang van begeleiding en vorming wordt ook bevestigd in onze studies over verslaglegging. In ons project over een veelgebruikte rapportagerichtlijn leken de auteurs zich bewust van het bestaan en het belang ervan, maar deze kennis werd niet noodzakelijk omgezet in actie. Ook bij het zoeken naar causale beweringen in observationeel onderzoek zien we vaak uitglijders, hoewel we ervan overtuigd zijn dat de causaliteitsbeginselen bij de auteurs goed bekend zijn.

In onze projecten over de onderzoeksmethode hebben wij precies vastgesteld op welke gebieden wij moeten ingrijpen. De projecten inzake rapportage daarentegen plaatsten redacteurs van wetenschappelijke tijdschriften in het middelpunt van de belangstelling. Wetenschappelijke tijdschriften bleken een sleutelrol te spelen bij de bevordering van het gebruik van RG's; maar hun rol moet waarschijnlijk worden uitgebreid, ook al beseffen we dat dit extra middelen zal vergen en dat alleen de grotere tijdschriften in staat zouden zijn om dergelijke veranderingen door te voeren. Tijdschriften die over extra middelen beschikken, zouden het gebruik van RG's kunnen vereisen, maar ook een meer proactieve rol kunnen spelen, door tools aan te bieden om auteurs te scholen en te begeleiden bij het gebruik van relevante RG's. Evenzo zou een krachtige redactionele inspanning moeten worden geleverd om het gebruik van ongefundeerde causale beweringen te vermijden; redacteurs die zich richten op de verspreiding van wetenschap zouden initiatieven voor het gebruik van de juiste bewoordingen in de wetenschap moeten steunen en ondersteunen.

7. Conclusies

Het in dit proefschrift beschreven onderzoek gaat uiteindelijk over *Waste in Research*, waarbij de nadruk lag op verschillende fasen in het onderzoeksproces. We richtten ons op de methodologie en op de rapportage. Bij de projecten waren methodologen, onderzoekers en tijdschriftredacteurs betrokken. Wij hebben aspecten van het beoordelingsproces van de *risk of bias* vastgesteld die voor verbetering vatbaar zijn. Wat de rapportering betreft, moeten tijdschriften zich bewust zijn van hun rol bij de toepassing van de richtlijnen voor rapportage en het vermijden van ongepaste causale beweringen.

Brief bibliography

1. Sackett, D.L., et al., *Evidence based medicine: what it is and what it isn't*. BMJ, 1996. **312**(7023): p. 71-2.
2. Eddy, D.M., *Clinical decision making: from theory to practice. Practice policies--guidelines for methods*. JAMA, 1990. **263**(13): p. 1839-41.
3. Bastian, H., P. Glasziou, and I. Chalmers, *Seventy-five trials and eleven systematic reviews a day: how will we ever keep up?* PLoS Med, 2010. **7**(9): p. e1000326.
4. Chalmers, I. and P. Glasziou, *Avoidable waste in the production and reporting of research evidence*. Lancet, 2009. **374**(9683): p. 86-9.
5. Macleod, M.R., et al., *Biomedical research: increasing value, reducing waste*. Lancet, 2014. **383**(9912): p. 101-4.
6. Ioannidis, J.P.A., et al., *Increasing value and reducing waste in research design, conduct, and analysis*. The Lancet, 2014. **383**(9912): p. 166-175.
7. Chalmers, I., et al., *How to increase value and reduce waste when research priorities are set*. The Lancet, 2014. **383**(9912): p. 156-165.
8. Chan, A.W., et al., *Increasing value and reducing waste: addressing inaccessible research*. Lancet, 2014. **383**(9913): p. 257-66.
9. Al-Shahi Salman, R., et al., *Increasing value and reducing waste in biomedical research regulation and management*. Lancet, 2014. **383**(9912): p. 176-85.
10. Hartling, L., et al., *Applying the risk of bias tool in a systematic review of combination long-acting beta-agonists and inhaled corticosteroids for persistent asthma*. PLoS One, 2011. **6**(2): p. e17242.
11. Hrobjartsson, A., et al., *Assessing risk of bias in randomised clinical trials included in Cochrane Reviews: the why is easy, the how is a challenge*. Cochrane Database Syst Rev, 2013(4): p. ED000058.
12. Hrobjartsson, A., et al., *Assessing Risk of Bias in Randomised Clinical Trials Included in Cochrane Reviews: The why is Easy, the how is a Challenge [editorial]*. Cochrane Database Syst Rev, 2013. **5**: p. ED000058.
13. Higgins JPT, Altman DG, Sterne JAC (editors). Chapter 8: Assessing risk of bias in included studies., in *Cochrane Handbook for Systematic Reviews of Interventions*, J.P. Higgins and S. Green, Editors., The Cochrane Collaboration, 2011. Available from www.handbook.cochrane.org.
14. Schulz, K.F., et al., *Empirical evidence of bias. Dimensions of methodological quality associated with estimates of treatment effects in controlled trials*. JAMA, 1995. **273**(5): p. 408-12.
15. Moher, D., et al., *Does quality of reports of randomised trials affect estimates of intervention efficacy reported in meta-analyses?* The Lancet, 1998. **352**(9128): p. 609-613.
16. Balk, E.M., et al., *Correlation of quality measures with estimates of treatment effect in meta-analyses of randomized controlled trials*. JAMA, 2002. **287**(22): p. 2973-82.
17. Page, M.J., et al., *Empirical Evidence of Study Design Biases in Randomized Trials: Systematic Review of Meta-Epidemiological Studies*. PLoS One, 2016. **11**(7): p. e0159267.
18. Dechartres, A., et al., *Empirical evaluation of which trial characteristics are associated with treatment effect estimates*. J Clin Epidemiol, 2016. **77**: p. 24-37.
19. Higgins, J.P.T., S. Green, and editors, *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]*. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org
20. Hartling, L., et al., *Testing the risk of bias tool showed low reliability between individual reviewers and across consensus assessments of reviewer pairs*. J Clin Epidemiol, 2013. **66**(9): p. 973-81.
21. Hartling, L., et al., *Risk of bias versus quality assessment of randomised controlled trials: cross sectional study*. BMJ, 2009. **339**(oct19 1): p. b4012.
22. Armijo-Olivo, S., et al., *Poor reliability between Cochrane reviewers and blinded external reviewers when applying the Cochrane risk of bias tool in physical therapy trials*. PLoS One, 2014. **9**(5): p. e96920.

23. Armijo-Olivo, S., et al., *Assessment of study quality for systematic reviews: a comparison of the Cochrane Collaboration Risk of Bias Tool and the Effective Public Health Practice Project Quality Assessment Tool: methodological research*. J Eval Clin Pract, 2012. **18**(1): p. 12-8.
24. Jordan, V.M., S.F. Lensen, and C.M. Farquhar, *There were large discrepancies in risk of bias tool judgments when a randomized controlled trial appeared in more than one systematic review*. J Clin Epidemiol, 2016.
25. da Costa, B.R., et al., *Effect of standardized training on the reliability of the Cochrane risk of bias assessment tool: a prospective study*. Syst Rev, 2017. **6**(1): p. 44.
26. Altman, D.G., D. Moher, and K.F. Schulz, *Improving the reporting of randomised trials: the CONSORT Statement and beyond*. Stat Med, 2012. **31**(25): p. 2985-97.
27. Begley, C.G. and J.P. Ioannidis, *Reproducibility in science: improving the standard for basic and preclinical research*. Circ Res, 2015. **116**(1): p. 116-26.
28. Glasziou, P., et al., *Reducing waste from incomplete or unusable reports of biomedical research*. The Lancet, 2014. **383**(9913): p. 267-276.
29. Mhaskar, R., et al., *Published methodological quality of randomized controlled trials does not reflect the actual quality assessed in protocols*. J Clin Epidemiol, 2012. **65**(6): p. 602-9.
30. Soares, H.P., et al., *Bad reporting does not mean bad methods for randomised trials: observational study of randomised controlled trials performed by the Radiation Therapy Oncology Group*. BMJ, 2004. **328**(7430): p. 22-4.
31. Altman, D.G., *Making research articles fit for purpose: structured reporting of key methods and findings*. Trials, 2015. **16**: p. 53.
32. Dechartres, A., et al., *Reviews assessing the quality or the reporting of randomized controlled trials are increasing over time but raised questions about how quality is assessed*. J Clin Epidemiol, 2011. **64**(2): p. 136-44.
33. Agha, R.A., et al., *Randomised controlled trials in plastic surgery: a systematic review of reporting quality*. Eur J Plast Surg, 2014. **37**: p. 55-62.
34. Bryant, J., et al., *A systematic review of the quality of reporting in published smoking cessation trials for pregnant women: an explanation for the evidence-practice gap?* Implement Sci, 2014. **9**: p. 94.
35. Chen, B., et al., *A retrospective survey of quality of reporting on randomized controlled trials of metformin for polycystic ovary syndrome*. Trials, 2014. **15**: p. 128.
36. Altman, D.G. and I. Simera, *A history of the evolution of guidelines for reporting medical research: the long road to the EQUATOR Network*. J R Soc Med, 2016. **109**(2): p. 67-77.
37. Chan, A.W., et al., *Empirical evidence for selective reporting of outcomes in randomized trials: comparison of protocols to published articles*. JAMA, 2004. **291**(20): p. 2457-65.
38. *A proposal for structured reporting of randomized controlled trials. The Standards of Reporting Trials Group*. JAMA, 1994. **272**(24): p. 1926-31.
39. Kendall, J.M., *Designing a research project: randomised controlled trials and their principles*. Emerg Med J, 2003. **20**(2): p. 164-8.
40. Begg, C., et al., *Improving the quality of reporting of randomized controlled trials. The CONSORT statement*. JAMA, 1996. **276**(8): p. 637-9.
41. *Call for comments on a proposal to improve reporting of clinical trials in the biomedical literature. Working Group on Recommendations for Reporting of Clinical Trials in the Biomedical Literature*. Ann Intern Med, 1994. **121**(11): p. 894-5.
42. Schulz, K.F., D.G. Altman, and D. Moher, *CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials*. PLoS Med, 2010. **7**(3): p. e1000251.
43. 20/11/2020]; Available from: <http://www.consort-statement.org/extensions>.
44. von Elm, E., et al., *The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies*. Epidemiology, 2007. **18**(6): p. 800-4.
45. Vandembroucke, J.P., et al., *Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): explanation and elaboration*. Epidemiology, 2007. **18**(6): p. 805-35.

46. da Costa, B.R., et al., *Uses and misuses of the STROBE statement: bibliographic study*. BMJ Open, 2011. **1**(1): p. e000048.
47. Cofield, S.S., R.V. Corona, and D.B. Allison, *Use of causal language in observational studies of obesity and nutrition*. Obes Facts, 2010. **3**(6): p. 353-6.
48. Thapa, D.K., et al., *Being honest with causal language in writing for publication*. J Adv Nurs, 2020. **76**(6): p. 1285-1288.
49. Kunz, R. and A.D. Oxman, *The unpredictability paradox: review of empirical comparisons of randomised and non-randomised clinical trials*. BMJ, 1998. **317**(7167): p. 1185-90.
50. Begg, M.D. and D. March, *Cause and Association: Missing the Forest for the Trees*. Am J Public Health, 2018. **108**(5): p. 620.
51. Hernan, M.A., *A definition of causal effect for epidemiological research*. J Epidemiol Community Health, 2004. **58**(4): p. 265-71.
52. Green, M.J., *Calculating Versus Estimating Causal Effects*. Am J Public Health, 2018. **108**(8): p. e4-e5.
53. Goetghebuer, E., et al., *Formulating causal questions and principled statistical answers*. Stat Med, 2020.
54. Dechartres, A., et al., *Evolution of poor reporting and inadequate methods over time in 20 920 randomised controlled trials included in Cochrane reviews: research on research study*. BMJ, 2017. **357**: p. j2490.
55. Sterne, J.A.C., et al., *RoB 2: a revised tool for assessing risk of bias in randomised trials*. BMJ, 2019. **366**: p. l4898.
56. *RoB 2.0 Tool*. 22/08/2018]; Available from: <https://sites.google.com/site/riskofbiastool/welcome/rob-2-0-tool>.

Acknowledgements

I would like to express my deep and sincere gratitude to my supervisor, Agnès Dechartres, for giving me the right advice on my work and providing me with guidance and structure despite our differences in views and my life choices, some of which complicated our collaboration.

My gratitude also goes to my co-supervisor Patrick Bossuyt for his meaningful insights not only on my thesis, but also on my professional life and general attitude.

I am grateful to Philippe Ravaud and Isabelle Boutron for having introduced me to the MiRoR project and for their valuable advice at various stages of my work on the PhD.

I also thank the members of the jury, Joerg Meerphol, Emilie Sbidian, Matthieu Resche-Rigon, Marcel Dijkgraaf, Marc Besselink and Lotty Hooft for sharing their comments on my thesis and allowing me to complete this journey and, therefore, join the scientific community.

I am thankful to the other authors and co-authors of the works included in this thesis: Ignacio Atal, Robin Haring, Matthew Page, Roser Rius, Elizabeth Wager, Guadalupe Gomez, Darko Hren, Sara Schroter, Els Goetghebeur. I would particularly like to thank Mona, Melissa, and Camila, who, besides being excellent research partners, became good friends.

Finally, I would like to express my gratitude to the members of the MiRoR team who were great companions in this adventure, and to the current and former-members of the METHODS team in Paris, as well as the team at AMC in Amsterdam.

Affiliations

1. Université de Paris, Paris, France
2. INSERM U1153, Paris, France
3. Centre d'Épidémiologie Clinique, Hôpital Hôtel-Dieu, Paris, France
4. Department of Clinical Epidemiology and Biostatistics, Academic University Medical Centers, University of Amsterdam, Amsterdam, the Netherlands.

Funding

EU Horizon 2020 Marie Skłodowska Curie Actions, Grant agreement number 676207

Publications

Bertizzolo L, Bossuyt P, Atal I, Ravaud P, Dechartres A. (2019), Disagreements in risk of bias assessment for randomised controlled trials included in more than one Cochrane systematic reviews: a research on research study using cross-sectional design, *BMJ Open*. 2019 Apr 1;9(4):e028382. doi: 10.1136/bmjopen-2018-028382.

Bertizzolo L, Bossuyt PM, Atal I, Ravaud P, Dechartres A. (2020), External factors may influence Cochrane reviewers when classifying the risk of bias of original reports. *J Clin Epidemiol*. 2020 Jul;123:1-8. doi: 10.1016/j.jclinepi.2020.03.008.

Haring R, Ghannad M, **Bertizzolo L**, Page MJ. (2020), No evidence found for an association between trial characteristics and treatment effects in randomized trials of testosterone therapy in men: a meta-epidemiological study. *J Clin Epidemiol*. 2020 Jun;122:12-19. doi: 10.1016/j.jclinepi.2020.02.004.

Sharp MK, **Bertizzolo L**, Rius R, Wager E, Gómez G, Hren D. (2019), Using the STROBE statement: survey findings emphasized the role of journals in enforcing reporting guidelines. *J Clin Epidemiol*. 2019 Dec;116:26-35. doi: 10.1016/j.jclinepi.2019.07.019.

Olarte Parra C, **Bertizzolo L**, Schroter S, Dechartres A, Goetghebeur E. (2020), Use of causal claims in observational studies: a research on research study. *medRxiv* 2020.09.17.20194530; doi: <https://doi.org/10.1101/2020.09.17.20194530>

Table of Contents

ABSTRACT	- 5 -
RESUME	- 7 -
SYNTHESE DES TRAVAUX DE THESE	- 9 -
1. INTRODUCTION ET OBJECTIFS	- 9 -
2. EXPLORATION DES DESACCORDS DANS L'ÉVALUATION DU RISQUE DE BIAIS	- 12 -
2.1. <i>Raisons de désaccords dans l'évaluation du risque de biais dans les essais inclus dans deux revues Cochrane..</i>	- 12 -
2.2. <i>Influence de facteurs extérieurs sur l'évaluation du risque de biais.....</i>	- 13 -
3. ASSOCIATION ENTRE CERTAINES CARACTERISTIQUES METHODOLOGIQUES ET L'EFFET TRAITEMENT : UN EXEMPLE CIBLE PORTANT SUR LA TESTOSTERONE CHEZ L'HOMME	- 14 -
4. QUALITE DES RAPPORTS D'ETUDES OBSERVATIONNELLES.....	- 15 -
4.1. <i>Enquête sur la connaissance et l'utilisation de l'outil STROBE pour rapporter une étude observationnelle..</i>	- 15 -
4.2. <i>Utilisation d'allégations causales dans les résumés d'articles scientifiques décrivant une étude observationnelle.....</i>	- 16 -
5. DISCUSSION.....	- 17 -
6. IMPLICATIONS	- 19 -
7. CONCLUSIONS	- 20 -
SUMMARY	- 21 -
1. INTRODUCTION AND OBJECTIVES	- 21 -
2. EXPLORATION OF DISAGREEMENTS WHEN ASSESSING ROB IN RCTs	- 23 -
2.1. <i>Reasons of disagreements in risk of bias assessment for RCTs assessed in two different Cochrane reviews..</i>	- 23 -
2.2. <i>Influence of external factors on RoB assessment.....</i>	- 25 -
3. ASSOCIATION BETWEEN METHODOLOGICAL CHARACTERISTICS AND TREATMENT EFFECT: A CASE EXAMPLE ON TESTOSTERONE THERAPY IN MEN	- 26 -
4. QUALITY OF REPORTING IN OBSERVATIONAL STUDIES	- 26 -
4.1. <i>A survey on the use and knowledge of STROBE</i>	- 26 -
4.2. <i>Use of causal claims in observational studies</i>	- 27 -
5. DISCUSSION.....	- 28 -
6. IMPLICATIONS	- 30 -
7. CONCLUSIONS.....	- 31 -
SAMENVATTING	- 33 -
1. INLEIDING EN DOEL	- 33 -
2. VERKENNING VAN DISCREPANTIES BIJ DE BEOORDELING VAN RISK OF BIAS IN RCT'S.....	- 36 -
2.1. <i>Redenen voor discrepanties bij de beoordeling van risico op vertekening voor RCT's die in twee verschillende Cochrane-reviews zijn beoordeeld</i>	- 36 -
2.2. <i>Invloed van externe factoren op de RoB-beoordeling.....</i>	- 38 -
3. ASSOCIATIE TUSSEN METHODOLOGISCHE KENMERKEN EN BEHANDELEFFECT: EEN VOORBEELD OVER TESTOSTERONTERAPIE BIJ MANNEN ..	- 39 -
4. KWALITEIT VAN DE RAPPORTAGE IN OBSERVATIONELE STUDIES.....	- 40 -
4.1. <i>Een enquête over het gebruik en de kennis van STROBE</i>	- 40 -
4.2. <i>Gebruik van causale beweringen in observationele studies.....</i>	- 41 -
5. DISCUSSIE	- 42 -
6. IMPLICATIES.....	- 44 -

7. CONCLUSIES	- 45 -
BRIEF BIBLIOGRAPHY	- 47 -
ACKNOWLEDGEMENTS	- 51 -
TABLE OF CONTENTS.....	- 53 -
ABBREVIATIONS.....	- 56 -
CHAPTER 1: INTRODUCTION	- 57 -
1.1. EVIDENCE BASED MEDICINE	- 57 -
1.1.1. <i>History of EBM</i>	- 57 -
1.1.2. <i>Determining best evidence</i>	- 58 -
1.2. EVIDENCE SYNTHESIS.....	- 58 -
1.2.1. <i>Systematic reviews</i>	- 59 -
1.2.2. <i>Meta-analyses</i>	- 60 -
1.2.3. <i>Cochrane</i>	- 61 -
1.2.4. <i>Cochrane Systematic Reviews</i>	- 61 -
1.2.5. <i>Limits of systematic reviews and meta-analyses</i>	- 62 -
1.3. RESEARCH WASTE	- 63 -
1.4. THE CONCEPT OF QUALITY OF RESEARCH.....	- 63 -
1.5. BIAS IN EPIDEMIOLOGY	- 64 -
1.5.1. <i>Measuring and preventing the bias</i>	- 65 -
1.5.2. <i>Assessing the Risk of Bias</i>	- 66 -
1.6. THE COCHRANE RoB TOOL.....	- 66 -
1.6.1. <i>Meta-Epidemiology and empirical evidence of bias</i>	- 67 -
1.6.2. <i>Assessment of the RoB through the tool</i>	- 67 -
1.6.3. <i>Domains of bias and RoB Tool Items</i>	- 68 -
1.7. REVIEW OF THE LITERATURE ON REPRODUCIBILITY OF THE RoB TOOL	- 71 -
1.8. REPORTING AND REPORTING GUIDELINES.....	- 74 -
1.9. OTHER ISSUES IN POOR REPORTING; USING CAUSAL LANGUAGE WHEN REPORTING OBSERVATIONAL STUDIES	- 75 -
1.10. RATIONALE FOR THE THESIS	- 77 -
1.11. THESIS STRUCTURE.....	- 78 -
CHAPTER 2: REASONS OF DISAGREEMENTS IN RISK OF BIAS ASSESSMENT FOR RCTS ASSESSED IN TWO DIFFERENT COCHRANE REVIEWS	- 79 -
2.1. BACKGROUND.....	- 79 -
2.2. METHODS.....	- 79 -
2.3. RESULTS	- 80 -
2.4. DISCUSSION.....	- 83 -
2.5. ARTICLE	- 85 -
CHAPTER 3: INFLUENCE OF EXTERNAL FACTORS ON RoB ASSESSMENT	- 97 -
3.1. BACKGROUND.....	- 97 -
3.2. METHODS.....	- 97 -
3.3. RESULTS	- 98 -
3.4. DISCUSSION.....	- 99 -
3.5. ARTICLE	- 101 -
CHAPTER 4: ASSOCIATION BETWEEN METHODOLOGICAL CHARACTERISTICS AND TREATMENT EFFECT: A CASE EXAMPLE ON TESTOSTERONE THERAPY IN MEN.....	- 111 -

4.1. BACKGROUND.....	- 111 -
4.2. METHODS.....	- 111 -
4.3. RESULTS	- 112 -
4.4. DISCUSSION.....	- 112 -
4.5. ARTICLE.....	- 115 -
CHAPTER 5: A SURVEY ON THE USE AND KNOWLEDGE OF STROBE	- 125 -
5.1. BACKGROUND.....	- 125 -
5.2. METHODS.....	- 125 -
5.3. RESULTS	- 126 -
5.4. DISCUSSION.....	- 127 -
5.5. ARTICLE.....	- 129 -
CHAPTER 6: USE OF CAUSAL CLAIMS IN OBSERVATIONAL STUDIES	- 141 -
6.1. BACKGROUND.....	- 141 -
6.2. METHODS.....	- 141 -
6.3. RESULTS	- 142 -
6.4. DISCUSSION.....	- 142 -
6.5. ARTICLE.....	- 145 -
CHAPTER 7: DISCUSSION AND IMPLICATIONS.....	- 175 -
7.1. DISCUSSION.....	- 175 -
7.1.1. <i>Multidisciplinary approach</i>	- 175 -
7.1.2. <i>Novelty and main results</i>	- 176 -
7.2. IMPLICATIONS	- 179 -
7.3. CONCLUSIONS.....	- 182 -
BIBLIOGRAPHY	- 183 -

Abbreviations

ABBREVIATION	Expansion
RoB	Risk of Bias
RCT	Randomized Controlled Trial
SR	Systematic Review
Evidence Based Medicine	EBM
CDSR	Cochrane Database of Systematic Reviews
Reporting Guidelines	RG

Chapter 1: Introduction

1.1. Evidence Based Medicine

Evidence based medicine (EBM), is defined as “the conscientious, explicit and judicious use of current best evidence in making decisions about the care of individual patients”[1]. EBM represented a new paradigm integrating clinical expertise, patient’s values and best available evidence in the process of decision making and health practice [57]. The central principle of EBM is that the foundation of any clinical medical decisions should rely on the best scientific evidence from clinical research[2]. EBM converts the exercise of reading and appraising the literature into a pragmatic process of using the literature to benefit individual patients while simultaneously expanding the clinician’s knowledge base [58]. As a result, EBM requires different skills from researchers and clinicians than traditional medicine does. These skills include efficient literature-searching and the knowledge and application of rules of epidemiology which permits to critically appraise the clinical literature. Each clinician will have during his/her professional life to evaluate scientific articles in order to provide the best level of care to patients according to the current best evidence from the scientific literature [59].

1.1.1. History of EBM

The origin of EBM principles is debated; some authors find traces of its origin in ancient Chinese medicine, while others think it was ancient Greece. We know for sure that testing medical interventions for efficacy has existed since the 11th century when Avicenna wrote The Canon of Medicine[60]. Conversely, the clear introduction of EBM in the last century can be attributed to Professor Archibald Cochrane, who wrote the book “Effectiveness and Efficiency: Random Reflections on health Services” published in 1972 and advocated for use of scientific evidence in medicine[61]. His work and ideas were honoured through the naming of an international organization devoted to EBM as Cochrane Collaboration (now Cochrane) [62]. The term “evidence based” itself was first introduced by David Eddy in 1989[2, 63], while speaking about the creation of clinical practice policies, which are important to facilitate and homogenise decisions in clinical practice. Eddy highlighted the limits of the traditional approach which identified “standard and accepted” practices creating a “dangerous tautology”: “what people *should do*, being based on what people *are doing*”. The approach hinted by Eddy, which suggested to have a more outcome-based view on issues, was then developed by the research group of David Sackett at the McMaster University in Canada, which first used the term “evidence-based medicine in 1992 [57, 59, 64]. The concept was initially proposed as a different approach to teaching the practice of medicine. Very

soon EBM moved its purpose from teaching to practice, extending its scope to clinical practice[62]. The development of EBM was facilitated by the creation of electronic health databases and the increased use of the internet[59]. In parallel, the research group led by David Sackett and Gordon Guyatt at the McMaster University also engaged a reflexion on how to hierarchize evidence and identify the best evidence[65].

1.1.2. Determining best evidence

EBM categorizes different types of studies and ranks them according to their level of evidence and risk of bias[66]. This classification is often translated into a pyramid [67] represented in Figure 1, which varies slightly across authors and research groups without changing the basic meaning. According to EBM the best evidences are obtained by systematic reviews and meta-analyses of randomized controlled trials (RCT). Just below there are evidences from only one of such RCTs; but according to some authors a large RCT could be considered as having also the highest level of evidence[68]. Immediately below this level is evidence from well-designed non-randomized research and evidences from quasi experimental research. At the lower level of this pyramid we found evidences from non-experimental studies (case control studies, case series...), and textbooks. At the lowest level we have opinions from experts.

Figure 1: Pyramid of evidences. (from Reddy KR, 2018[69])

1.2. Evidence Synthesis

As previously discussed, systematic reviews and meta-analyses are generally considered to provide the highest level of evidence. Apart from their role in providing best evidences, SRs are also useful to avoid replication of studies, therefore preventing research waste.

Nowadays, because of the exponential growth in production of medical knowledge, the widespread interest in the field of evidence synthesis have become a necessity. For a modern physician it would

be unfeasible to follow the huge and increasing mole of medical information in a timely way and in her/his busy daily activity. In 2010, 75 new randomized trials and 11 systematic reviews were published every day[3]. A search of Medline, in January 2021, shows more than 32 million entries in the database [70]. Moreover, many clinicians still lack the skills to differentiate what literature is worth considering and what is garbage.

Continuing from the teachings of Archibald Cochrane, the process of Evidence Synthesis has become one of the tasks of EBM-oriented professionals [60, 71-74]. Beyond the need to synthesize a huge amount of data, such synthesis is also essential to limit waste in research[6, 7, 28, 75].

Previous researchers highlighted that all new research studies should begin and end with a carefully conducted evidence synthesis [76-78]. The reasons behind this is that, in a first moment, a synthesis of the literature is necessary to evaluate whether the research question has been answered in the past. If the clinical question has not been properly addressed yet, then a SR could help in shaping in the best way the research question [6, 7, 75].

1.2.1. Systematic reviews

In the field of evidence synthesis, systematic reviews and meta-analyses are considered the gold standard[78]. A systematic review is “a review of a clearly formulated question that uses systematic and explicit methods to identify, select, and critically appraise relevant research, and to collect and analyse data from the studies that are included in the review”[79]. The methods should also be rigorous and reproducible, allowing to minimize bias[79]. Therefore, SRs are characterized by the following process:

- A clear clinical question, with precise definition of objectives;
- A systematic search of evidences, which will identify all studies meeting the eligibility criteria;
- The critical appraisal of studies identified including an evaluation of their risk of bias;
- A qualitative or quantitative synthesis to communicate the results, which should be effective in answering the clinical question of the objective and thus make the SR useful for clinical practice.

The cue for the creation of a review is the need for a clinician to find the best and most recent synthesis of existing data that will allow to solve an individual patient clinical problem, in particular "Which treatment to choose?" [80-83].

The clinical question needs to be translated in a good research question, which should be clear, directly aimed to the problem, and answerable. EBM advises to use the so-called PICO format. The

PICO format identifies four critical components in every research question. These components are Patient or problem, Intervention, Comparison and Outcome[59, 84].

1. P identifies the patient or problem. This concept aims at the precise identification of who are the relevant patients, meaning who are the patients that may benefit from the intervention. If the research does not involve patients then it should be the problem being the focus;
2. I is for intervention, which identifies the management strategy, a treatment either pharmacological or not or a diagnostic test or exposure of interest. This aspect concerns the definition of the drugs, diagnostic test or even a particular food or a surgical procedure that will be examined in relationship to the defined patient;
3. C is for comparison. This is the identification of the control or alternative management strategy, test or exposure that will be compared with the intervention. This will define against what the results of the intervention will be measured;
4. O is the outcome, which specifically identifies the patient-relevant consequences of the intervention evaluated.

The second part of the SR process is the search for adequate information. Information should be valid (containing high quality data), relevant (for EBM this essentially means to be clinically applicable), comprehensive (including every data on benefits but also on harms of interventions analysed). The process of writing a SR includes the search for less accessible source of information including grey literature (referring to unpublished data). The search of different sources is very important, because it may greatly impact the results of a SR.

The next step in producing a systematic review is the critical evaluation of the information. This step involves the careful reading and appraisal of methodology, contents and conclusions of individual studies.

The last step is the translation of the evidence to the practice, which means to apply information to the patient. This question is not minor because a doctor needs to decide whether the evidence could apply to his/her patients.

1.2.2. Meta-analyses

A systematic review may or may not be followed by a meta-analysis. A meta-analysis consists of a statistical analysis that allows one to combine the results of several studies in order to obtain a single summary estimate of the effect [85, 86]. A meta-analysis can be done for each outcome defined in the review if there is a sufficient number of studies. The meta-analysis produces a weighted average of the intervention effect estimated in the individual studies. The weighting

accounts for the study precision (determined by the variance of a study) so the studies with the most precise intervention effect will have the most “weight” in the meta-analysis. To conduct a meta-analysis, the studies must conceptually have evaluated the same population and interventions even if there are some differences across studies. Therefore, statistical heterogeneity should be systematically assessed when conducting a meta-analysis and it could be explored thereafter.

1.2.3. Cochrane

Cochrane[87] (previously known as “Cochrane Collaboration”) is an international not-for-profit and independent organization. The mission of Cochrane is making readily available worldwide up-to-date, accurate information about the effects of healthcare interventions. Cochrane produces and disseminates systematic reviews of healthcare interventions and has a pivotal role in guaranteeing the importance of EBM in research panorama.

Cochrane mission is very broad and cohesively aimed to “promote evidence-informed health decision-making by producing high-quality, relevant, accessible systematic reviews and other synthesized research evidence” [87, 88]. For reaching this purpose Cochrane is active in a series of activities concerning EBM[89, 90]. First, Cochrane worked to formalize and coordinate the efforts in the field of Evidence Synthesis. Cochrane produced and revised through successive editions an handbook which is regarded as one of the most important guidelines on how to conduct SRs[79]. Together with this is the creation of a software called Revman[91], which is a widely used software for SRs. To channel the adoption of EBM and promote the use of SRs, Cochrane has a widespread worldwide network ensuring the teaching of the fundamentals of EBM and of Evidence Synthesis. The Cochrane Library[92] is the expression of the mandate of Cochrane to produce high quality research and importance and its most visible accomplishment. The Cochrane Library is a collection of different databases; it includes the Cochrane Database of SRs (CDSR), the Cochrane Central Register of Controlled Trials (CENTRAL) and Cochrane Clinical Answers (CCA). CENTRAL includes reports of RCTs and quasi-randomized trials, CCA represents a readable compendium of important clinical questions. The main product is the CDSR which is one of the most important collections of SRs and includes all Cochrane Reviews and protocols written since the inception of Cochrane. The CDSR itself had an impact factor of 7.890 in 2019[93].

1.2.4. Cochrane Systematic Reviews

Cochrane Systematic Reviews are recognized for their quality [94]. They are an important source of information when developing clinical practice guidelines and they cover the broadest magnitude of

topics[78]. The organization of Cochrane is through different Cochrane Review Groups, each dedicated to a specific topic area. The Cochrane groups at the time of this document (early 2021) are organized in eight networks: Abdomen and Endocrine, Acute and Emergency Care, Cancer, Children and Families, Circulation and Breathing, Mental Health and Neuroscience, Musculoskeletal, Oral, Skin and Sensory and Public Health and Health Systems. The organization in specialized group mirrors the commitment for Cochrane to have SRs to be conducted by experts in the topic of interest[87]. Cochrane SRs are available in the CDSR, which includes more than 8000 reviews[92]. Cochrane SRs follow a shared and rigorous methodology and are conducted by members with training and expertise. Cochrane SRs are also periodically updated [79].

1.2.5. Limits of systematic reviews and meta-analyses

In the previous paragraphs we explained the importance of SRs to synthesize existing research by using a precise and rigorous process. Despite all this, SRs and meta-analyses can have numerous limitations, which are mainly related to the fact that both SRs and meta-analyses use a retrospective approach and are, in their essence, observational studies. Although the process is precisely formalized, they are highly dependent on the quality of primary studies they include. The main factors are the quality and completeness of reporting and the absence of bias in primary research. SRs are affected by reporting bias in the scientific literature. Numerous studies have demonstrated how certain studies tend to be more often published and therefore more easily accessible than others. Previous evidence showed that studies with statistically significant results are more frequently published (the so-called “publication bias”) [95-100] and also, more rapidly published[101, 102]. Such studies are also more often published multiple times [103, 104]. Another aspect of reporting bias is related to selective outcome reporting. Selective outcome reporting happens when one outcome of the study is changed or omitted in the publication, such as outcomes reaching statistical significance are highlighted, while other are omitted or degraded to secondary[37, 105, 106].

SRs are also affected by the methodological quality of included studies. The quality of research is mostly defined by their absence of bias. Bias are systematic errors leading to an overestimation or underestimation of treatment effect. If the studies included in a SR are biased the SRs pooling their results will also be biased. They are also affected by the poor quality of included studies preventing researchers to extract the data or to assess the main characteristics of the studies including their methodological quality. These two aspects of bias and poor quality of reporting result is an

important source of waste in research affecting not only primary research but also systematic reviews and meta-analyses.

1.3. Research waste

Recent advances in research methodology and in requirements by regulatory agencies have led to an increase in quality of research[3]. Despite these recent efforts, the quality of scientific literature is still considered to be overall not optimal[3]. It has been estimated that up to 85% of biomedical research could be wasted[4, 5]. Such waste can happen at every step of the research process from the planning to the reporting of the findings. At the planning stage, researchers may formulate the wrong research question, setting irrelevant research priorities or using an inappropriate design. Research waste can also happen during the study conduct, by using methods that are prone to induce bias in the results. Finally, there may be waste caused by the poor reporting of results, the scientific communication could be incomplete or for example use wrong terminology[6-9].

1.4. The concept of quality of research

Evaluating the quality of individual studies is a key step of the systematic review process. The process of determining the quality of trials has followed a long road. For a long period, the efforts of methodologists were directed toward the creation of “quality scales”. Such scales evaluated different notions and used different types of scoring system to give a quantitative information regarding either the research conduct and/or the research reporting. The number of such scales increased rapidly and in a SR in 1996, Moher et al. described the existence and use of 25 scales and 9 lists of items to assess quality. Among those, only a small number were developed using scientifically recognized methods [107].

The study of Moher highlighted the limits of these quality scales. These scales were in fact mixing very different concepts such as internal or external validity of a study and the profound difference between the study conduct and the study reporting. But the final word on the question was posed in 1999 by Peter Jüni et al. In a famous meta-analysis the research group completely questioned the use of scales to assess the quality of trials[108]. In the study, the researchers used 25 different quality scales to assess 17 RCTs included in the meta-analysis. The study showed that with different scales the results would have been completely different. The conclusions of the study of Peter Jüni discouraged the use of quality scales any further; their recommendation was to evaluate

methodological characteristics of the studies as a function of their influence on the treatment effect estimates[108]. The study marked a turning point in the way of evaluating the validity of trials, opening the way to the RoB assessment.

1.5. Bias in epidemiology

In epidemiology “bias” represents a systematic error. The bias translates in a deviation from the truth, which will consequently be reported in the results of a study or its inferences. One of the main characteristics of bias is that it can operate in either direction, which means the bias can lead to underestimation or overestimation of the true intervention effect. The magnitude of bias also varies from small to substantial[13, 109, 110].

Bias is different from imprecision. Bias refers to a systematic error, while imprecision represents a random error. One of the main differences between the two is that the systematic error cannot be corrected by multiple replications. With a random error, multiple replications of the same measure will produce different estimates, scattering in different directions and of different magnitude; therefore, repeating an imprecise study over and over would eventually give the right answer because of sampling variation. Quite on the contrary, multiple replications of the same biased study would reach a similarly wrong answer, constantly. In practical terms, bigger sample size and repetition should resolve imprecision, but cannot affect bias.[21]

Bias is different from quality. The terms have been confused often, but they represent different concepts and have different implications. Bias concerns a measurement and it defines how close the measurement is to the true value. Quality, on the other hand, concerns a series of characteristics, which are perceived as optimal. The definition of quality does not aim directly to a measure, and therefore to a practical aspect. To summarize the definition of quality is close to the concept of good research, or good clinical practice. Therefore, there exist some well-known markers of “good research”, such as obtaining an informed consent from a patient, declaring conflict of interest in research, publishing even negative results which, most probably, do not affect the measurement of an outcome[111, 112].

The bias is one of the main reasons of heterogeneity of results among different studies. The bias explains why and how the studies which are more methodologically accurate, show results that are closer to the “truth” or better, the true value of interest. Bias involves several aspects of research, impairing the study on several levels. Cochrane used to classify bias in four or five different domains[113]:

- Selection bias, which represent systematic differences between the baseline characteristics of the study groups. This type of bias can usually be counteracted through an adequate generation of the randomization sequence as well as allocation concealment.
- Performance bias, which represent systematic differences in the administration of every additional care (apart from the intervention of interest) between the study groups to be compared. It happens when, for example, a group is administered a co-intervention more often (e.g. corticoids when evaluating a biotherapy). Usually, administering the treatment without knowledge of which treatment the patient receives (the “masking” or “blinding”) limit this bias.
- Detection bias. It represents systematic differences in the assessment of the study outcomes between the study groups. This bias is particularly present for subjective measures[114]. This bias can also be corrected by blinding of the treatment allocation.
- Attrition bias represents systematic differences between study groups regarding withdrawals from study participation. There are different causes of withdrawals, such as lost to follow-up, drop out of the study, non-compliance with treatment. The attrition bias is by limiting as much as possible the number of participants lost to follow-up and using an intention-to-treat analysis (ITT) associated with an adequate management of missing data.[115]
- Reporting bias, which concerns the selective reporting of the information from research. Information regarding methodology or results may be omitted or reported in a misleading way.

1.5.1. Measuring and preventing the bias

The identification and measurement of bias is difficult. Paradoxically, a study with numerous flaws in its methodology may by chance be unbiased, while the contrary is also possible. As we explained in a paragraph above, bias represents a systematic error from the truth, therefore, examining a study, we cannot infer whether bias has occurred and establish the potential direction and magnitude of this error as the true value of the parameter is unknown. In brief, we cannot measure bias by itself. Researchers in clinical epidemiology have therefore established that it is more appropriate to focus on the risk of bias. RoB recognises and measures the likelihood of bias to happen. This translates in the identification and assessment of features of a study, which could possibly lead to misleading results. This characteristic concern the design or the conduct of the study [10-13]. The evaluation of RoB in studies is therefore a possible solution to reveal the occurrence and the magnitude of bias that may have occurred.

1.5.2. Assessing the Risk of Bias

In 2005, Cochrane identified a working group of methodologists, medical journal editors and authors of systematic reviews to develop a tool to assess RoB in RCTs; this effort produced the RoB tool [19]. The RoB tool developed by Cochrane is nowadays used in all Cochrane SRs and has also been increasingly common among non-Cochrane reviews. A study of 2013[116], showed that Cochrane reviews were already using the tool extensively (82% of the Cochrane SRs used the RoB tool, and 9% a slightly modified version), while the non-Cochrane reviews were just beginning the adoption (20% of them was using the RoB tool while 38% of the non-Cochrane reviews was still using quality scales).

1.6. The Cochrane RoB tool

The panel identified by Cochrane agreed on main principles that became the basis of the new tool for assessing the RoB in RCTs[19, 113]. These main principles are:

- To avoid the use of scales to assess quality, because such scales have already proved misleading;
- To focus on internal validity. Internal validity of a study is mainly defined as the absence of bias. The tool does not evaluate external validity of a study or its precision;
- To evaluate the RoB of the results of a study. The tool does not assess the completeness of the study report or the presence of methodological flaws which are not directly associated to RoB (such as the trial registration). The tool should focus on study “conduct” and not study “reporting”;

These principles guided the creation of the tool. At the same time, the methodologists working on the tool decided to address the issues that concerned most of the previous attempts to measure the validity of a study. Therefore, they provided indication to make the tool more replicable and give it a stronger methodological background.

It was decided that, because of the subjectivity of every judgement, the assessment should provide every element that was used to deliver the judgement (support for judgement).

The methodological background of the tool was chosen to be both theoretical and empirical. The items to be evaluated in the RoB tool were chosen through evidences from meta-epidemiological studies evaluating the association between study characteristics and treatment effect.

1.6.1. Meta-Epidemiology and empirical evidence of bias

The tool developed by Cochrane is based on empirical evidence of characteristics associated with intervention effect, which is known as meta-epidemiology[13-18, 117, 118]. The starting point of a meta-epidemiological study is a collection of meta-analyses. The studies included in these meta-analyses are then analysed according to some study-level characteristics. One of the first examples of meta-epidemiological studies was the one of Schulz[14], which in 1995 showed that RCTs in which randomization was inadequately concealed or reported lead to inflated estimates of intervention effect compared with trials with adequate concealment.

One of the basic processes of meta-epidemiological studies is the calculation of the ‘ratio of odds ratios’ within each meta-analysis. This means, for example, to evaluate the intervention odds ratio in trials with inadequate randomization process by dividing it by the odds ratio in trials with adequate randomization. These ratios of odds ratios are then combined across meta-analyses, with a meta-analysis, which explains what this method is sometimes also called ‘meta-meta-analysis’. The ratio of odds ratios represents the average difference on treatment effect between trials with and without the characteristic of interest, for example here, inadequate randomization process.

1.6.2. Assessment of the RoB through the tool

The RoB assessment of each item consists of two parts:

- The judgment, which is expressed as a level of risk. The risk can be: low risk of bias, high risk of bias and unclear risk of bias if there is insufficient information to issue a judgment[13]. It is relevant to stress that for the Cochrane RoB tool, the risk of bias can be diminished to a low level but can never be “zero”; the risk of bias is a feature of a study that cannot be removed.
- The support for judgement, which is a brief text summary from which judgements of risk of bias can be made. The purpose is to ensure transparency in how the judgement is reached. The text may include verbatim quotes from the source or alternatively another known fact regarding the study or the author, or a comment from the review authors

1.6.2.1. *Retrieving essential information for a judgement*

The RoB tool aims to assess the conduct not the reporting of a study. Nevertheless, Cochrane acknowledges that in most cases the source for information is going to be represented by published reports, usually in the form of journal articles. Cochrane provides precise guidance regarding how to retrieve and collect the relevant data. Among preferred sources of information apart from scientific articles already cited there are book chapters, dissertations, conference abstracts and web

sites including trials registries and research protocols. Given the high frequency of incomplete and unclear reporting, an important advice given by Cochrane is to search for additional sources and also contact the study author with open-ended questions regarding the study conduct.

1.6.3. Domains of bias and RoB Tool Items

The different types of bias were exposed previously. In the Cochrane RoB tool, each different type of bias represents a domain of the tool. Within a domain are present the different items that are the actual basis of assessment.

1.6.3.1. *Selection bias; Random Sequence Generation and Allocation Concealment*

Selection bias refers to systematic differences at baseline between the characteristics of the groups that are compared. The randomization correctly and successfully performed is the best way to avoid such differences both for measurable and unmeasurable or unknown factors.

The key moments in the process of a correct randomization procedure are two: the generation of a randomized sequence and the concealment of such allocation rule.

The randomized sequence is the rule for allocating interventions to participants. To be valid it should be based on a random process. This corresponds to the item in the RoB tool: Random Sequence Generation.

The second moment in the process of an effective randomization is represented by every step taken to secure the implementation of the randomized sequence by preventing foreknowledge of the forthcoming allocations. This process is called allocation concealment, although should more correctly be called allocation sequence concealment. The item in RoB tool identifying this moment is called Allocation Concealment.

The concealment of the allocation sequence is linked to the generation of the random sequence, this is easily explained by examples. A simple assignment rule, such as alternation could achieve a balance in the characteristics of participants[119], similarly to a random rule. However, if future assignments can be anticipated then bias can arise due to the selective enrolment and non-enrolment of participants into a study. Conceptually, the patients who would have been assigned to an intervention considered ‘inappropriate’ by their carer may be rejected or put ‘on hold’ until the ‘right’ choice come up in the sequence. From a practical point of view, knowing the next assignment could completely invalidate the randomization of the sequence.

The empirical importance of random sequence generation has been assessed in meta-epidemiological studies[14, 15, 120, 121], similarly has happened for allocation concealment[114, 122, 123].

1.6.3.2. Performance bias; Blinding of participants and personnel

Performance bias happens when there are systematic differences (other than the one of interest) in the intervention between the groups included in the RCT. The difference for example could be a change in the care that is provided, or the exposure to factors other than the one of interest. Such differences tend to happen when the participants or the personnel caring of them are aware of their allocation to either intervention of interest. For this reason, after enrolment into the study, the masking (blinding) of study participants and personnel to the allocation may reduce the risk that knowledge of which intervention was received affects outcomes. The blinding should ensure that the different groups receive similar attention, in the form of ancillary treatment or additional diagnostic investigations. The item in the RoB tool relative to this bias is called “Blinding of Participants and Personnel”.

There are empirical evidences from meta-epidemiological studies that the lacking of blinding of participants or personnel or both is associated with differences in treatment effect estimates [17, 114, 123-126].

It is important to underline the difference between allocation concealment with blinding of allocated interventions. Allocation concealment happens before blinding; it is aimed to prevent the knowledge of the allocation sequence *before and until* assignment. It can always be successfully implemented, regardless the study topic and design [14, 127]. Blinding happens *after* assignment and cannot always be implemented, for example when the intervention is surgery[127-129]. Briefly, allocation concealment is up to the point of assignment of the intervention and blinding just after that point.

1.6.3.3. Detection bias; Blinding of outcome assessment

Detection bias happens when there are systematic differences in how outcomes are determined (measured) between groups. The way to prevent this to happen is usually through masking (blinding) the outcome assessors of which intervention was received. It is important to notice that the assessors may be the participants themselves, their healthcare providers, or other possibly independent assessors. Similarly, to blinding of participants and personnel, in some cases the blinding of outcome assessors can be impossible. The lack of blinding is especially important for assessment of subjective outcomes (e.g. pain), while in cases of objective outcomes the importance of this item is negligible (e.g. all-cause mortality). Consequently, when considering this bias there are two aspects to determine: who is assessing that specific outcome and how “subjective” or “objective” the assessment of the outcome is. For this item is also relevant that the assessors of some outcomes may be blinded while assessors of other outcomes are not. For example, patients who received a certain type of surgery could not be blinded for a pain outcome, but the personnel

taking care of her/his recovery may be unaware of that. For all these reasons Cochrane suggests assessing this item for each outcome, or at least “type” of outcome (objective/subjective) in the study.

Empirical evidence from meta-epidemiological studies, shows that the lack of blinding of outcome assessors in RCTs is associated with more exaggerated estimated intervention effects[114, 123],

1.6.3.4. Attrition bias; Incomplete Outcome Data

Attrition bias characterises the situation when there are systematic differences between groups in withdrawals (dropouts) from a study which may result in a bias in the observed effect estimate. In the risk of bias tool, this aspect is examined in the “incomplete outcome data” item. It is important to differentiate between different mechanisms that cause outcome data to be incomplete; a main difference is between exclusions and attrition.

Exclusions refer to situations in which data from certain participants are omitted from analyses, despite being available to the trialists; an example for exclusion is when enrolled participants are later found to be ineligible.

Attrition happens when outcome data are not available. Possible reasons for attrition are related to participants withdrawing from the study, not attending scheduled appointments, or not completing evaluations. Even if attrition happens, it is important to consider that the risk of bias related to this aspect depends on several factors. A partial list of those includes: the amount and distribution of missing data across the different intervention groups, the reasons for outcomes being missing, the likely difference in outcome between participants with and without data, the strategy that authors used to deal with the problem, the clinical context.

To prevent this type of bias the first approach is to diminish attrition as much as possible, boosting adherence and retainment of participants. Another systematic approach often used to minimize bias is to perform an intention-to-treat (ITT) analysis. The basics of ITT analyses are used inconsistently in study reports[130]; nevertheless a couple of principles should always be considered: to keep participants in the intervention groups to which they were randomized (regardless of the intervention they actually received or their adherence to intervention); and include all randomized participants in the analysis. The first principle can always be applied, while often not all participants can be included in the analysis, attrition cannot always be controlled. Therefore, researchers need to make assumptions about the missing values, using the principles of imputations. These principles are statistical ways to account for missing data in the analysis. In contrast to ITT analysis, the “per protocol” or “as treatment received” analysis have been found to exaggerate estimates of the effect of assignment to intervention compared with ‘intention-to-treat’ analyses[131, 132].

Meta-epidemiological evidences regarding this bias are not strong and not concordant[14, 16, 114, 120, 131-134]; concerns over bias resulting from incomplete outcome data are mostly driven by theoretical considerations.

1.6.3.5. Reporting bias; Selective Reporting

Reporting bias refers to systematic differences between reported and unreported findings; this may for example refers to report analyses with statistically significant results, while the ones with non-significant ones are omitted from publication. In the RoB tool this type of bias is assessed in the item “selective reporting”. The reporting bias includes different ways of altering the reporting such as using only certain timepoints for analysis, using a variable as continuous or dichotomous, using different scales or measures, changing primary outcome to secondary (or vice versa)

The problem of assessing this bias relies in the difficulty in detecting it. To understand if authors changed something in planned analyses most of the times is necessary to compare the published report with a protocol or registration form. Especially for older studies, before the CONSORT statement [135], finding protocols is often impossible.

Meta-epidemiological evidences supporting the inclusion of this item in the tool are present [37, 105, 136-139].

1.7. Review of the literature on reproducibility of the RoB tool

An important limitation of the RoB tool outlined by previous studies is related to its suboptimal reproducibility [10, 20-25]. The majority of these studies evaluated agreement between 2 independent reviewers, using the K-Statistic test [140] or across different groups using the Fleiss’ K-Statistic [141]. The classification criteria for interpreting kappa values vary slightly across publications; but mostly followed the criteria proposed by Byrt [142]: values between 0.93 and 1.00 were considered excellent, between 0.81 and 0.92 very good; 0.61–0.80 good; 0.41–0.60 fair; 0.21–0.40 slight; 0.01–0.20 poor and less than 0.01, no agreement. The findings of the literature on the topic are exposed below.

Hartling, 2009 [21], studied the agreement between two different reviewers (“inter-rater reliability”) on 163 full manuscripts. They found a variation of inter-rater agreement for the seven items that ranged from good agreement for Random Sequence Generation ($k=0.74$) to poor agreement for the Selective Outcome Reporting item ($k=0.13$). This study reached the conclusion that the differences in the RoB agreement for the different items were caused by the way the assessment is conducted for that specific item. For judging the RoB of some items, the assessors rely more on the actual reporting

from the manuscript, while other items require more interpretation and leave assessors “freer” in their judgement. Consequently, domains as “Selective Outcome Reporting”, relying more in interpretation and open judgment perform worse in agreement.

Hartling 2011 [10], used same methodology on a different subset of 107 manuscripts. They also compared the RoB assessment with the use of another “quality” scale (Jadad scale) The results were slightly better than in the previous study as inter-rater agreement varied from slight (0.40) for Selective Outcome Reporting to very good (0.86) for Random Sequence Generation. Remaining domains shown fair agreement ($k = 0.44\text{--}0.60$).

Hartling 2013 [20], used a similar approach on 154 full manuscripts but also evaluated agreement between couples of researchers (“inter-consensus reliability”) on a subset of 30 manuscripts. The inter-rater agreement was slight for most of the domains ($k = 0.24\text{--}0.37$) except for sequence generation which showed good agreement ($k = 0.79$). Across reviewer pairs agreement was poor ($k = 0.05\text{--}0.09$) for Incomplete Outcome data, Selective Outcome Reporting and Blinding, slight ($k = 0.37$) for Allocation Concealment and fair ($k = 0.60$) for Random Sequence Generation. The agreement was therefore worse when comparing the reconciled results of different couples (inter-consensus reliability) than between two independent assessors.

Armijo-Olivo 2012 [23], evaluated inter-rater agreement on 20 different RCTs, reported in 25 manuscripts. They report inter-rater reliability across individual domains of 0.30 (slight), without detailed results by domain.

Armijo-Olivo 2014 [22], used a different methodology: 17 Cochrane SRs were identified and 109 trials were extracted. Two independent researchers evaluated RoB and inter-rater agreement was estimated. After consensus, the RoB assessment was compared to the results reported in the SRs (agreement between blinded external reviewers and Cochrane review authors). Inter-rater agreement was fair ($k = 0.50\text{--}0.56$) for Selective Outcome Reporting, Blinding of Participant and Personnel and Blinding of Outcome Assessment and was good ($k = 0.71\text{--}0.79$) for Incomplete Outcome Data, Random Sequence Generation and Allocation Concealment. After-consensus agreement with the Cochrane results was poor ($k = 0.13\text{--}0.15$) for Selective Outcome Reporting and Blinding of Participants and Personnel and slight ($k = 0.24\text{--}0.41$) for Incomplete Outcome Data, Allocation Concealment, Blinding of Outcome Assessment; only Random Sequence Generation showed good agreement ($k = 0.62$). In the results, the authors acknowledge that the tool itself may be subjective and not reliable, being dependent on the reviewers’ level of knowledge and familiarity with the information provided in the individual trials.

Jordan 2016 [24] used solely data from Cochrane reviews, comparing RoB assessment when a single study was evaluated in two different SRs. Their sample size consisted of 46 studies, from a single

Cochrane group (“Menstrual disorders and Subfertility”). Not every study in their sample reported RoB assessment for each domain in the tool. The agreement was calculated as percentage. In the results, on average, the agreement was comparable to previous findings; in this sample, nevertheless, agreement was low (35%-43%) for Blinding items and Selective Outcome Reporting respectively, mediocre (59%) for Allocation Concealment and good (70%-71%) for Incomplete Outcome Data and Random Sequence Generation.

Da Costa, 2017 [25], proposed an experimental approach; 3 groups of pairs of reviewers were invited to evaluate 56 different trials. Groups were formed as following: a couple of experienced researchers which served as reference, a couple of unexperienced reviewers who got intensive training in the use of RoB tool and a second couple of unexperienced researchers who had minimal training. The study evaluated both the inter-reviewer agreement and the one between groups after consensus. Within group agreement was as following: for minimal training group there was no agreement ($k=0$) for Blinding of Outcome Assessment, poor/slight agreement for Blinding of Participants ($k=0.14$) and Personnel ($k=0.30$), slight agreement for Random Sequence Generation and Allocation Concealment ($k=0.30-0.38$). In the training group agreement was excellent for every domain ($k=0.93-1.00$). The between group agreement was as following; between inexperienced group and reference was poor ($k=0.10$) for Incomplete Outcome Data, slight ($k=0.25$) for Blinding of Outcome Assessment, slight/good for Blinding of Personnel ($k=0.40$) and Participants ($k=0.81$), fair ($k=0.57$) for Allocation Concealment and good ($k=0.71$) for Random Sequence Generation. The comparison between the training group and reference showed constantly better results. Agreement was fair ($k=0.41$) for Blinding Outcome Assessment, fair/very good for Blinding of Personnel ($k=0.50$) and Participants ($k=0.90$), good ($k=0.62$) for Incomplete Outcome Data, very good for Random Sequence Generation ($k=0.82$) and Allocation concealment ($k=0.87$). Conclusions underline the importance of training in improving inter-rater agreement but the effect on between-group agreement and therefore on reproducibility of the tool is limited.

Overall, the literature on the topic agrees that the tool has a suboptimal reproducibility. Nevertheless evidences, although numerous, do not cover the issue completely. Most studies were set in an experimental setting and often only assessed the agreement between two different reviewers. Only the study of Jordan [24] used a real-life approach, but the sample size was very limited especially for certain items of the tool. Moreover, none of the study clarifies what could be the reasons of disagreements in RoB assessment.

1.8. Reporting and reporting guidelines

Another source of the poor quality of research and a barrier of good evidence synthesis is related to the poor quality of reporting of studies. Study reporting should respond to three main characteristics: it should be clear, complete and easily available. If important information of a study is poorly reported or not reported this usually makes the replication, critical appraisal and interpretation of results difficult or even impossible. Poor reporting also limits the RoB assessment validity, making poorly reported items to be generally evaluated at “unclear RoB” in the assessment. With inadequate reporting, studies are also less likely to be properly considered for inclusion in SRs and meta-analysis with the results that studies cannot properly inform clinical practice or health policies, contributing again to the overall problem of waste in research[26-28]. Although the quality and completeness of reporting has been improving in the recent years, numerous studies have shown the poor quality of reporting across various topics and study types[15, 29-35]. To address the issue of poor reporting, the research community has developed structured guidance to scientific writing with the creation of tools such as reporting guidelines (RG) [4, 36, 37]. The first structured efforts to improve the quality of reporting was aimed toward RCTs and can be traced back to 1993 when the Standardized Reporting of Trials (SORT) Statement was created [38, 39]. Simultaneously and independently, a different group of researchers, the Asilomar group, was also working on providing guidance for reporting. As a result of their shared effort, in 1996, the SORT group, in collaboration with the Asilomar Working Group, published the Consolidated Standards of Reporting Trials (CONSORT) Statement[40-42]. The CONSORT Statement is vastly considered one of the first structured RG. The RG, although initially conceived for RCTs, has inspired many offshoots, which are known as “extensions” of the CONSORT Statement, which provide specific guidance for different study designs, interventions, or data types[43].

RGs are often organised as checklists, containing essential items that should be reported in a publication. The checklists may also be accompanied by a flow diagram, which allows the detailing of additional information, such as how many people were initially contacted, how many declined to participate, and the number of dropouts. The RGs are also frequently associated with an “Elaboration and Explanation” document. The document provides a detailed description of the items in the checklist and their use. It also includes supporting background for the tool. Although such document is not a training support for the use of the tool, it gives additional information in the attempt to provide some education alongside the strict writing guidance.

In the field of observational research, one of the most used RGs is the STrengthening the Reporting of OBservational studies in Epidemiology (STROBE) Statement, published in 2007[44]. STROBE is a 22-item checklist that details the key information needed when reporting the results of an observational study. STROBE is also accompanied by an “Explanation and Elaboration” document, providing further details and guidance for each checklist item giving examples of good reporting[45]. Because of the completeness and depth of its explanatory document, STROBE has been -improperly- used by researchers as an educational tool, both for the use of STROBE, but also for correct methodology of reporting[46]. Although STROBE has been widely used by researchers for more than 10 years, there is lack of evidence regarding the knowledge and the perception of researchers.

1.9. Other issues in poor reporting; using causal language when reporting observational studies

Another cause of inadequate reporting is related to the use of inappropriate terminology that may be methodologically incorrect or may mislead the readers.

Showing causal inference is the ultimate goal of most clinical research[50, 51]. It is generally accepted that causal inference could only be drawn from RCTs [47, 48]. This is because of the unique ability of RCTs to avoid confounding bias. This ability is granted by the randomization process, which allows a comparable distribution of measured and unmeasured confounding factors between experimental and control groups[49].

RCTs however cannot be performed in every occasion; in many settings RCTs may be not feasible, ethical or timely. For these reasons, researchers are often involved in observational studies and, despite the limitations of observational designs, authors may be tempted to draw causal conclusions in their reports[50, 51]. As we already discussed, the knowledge that association does not mean causation is well established within the medical community; therefore, authors may deliberately avoid causal statements on a hoped-for causal answer. Such writing strategy has been shown to bring ambiguity and contrived reporting [143, 144].

Researchers have proposed alternative ways to present observational research correctly; a suggestion is that authors should include in their report an open disclosure regarding the differences between the study designs and the likely distance in meaning and magnitude between their data-based estimates and the desired targeted causal effect [52, 53]. To promote similar efforts, Lederer et al. (2019)[145] recently published a guide for authors and editors on how to report causal studies in a specific field (Respiratory, Sleep and Critical Care). Their advices, similarly to other recent

authors, are in the direction of providing in the discussion key elements to ensure the validity of causal claims, rather than circumventing the problem by asking to avoid causal language[146]. An example that was given, involved multiple passages: briefly explaining what causal inference is, then provide a definition of a confounder, outline how to identify confounding through a directed acyclic graphs and finally discuss how p-values are often misinterpreted and how their value does not reflect the magnitude, direction or clinical importance of a given association. Such elements would empower the target audience to critically assess observational studies.

1.10. Rationale for the thesis

The thesis addresses different aspects related to research waste, exploring the RoB assessment and the reporting. The two core projects focus on the exploration of reasons of disagreements when assessing RoB for the same trials included in two different systematic reviews (SRs). A third project evaluates the association between RCT characteristics including RoB items (but also year of publication, sample size, trial registration status, centre status, region, funding source and conflict of interest) and the reported treatment effects of testosterone therapy in men using a meta-epidemiological approach. The other works explore the quality of reporting in observational studies, through the examination of the knowledge and adherence to reporting guidelines and the use of appropriate wording regarding causal claims in abstracts of observational studies.

For the core topic of this thesis, the RoB assessment, we vastly expanded current research in the issue. Although the low reproducibility of the RoB tool has been well demonstrated in previous studies, some important aspects such as the reasons of disagreements were not explored before. We decided to explore the uncertainty regarding the reproducibility of the RoB tool outside the experimental setting, which was rarely considered previously.

1.11. Thesis structure

The rest of this thesis is structured as follows:

Chapter 2 presents a research-on-research study evaluating the agreement on the RoB assessment when the same study was included in more than one Cochrane SRs and exploring the likely reasons in case of disagreements.

Chapter 3 presents a research-on-research study that continues from the results of the study reported in Chapter 2. In case of disagreements based on a different interpretation of the same information recovered from the source we examined if and how factors/characteristics that are not part of the RoB assessment may have influenced researchers in their assessment.

Chapter 4 presents a meta-epidemiological study that explores the association of certain trial characteristics, including RoB assessment, with treatment effect estimates in RCT of testosterone therapy in men.

Chapter 5 presents the results of a survey aimed to researchers involved in the field of observational research. The survey evaluated authors knowledge and use of the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement.

Chapter 6 presents a research on research study regarding the use of causal claims in observational studies, analysing how authors reports their finding in the abstracts sent to a prestigious journal and how this was mirrored in the full-text articles that were published.

Chapter 7 concludes the thesis by summarising the results, discussing the implications of the work presented in this thesis, and providing suggestions for future research.

Chapter 2: Reasons of disagreements in risk of bias assessment for RCTs assessed in two different Cochrane reviews

2.1. Background

Systematic reviews are synthesis of all the existing knowledge concerning a clinically relevant question. Assessing the methodological quality of individual studies is a crucial step when conducting a systematic review because if the individual studies are biased, the SR will be also biased. The tool developed by Cochrane (the RoB tool) has rapidly become the reference for assessing RoB in RCTs. The RoB tool is based on specific characteristics related to study design and conduct, selected on theoretical grounds and on empirical evidence from meta-epidemiological studies that these characteristics are associated with differences in treatment effect-estimates. Despite its rigorous development and the existence of guidelines on how to use it, previous studies showed that the reproducibility of assessments is often suboptimal. The reasons why disagreements in the assessment happened have not been explored in depth. For example, reviewers may base their judgment on additional information such as protocols or they may contact study authors. The aim of the project is to assess the frequency and reasons for disagreements in RoB assessment for RCTs included in more than one Cochrane SR.

2.2. Methods

This is a research-on-research study on RoB assessment using a dataset of recent Cochrane SRs to identify clinical trials that were assessed for RoB in two different SRs. For five key RoB items we evaluated agreement between two different reviews.

Data sources: All the 2,796 Cochrane reviews published between March 2011 and September 2014. All data regarding the SRs were retrieved in electronic format as XML files and were then used to retrieve the relevant data for the study. Reviews that were withdrawn or not containing any study were excluded.

Data selection: RCTs included in more than one review were identified among all selected reviews. The same studies were identified by matching them through the primary reference used to identify them in the SRs. A manual check of the results of the automatic matching was conducted. When more than two different SRs assessed RoB for a study, we kept only two reviews chosen at random. The situation concerned less than 10% of the total number of assessments.

Data extraction: RoB assessment in term of low, high and unclear and support for judgement for five key risk of bias items (Random Sequence Generation, Allocation Concealment, Blinding of

Participants and Personnel, Blinding of Outcome Assessment, Incomplete Outcome Data) was extracted. We checked the vocabulary related to RoB items and manually re-classified them when the terminology used was inconsistent (e.g. “allocation masking” instead of “allocation concealment”).

Data synthesis: For each item, we compared risk of bias assessment made in each review and calculated proportion of agreement. We calculated the percentage agreement for each risk of bias item, as the proportion of studies with a concordant assessment in both reviews (e.g. “low” risk of bias AND “low” risk of bias). Not all reviews assessed all five key risk of bias items for each RCT included; consequently, the number of RCTs evaluated for discrepancies varies depending on the item considered.

Two reviewers independently analysed 50% of all eligible RCTs identified in the previous step. The in-depth analysis was done by comparing support for each judgement with the information from the study report(s). The aim was to evaluate whether disagreements were related to a difference in information (e.g., one reviewer contacted the study author obtaining additional information) or a difference in interpretation (same support for judgement but different interpretation). Other possible reasons were “input mistake” (support for judgement contradict the judgement) and “unclear reason” if a reason could not be found. They also highlighted the main reasons for disagreements caused by different interpretation.

Statistical analysis: Analysis was descriptive with use of frequencies and percentages for qualitative variables.

2.3. Results

Figure 2 shows the selection process. From the 2,796 systematic reviews published between March 2011 and September 2014, 2,291 reviews included RCTs only and reported a risk of bias assessment. Of these, 797 included at least one RCT whose primary reference was shared with another review for which a risk of bias assessment was reported. These 797 reviews included 1,604 single RCTs evaluated for the same risk of bias item in more than one review. Proportion of agreement ranged from 57% (770/1,348 trials) for incomplete outcome data to 81% for random sequence generation (1,193/1,466) (Figure 3).

Classification of disagreements: The analysis was conducted on 802 studies selected at random. The most common source of disagreement was difference in interpretation of the same information, ranging from 65% (88/136) for random sequence generation to 90% (56/62) for blinding of

participants and personnel. Details are reported in Figure 4. The access to different or additional information explained 32/136 (24%) disagreements for random sequence generation and 38/205 (19%) for allocation concealment. Access to additional information was less common for the remaining items, with proportions ranging from 2% to 4%. In 80% of the cases, the access to additional information was obtained through the contact of the study author.

The other sources of disagreement we identified were less common; input mistake ranged from 1% to 6%, missed information from 1% to 6%. We could not determine the source of disagreement in 5% of our disagreements.

Main reasons of disagreements for different interpretation: The most common reason across items was related to incomplete or unclear reporting in the RCT. For random sequence generation, disagreements in 73/88 (83%) trials were related to lack of a precise description of the randomization process. For allocation concealment, the most common reason was due to the lack of description of the envelopes used to conceal allocation (17%, n=26/149 trials). For the two blinding items, a common reason was when an article mentioned only “double blind” without an additional description (16% of cases, n=9/56 trials for blinding of participants and personnel, 13%, n=9/70 for blinding of outcome assessment). For incomplete outcome data, a common reason was due to the use of formulas such as “no missing data” or “all data reported” (10%, 22/220 trials). Finally, a common reason for a difference in interpretation was confusion with another item. Allocation concealment was confused with blinding (10%, n=15/149 trials) but also with random sequence generation (4%, n=6/149). For blinding of participants and personnel, the most common cause for disagreement concerned the interpretation of cases when blinding was not feasible (36%, n=20/56 trials). For incomplete outcome data, the use of different cut-offs for the rate of missing data is the most common reason for disagreement (26%, n=57 trials).

Figure 2 Selection process

Figure 3 Agreement and different types of disagreement

Figure 4 Causes of disagreement

2.4. Discussion

Our results confirm that the agreement for risk of bias assessments is generally suboptimal, with better agreement for random sequence generation and allocation concealment and less agreement for incomplete outcome data. Access to different sources of information explained why 24% of the trials had disagreements in the assessment of risk of bias for random sequence generation and 19% for allocation concealment, while this reason was less common for other items. The main source of disagreements was a difference in interpretation of the same information, which was frequently related to incomplete or unclear reporting in the study report.

An implication of our study is that the agreement in risk of bias assessment would be enhanced by more detailed guidance in use of the RoB tool with focus on common causes of disagreements. We showed that in many cases, the unclear reporting from source material allows reviewers ample space for personal judgement and explained differences in judgement. A suggestion could be to give clearer instruction on how to evaluate common cases of incomplete reporting.

Following the suggestions based on the findings and comments provided by Jorgensen et al.[147] and Savovic et al.[148], Cochrane proposed a new version of the risk of bias tool, which has recently been released[56, 149]. The new version has a different approach to the risk of bias assessment, guiding reviewers through the process with the use of “signalling questions”, which

might leave less room for subjectivity. However, the new tool does not cover some of our concerns, especially those related to incomplete outcome data: quota for missing data that are considered acceptable, and whether reviewers should focus more on the reasons for the missing data or their magnitude. Research-on-research studies are needed to evaluate whether this new version of the tool results in improved reproducibility.

2.5. Article

Bertizzolo L, Bossuyt P, Atal I, Ravaud P, Dechartres A.

Disagreements in risk of bias assessment for randomised controlled trials included in more than one Cochrane systematic reviews: a research on research study using cross-sectional design.

BMJ Open. 2019 Apr 1;9(4):e028382.

doi: 10.1136/bmjopen-2018-028382.

BMJ Open Disagreements in risk of bias assessment for randomised controlled trials included in more than one Cochrane systematic reviews: a research on research study using cross-sectional design

Lorenzo Bertizzolo,^{1,2} Patrick Bossuyt,² Ignacio Atal,^{1,3} Philippe Ravaud,^{1,3,4,5,6} Agnes Dechartres⁷

To cite: Bertizzolo L, Bossuyt P, Atal I, et al. Disagreements in risk of bias assessment for randomised controlled trials included in more than one Cochrane systematic reviews: a research on research study using cross-sectional design. *BMJ Open* 2019;**9**:e028382. doi:10.1136/bmjopen-2018-028382

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-028382>).

Received 5 December 2018
Revised 14 February 2019
Accepted 15 February 2019

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to
Dr Lorenzo Bertizzolo;
lorenzo.bertizzolo@gmail.com

ABSTRACT

Objectives Assess the frequency and reasons for disagreements in risk of bias assessments for randomised controlled trials (RCTs) included in more than one Cochrane review.

Design Research on research study, using cross-sectional design.

Data sources 2796 Cochrane reviews published between March 2011 and September 2014.

Data selection RCTs included in more than one review.

Data extraction Risk of bias assessment and support for judgement for five key risk of bias items.

Data synthesis For each item, we compared risk of bias assessment made in each review and calculated proportion of agreement. Two reviewers independently analysed 50% of all disagreements by comparing support for each judgement with information from study report to evaluate whether disagreements were related to a difference in information (eg, contact the study author) or a difference in interpretation (same support for judgement but different interpretation). They also identified main reasons for different interpretation.

Results 1604 RCTs were included in more than one review. Proportion of agreement ranged from 57% (770/1348 trials) for incomplete outcome data to 81% for random sequence generation (1193/1466). Most common source of disagreement was difference in interpretation of the same information, ranging from 65% (88/136) for random sequence generation to 90% (56/62) for blinding of participants and personnel. Access to different information explained 32/136 (24%) disagreements for random sequence generation and 38/205 (19%) for allocation concealment. Disagreements related to difference in interpretation were frequently related to incomplete or unclear reporting in the study report (83% of disagreements related to different interpretation for random sequence generation).

Conclusions Risk of bias judgements of RCTs included in more than one Cochrane review differed substantially. Most disagreements were related to a difference in interpretation of an incomplete or unclear description in

Strengths and limitations of this study

- Use of a very large and comprehensive collection of Cochrane reviews to assess the agreement in risk of bias assessment and to understand reasons of disagreement.
- Analysis of the full text of study reports to underline what information was available to review authors and how they used them while assessing risk of bias.
- Focus on disagreements only. Possible that a proportion of agreements happened 'by chance'. For example, review authors may express the same risk of bias judgement while using different information or interpreting information differently.
- No evaluation of the potential impact of disagreements in conclusion making at the review level.

the study report. A clearer guidance on common causes of incomplete information may improve agreement.

INTRODUCTION

Systematic reviews aim to synthesise all existing evidence for a research question by the use of a rigorous and reproducible methodology.¹ Because reviews may be affected by bias at the level of individual studies,² an assessment of the risk of bias in these studies is a crucial step in conducting a systematic review.^{3,4}

Cochrane has developed a tool to provide a standardised approach to the assessment of the risk of bias in randomised controlled trials (RCTs).⁵ The risk of bias tool is based on specific characteristics related to study design and conduct, selected on theoretical grounds and on empirical evidence from

meta-epidemiological studies that these characteristics are associated with differences in treatment effect estimates.^{6–11} The tool includes seven items (random sequence generation, allocation concealment, blinding of participants and personnel, blinding of outcome assessment, incomplete outcome data, selective reporting, other source of bias), which the researchers assess and judge as either 'high', 'low' or 'unclear' risk of bias.^{11 12}

Although Cochrane provides detailed guidance on how to use the tool and recommends consensus between two independent reviewers,¹¹ personal judgement is also involved, which may bring variability. Several studies have evaluated the reproducibility of the risk of bias tool, generally shown to be poor.^{12–19} However, there is some uncertainty about the main causes of disagreements. For example, some reviewers may search for additional information such as protocols or contact study authors and this difference in available information, rather than a difference in judgement, may explain some of the disagreements.

In this study, we used a large collection of Cochrane reviews to evaluate the reproducibility of risk of bias assessments by identifying RCTs included in more than one Cochrane review and comparing the assessments. In addition, we examined the likely reasons for any disagreements. In particular, we evaluated whether disagreements were related to differences in information available to reviewers or differences in interpreting the same information and what could explain such different interpretation.

METHODS

This is a research on research study on risk of bias assessment, which used a cross-sectional design. We identified RCTs included in more than one reviews included in a large collection of Cochrane reviews. For key risk of bias items, we evaluated agreement between the different systematic reviews; analysed whether disagreements were related to a difference in information available to reviewers or a difference in interpretation of the same information and highlighted the main reasons for disagreements by an in-depth, one-by-one evaluation of disagreements.

Data sources

We obtained data from the 2796 Cochrane reviews, which correspond to all the reviews available in the Cochrane library between March 2011 and September 2014, including updates (March 2011 corresponds to the last update of the risk of bias tool⁵). Data consisted of one XML file per review, each file containing all data entered by review authors in RevMan, the software used for managing Cochrane reviews.²⁰ All individual XML files were merged in a single database by using R V.3.2.2²¹ with the XML package.²² The vocabulary used for risk of bias items slightly varied across reviews (eg, some reviews could refer to 'allocation concealment' as 'allocation masking'). For this reason, two authors independently

evaluated all terms used and classified them according to the vocabulary of the tool. Disagreements were resolved by consensus. This standardisation was done for a previous publication.²³

Selection of eligible reviews

We excluded withdrawn or 'empty' reviews (ie, systematic reviews not including any study) as well as reviews including observational or non-randomised studies and considered only reviews with an assessment of risk of bias for at least one item of the risk of bias tool.

Selection of eligible RCTs

To identify single RCTs included and assessed for risk of bias in more than one systematic review, we proceeded as follows. For each RCT, we identified the primary reference(s), which was the reference identified by review authors as the main reference(s) for an included study. Then, we used a matching algorithm²⁴ to identify studies that shared the same primary reference. If several primary references were reported, we considered all of them. We manually checked that the studies sharing the same primary reference in the reviews corresponded to the same RCT.

Extraction of risk of bias assessment

For each eligible RCT, we extracted the risk of bias assessment and the corresponding support for judgement for each risk of bias item in each review. Whenever a single RCT was included in three or more reviews, we considered only the risk of bias assessment from two reviews chosen at random; this was decided because of workload and to facilitate direct comparison of two assessments and concerned less than 10% of our included RCTs. We focused on five risk of bias items: random sequence generation, allocation concealment, blinding of participants and personnel, blinding of outcome assessment and incomplete outcome data. We did not consider selective reporting because it is difficult to evaluate in the absence of the study protocol, which is frequently lacking, especially for older studies.^{11 12 14} We also did not consider the item other bias because the definition is very wide (ie, 'any important concerns about bias not covered in the other domains in the tool'¹¹), so comparisons across reviews are difficult.

Comparison of risk of bias assessment between reviews

For each item, we compared the risk of bias assessment in terms of 'high', 'low' or 'unclear' risk of bias between the two reviews. According to the Cochrane handbook, the items blinding of outcome assessment and incomplete outcome data should be assessed for each outcome. Therefore, when the reviews reported an assessment of these items at the outcome level, we manually checked that outcomes were identical in both reviews and we retained for our analysis only the assessments that focused on the same outcomes. For blinding, we followed the last version of the Cochrane handbook and we retained only assessments of blinding of participants and personnel

and blinding of outcome assessment as two independent items, excluding different types of assessment (ie, blinding as a single item, blinding of only participants or of only personnel).

We calculated the percentage agreement for each risk of bias item, as the proportion of studies with a concordant assessment in both reviews (eg, 'low' risk of bias AND 'low' risk of bias). Not all reviews assessed all five key risk of bias items for each RCT included; consequently, the number of RCTs evaluated for discrepancies varies depending on the item considered.

Selection of studies for in-depth analysis of disagreements

For workload reasons, we in-depth evaluated the reasons for disagreements for 50% of the studies analysed in the previous step. In cases of more than one shared RCTs within a given pair of Cochrane reviews, we selected only one RCT at random. To reach 50% of the total sample, we used a simple random selection in the remaining database.

Classification of disagreements

For the random selection, two reviewers (LB and AD) independently evaluated all disagreements in the risk of bias assessment in the two systematic reviews. They first scrutinised the support for the judgement in each review and evaluated whether it was the same or 'conceptually' the same in both reviews (eg, 'randomised, probably done'; 'randomised, probably not done'; 'study only mentions randomisation, but does not specify how randomisation was performed; unclear'; 'study states it is randomised; low risk'). If the support differed, they assessed any other information regarding the study as reported in both reviews, systematically searching and evaluating the full-text study report indicated in the primary reference. A formalised data extraction process for full texts was not used. Full texts were examined, looking primarily for correspondence between information reported by the reviewers in their support for judgement and the text.

They independently classified each case of disagreement as follows:

- ▶ Disagreement related to differences in interpretation:
 - The support for judgement was the same (or 'conceptually' the same) in both reviews, but the interpretation differed.
 - One review clearly confused one item of the risk of bias tool with a different one or the review authors misunderstood the definition of the item (eg, for random sequence generation, support for judgement reports '600 opaque envelopes, 1 was drawn every time').
- ▶ Disagreement related to differences in information: the support for judgement cites information that is not available in the study report; additional sources are cited (eg, protocol) or the review authors reported that they had contacted the RCT author for additional data.

- ▶ Disagreement related to information missed by the review authors: the study report clearly describes the information, but some review authors seemed to have missed this information in the study report.
- ▶ Disagreement related to input mistakes: risk of bias assessment in terms of 'high'/'low'/'unclear' did not match the support for the judgement (eg, 'randomisation described explicitly', judgement 'unclear').
- ▶ Unclear: when it was not possible to classify the disagreement because the support for the judgement was empty or because we could not retrieve the full-text study report.

Any disagreements between reviewers were solved by discussion to reach consensus. In the online supplementary appendix 1, we report a figure synthesising how the in-depth analysis process was conducted.

Identification of main reasons for different interpretation

For each disagreement related to a difference in interpretation, we evaluated the probable reason for disagreement. For example, the interpretation could differ because of confusion with another risk of bias item (eg, random sequence generation and allocation concealment) or because the information was unclear or insufficiently detailed in the article. When we were unsure about the reason, we classified the reason as unclear. Two authors (LB and AD) conducted this process in duplicate by using all available information (ie, support for the judgement, characteristics of the study reported in the review, full-text article), with disagreements resolved by discussion.

Statistical analysis

Analysis was descriptive with use of frequencies and percentages for qualitative variables. Statistical analysis was conducted with Stata V.13.1.²⁵ We decided to use simple per cent agreement because other static approaches were problematic. The Kappa statistic requires having defined reviewers, which is not the case of our approach. Another statistic, the intraclass correlation coefficient is not suitable, because it requires assessments to be in an ordinal order, which is not our case. There is no continuum between the assessments of low, unclear and high risk of bias.

Patient involvement

Patients were not involved in any aspect of the study design, conduct or the development of the research question or outcome measures. This is a research-on-research study, and therefore, there was no active patient recruitment for data collection.

RESULTS

Selection process

Figure 1 shows the selection process. From the 2796 systematic reviews published between March 2011 and September 2014, 2291 reviews included RCTs only and

Figure 1 Flow chart of the selection process. RCT, randomised controlled trial.

reported a risk of bias assessment. Of these, 797 included at least one RCT whose primary reference was shared with another review for which a risk of bias assessment was reported. These 797 reviews included 1604 single RCTs evaluated for the same risk of bias item in more than one review. The online supplementary appendix 2 reports the frequency of the different Cochrane groups among those reviews.

Among the 1604 selected RCTs 1603 had duplicate evaluation for allocation concealment, 1466 for random sequence generation, 375 for blinding of participants and personnel, 583 for blinding of outcome assessment and 1348 for incomplete outcome data.

Evaluation of agreement and distribution of disagreements

The agreement of risk of bias judgements ranged from 57% (770/1348 trials) for incomplete outcome data to 81% (1193/1466 trials) for random sequence generation (figure 2). We identified most disagreements for 'low' and 'unclear' risk of bias judgements, especially for random sequence generation (231/273 trials, 85%). Disagreements between 'low' and 'high' risk of bias were generally rare, for example 8/273 of disagreements (3%) for random sequence generation, with the exception of incomplete outcome data for which they were more frequent (190/578, 33%). For blinding of participants and personnel, the most frequent disagreement was

between 'unclear' and 'high' risk of bias (50/107, 47%), then 'low' versus 'unclear' (34/107, 32%), and 'low' versus 'high' (23/107, 21%) (figure 2).

Classification of disagreements

The in-depth analysis of disagreements included 802 studies: 799 for allocation concealment, 747 for random sequence generation, 206 for blinding of participants and personnel, 297 for blinding of outcome assessment and 660 for incomplete outcome data. The agreement results of this sample and the distribution of disagreements are reported in the online supplementary appendix 3.

For all items, the most common source of disagreement was a difference in interpretation, with frequencies ranging from 88/136 (65%) for random sequence generation to 56/62 (90%) for blinding of participants and personnel (figure 3). The access to additional or different information accounted for disagreements in 32/136 (24%) trials for random sequence generation and 38/205 (19%) for allocation concealment. Access to additional information was less common for the remaining items, with proportions ranging from 2% to 4%. In 80% of the cases, the access to additional information was through the contact of the study author.

The other sources of disagreement were less common; input mistake ranged from 1% to 6%, missed information from 1% to 6%. We could not

Figure 2 Distribution of agreements and disagreements for the different risk of bias items analysed; raw number and percentage of the total. For disagreements, distribution of the different discrepancies.

determine the source of disagreement in 5% of our disagreements. For this analysis, we accessed the full text of 216 different trials to help us in the process. The online supplementary appendix 4 reports some examples of disagreements in which the access to the study report helped us in the classification and

the analysis of reasons of disagreement. We could not retrieve or access 19 full texts we deemed necessary for the categorisation of disagreements and this explains the majority of cases where we were unable to categorise the source of disagreement ('unclear' source in figure 3).

Figure 3 Classification of disagreements for the different items; raw number and percentage of the total.

Main reasons of disagreements for different interpretation

The main reasons for a difference in interpretation for each item are reported in [table 1](#). Additional examples are provided for each item for the high-low disagreements (online supplementary appendix 5). The most common reason across items was related to incomplete or unclear reporting in the RCT. For random sequence generation, disagreements in 73/88 (83%) trials were related to lack of a precise description of the randomisation process with reviewers evaluating 'low', 'high' or 'unclear' risk of bias the reporting of 'randomised' in the text. For allocation concealment, the most common reason for disagreement was a different interpretation of description of the envelopes used to conceal allocation (17%, n=26/149 trials). For the two blinding items, many disagreements occurred when the article mentioned only 'double blind' in RCTs without an additional description (16% of cases, n=9/56 trials for blinding of participants and personnel, 13%, n=9/70 for blinding of outcome assessment). For incomplete outcome data, reviewers assessed differently the statement from the study report of 'no missing data' or 'all data reported' (10%, 22/220 trials). Another common reason for a difference in interpretation was confusion with another item. Allocation concealment was confused with blinding (10%, n=15/149 trials) but also with random sequence generation (4%, n=6/149). For blinding of participants and personnel, the most common cause for disagreement concerned the interpretation of cases when blinding was not feasible (36%, n=20/56 trials), assessed at high risk by some reviewers and low by others. Another common cause of disagreement for the two blinding items related to the assessment of outcomes that should not be affected by blinding (eg, mortality); it explained 21% (n=12 trials) of disagreements for blinding of participants and personnel and 23% (n=16 trials) for blinding of outcome assessment, often low versus high disagreements.

For incomplete outcome data, the use of different cut-offs for the rate of missing data is the most common reason for disagreement (26%, n=57 trials); also common is considering the explanation of reasons for missing data enough to attribute a low risk of bias (13%, n=28 trials).

DISCUSSION

In this study, we took advantage of a very large sample of Cochrane reviews to explore the sources of disagreements in risk of bias assessment for trials included in several reviews. We decided to focus on Cochrane reviews because as these reviews are produced within a single organisation, therefore, we expected results and procedures to be more appropriately comparable. Authors compiling Cochrane reviews are members of the organisation and, in most cases, they underwent a similar training for assessing risk of bias. Our results confirm that the agreement for risk of bias assessments is generally suboptimal, with better agreement for random sequence generation and allocation concealment and less agreement for incomplete

outcome data. Access to different sources of information explained why 24% of the trials had disagreements in the assessment of risk of bias for random sequence generation and 19% for allocation concealment. However, the main source of disagreements was a difference in interpretation of the same information, which was frequently related to incomplete or unclear reporting in the study report.

Strengths and weaknesses

Our study goes beyond previous literature on the topic.^{3 12–18 26} As compared with most other studies,^{12–17} we used real-world data to explore agreement of risk of bias assessments in real scenarios. We evaluated a very large and comprehensive collection of Cochrane reviews that spanned multiple specialties and topics, including a number of trials about 10 times larger than the largest study on the topic.¹² We completed our analysis by searching individual study reports to give support to our comments on reasons for disagreements, which, to our knowledge, has not been done in previous, smaller works that used a similar methodology.¹⁸ While doing this, we developed a suitable classification scheme for sources of disagreements and conducted, in duplicate, an extensive analysis to understand the risk of bias assessment process and explored the most common reasons for disagreements.

Our study has limitations. Whenever a single RCT was included in three reviews or more, we considered only the risk of bias assessment from two reviews chosen at random. Nevertheless, we cannot exclude that different combinations of two chosen evaluations could have produced slightly different results. Although the classification of disagreements was conducted in duplicate following a formalised process, there remains a component of personal judgement. We evaluated only disagreements, but a number of agreements might have occurred 'by chance'. In our analysis of likely reasons for disagreements, some resulted from confusion between risk of bias items. Similar discrepancies might have occurred among agreements; indeed, previous literature on the topic demonstrated that reviewers do not accurately follow the risk of bias tool.²⁷ We also did not assess the selective reporting item that is frequently judged on incomplete information. We did not evaluate whether disagreements varied depending on the Cochrane review group or year of publication. Finally, we did not evaluate the impact of disagreements and the extent to which the evidence base for making conclusions and providing summary statements of effectiveness may have been affected by changing the rating.

Comparison with other studies

Our findings confirm the importance of issues that were previously identified by Jørgensen *et al*⁸ and Savović *et al*.²⁶ In particular, Savović *et al*,²⁶ surveying users of the risk of bias tool, reported on the possibility of confusion between random sequence generation and allocation

Table 1 Main reasons for disagreements in cases of a different interpretation of the same information

Risk of bias Item	Main reasons for disagreements	N (%)	Examples of support for judgement from the review†
Random sequence generation	Consider differently incomplete or unclear description	73 (83)	'States 'cluster randomisation by computer'; Low risk of bias 'Cluster randomisation by computer. No further information provided'; Unclear risk of bias
	Confusion with allocation concealment	9 (10)	'Allocation was done using sealed envelopes containing name of one of the two groups.'; Low risk of bias
Allocation concealment	Consider differently incomplete or unclear description	49 (33)	'Not specified.'; High risk of bias 'Method of concealment not described.'; Unclear risk of bias
	Consider differently envelopes description	26 (17)	'Sequentially numbered sealed envelopes'. Does not state if opaque envelopes.'; Unclear risk of bias 'Sequentially numbered sealed envelopes.'; Low risk of bias
	Random sequence generated by computer or external centre considered enough for low risk	21 (14)	'Treatment was allocated based on the computer-generated no list.'; Low risk of bias
	Confusion in the definition of the item	19 (13)	'Researchers attempted to contact all patients seen by physicians during 1 month'; High risk of bias
	Confusion with blinding	15 (10)	'Participants were told to which compound they had been allocated.'; High risk of bias
	Confusion with random sequence generation	6 (4)	'Computer-generated randomised lists.'; Low risk of bias
Blinding of participants and personnel	Assess risk differently if blinding was not feasible because of the type of intervention	20 (36)	'Not possible to blind participants'; Low risk of bias 'Participants were not blinded for provided treatment. This is inherent to study design'; High risk of bias
	Outcome considered not influenced by blinding	12 (21)	'No information given about whether patients were blind to physician allocation but treatment outcomes judged unlikely to be affected by lack of blinding'; Low risk of bias
	Consider differently information of 'double blind'	9 (16)	'Quote: '... patients were randomised in double-blind conditions ...'. Comment: probably done'. Low risk of bias 'Quote: 'double blind conditions'. No further details'; Unclear risk of bias
	Consider differently incomplete or unclear description	7 (12)	'Researchers were blind until after the baseline assessment. participants were not blinded.'; Unclear risk of bias 'Not possible to blind participants to intervention. Insufficient information to make a judgement about blinding of therapists'. High risk of bias
Blinding of outcome assessment	Confusion in the definition of the item	5 (9)	'Described as an 'open-label' pilot study.'; Low risk of bias
	Consider differently incomplete or unclear description	24 (34)	'Not explicitly discussed in the publish study, it was assumed to be open label'; High risk of bias 'Not described in published study'; Unclear risk of bias
	Outcome considered not influenced by blinding	16 (23)	'Not stated, but it was unlikely that the outcome was influenced by lack of blinding'; Low risk of bias
	Consider differently patient-reported outcomes when patients are blinded or not to the intervention	9 (13)	'Comment: depression assessed by patient self-report'; High risk of bias 'Insufficient information available to assess'; Low risk of bias
	Consider differently information of 'double blind'	9 (13)	'Quote: '... double blind' Comment: probably done'; Low risk of bias 'Quote: 'double blind conditions'. No further details'; Unclear risk of bias
	Assess risk differently if blinding was not feasible because of the type of intervention	6 (9)	'Blinding not possible due to intervention'; High risk of bias 'Unclear blinding of outcome assessment'; Low risk of bias

Continued

Table 1 Continued			
Risk of bias item	Main reasons for disagreements	N (%)*	Examples of support for judgement from the review†
Incomplete outcome data	Use different cut-off for the rate of missing data	57 (26)	'11 withdrawals (10%); Low risk of bias 'Comment: there were post-randomisation drop-outs; High risk of bias
	Focus on no versus reasons/precise report of missing data	28 (13)	'20 drop-outs (27.2%) with 4 deaths (3 males, 1 female) from cardiovascular events; High risk of bias 'Numbers and reasons for drop-outs and withdrawals in all intervention groups were described; Low risk of bias
	Consider differently incomplete or unclear description	27 (12)	'Women who were untraceable or unsuitable for follow-up were excluded, other losses included as smokers; Low risk of bias '167/1287 (12.9%) (C=83, I=84) excluded from analysis due to moving away, being untraceable or deemed unsuitable for follow-up (eg, miscarriage). 1120 in sample. 51/1287 non-responders were included as continuing smokers; High risk of bias
	Consider differently intention-to-treat (ITT) analysis	25 (11)	'147 randomised; 14 in the letrozole group and 3 in the laparoscopic ovarian drilling dropped out of the trial, all for non-compliance. However, ITT analysis was not conducted; Unclear risk of bias '7 women lost to follow-up, but similar (3 vs 4) in both groups; losses due to non-compliance; Low risk of bias
	Consider differently report of 'no missing data'	22 (10)	'Did not report number of withdrawals. Comment: all patients who were randomised were included in the final analysis. ITT analysis was conducted; Unclear risk of bias 'It does not appear that there were any withdrawals or drop-outs; Low risk of bias
	Consider differently imputation of missing data	20 (9)	'Imputation method not described; Unclear risk of bias 'Drop-out rate was not significant; Low risk of bias
	Use different cut-off for difference in the rate missing data between different arms/comparisons	13 (6)	'Drop-out higher in placebo group (35% vs 25% in budesonide group). ITT used; High risk of bias 'Similar rates of withdrawal between arms. Withdrawals: 36 BUD, 51 placebo; Low risk of bias

*Number of RCTs disagreeing for this reason; percentage over the total of disagreements for different interpretation.

†When two extracts are reported, they refer to the same study.
RCT, randomised controlled trial.

concealment and between allocation concealment and blinding; the uncertainty on how to address unfeasibility of blinding; and the difficulties in assessing incomplete outcome data especially regarding the acceptable rate of missing data. More recently, Jørgensen *et al.*,³ evaluating comments on the use of the risk of bias tool, highlighted how authors complained that judgement often originates from incomplete or missing information.

A previous study identified 46 RCTs included in different systematic reviews in the field of fertility and evaluated the percentage agreement in risk of bias assessment. That analysis showed generally worse agreement than in our study, with percentage agreement ranging from 35% to 71%. Differences in sample size and the particular topic may explain these discrepancies. In addition, although the authors had compared supports for judgement between reviews, this evaluation may have been incomplete, because they did not evaluate the primary study reports.¹⁸

Implications

Our results confirm that the agreement in risk of bias assessment would be enhanced by more detailed guidance in use of the risk of bias tool with particular focus on common causes of disagreements. We showed that in many cases, the unclear reporting from source material allows reviewers ample space for personal judgement and differences in judgement.

The scientific community continues to stress the importance of improving the reporting of trials,^{28–31} which may limit disagreements when assessing risk of bias. In parallel, we could also work on restricting the space for personal interpretation when assessing risk of bias. A suggestion could be to give clearer instruction on how to evaluate common cases, for example, when confronted with nothing more than the term ‘randomised’ or ‘double blind’ in the study report. Similarly, a threshold could be set on the quota for missing data and indications on which imputation methods are appropriate and in which situations.

To minimise research waste, it could be interesting to have access to risk of bias assessments from other Cochrane groups and the supports they used, including information from authors or from protocols to help reviewers in their assessments. This process would imply having a unique study identification number across reviews and a central shared repository for all studies included in any Cochrane reviews.

Following the suggestions based on the findings and comments provided by Jørgensen *et al.*³ and Savović *et al.*,²⁶ Cochrane has been working on a new version of the risk of bias tool, which has recently been released.^{32–33} The new version has a different approach to the risk of bias assessment, guiding reviewers through the process with the use of ‘signalling questions’, which might leave less room for subjectivity. In addition, there is more guidance in assessing some items. For example, the new tool better clarifies some aspects of the randomisation process,

especially about what to do in some cases of incomplete information (eg, randomisation list created by an external centre with no other indication). The new tool also has a different approach to the blinding aspect, oriented to the implications of the masking process. However, the new tool does not cover some of our concerns, especially those related to incomplete outcome data: quota for missing data that are considered acceptable, and whether reviewers should focus more on the reasons for the missing data or their magnitude. It also does not address the common case of authors reporting ‘no missing data’. Research-on-research studies are needed to evaluate whether this new version of the tool results in improved reproducibility.

CONCLUSION

This analysis of risk of bias assessment for more than 1600 trials included in more than one reviews showed that agreement remains suboptimal. Most disagreements come from a difference in interpretation of an incomplete or unclear description in the study report. In some cases, the difference in the assessment was due to some but not all review authors obtaining additional information, from a protocol or from contacting study author.

Author affiliations

¹U1153, Epidemiology and Biostatistics Sorbonne Paris Cité Research Center (CRESS), Methods of therapeutic evaluation of chronic diseases team (METHODS), INSERM, Paris, Île-de-France, France

²Department of Clinical Epidemiology and Biostatistics, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands

³Cochrane France, Paris, France

⁴Centre d'Épidémiologie Clinique, INSERM U738 Hôpital Hôtel Dieu AP-HP (Assistance Publique des Hôpitaux de Paris), Paris, France

⁵Faculté de Médecine, Université Paris Descartes, Sorbonne Paris Cité, Paris, France

⁶Department of Epidemiology, Columbia University, Mailman School of Public Health, New York, USA

⁷Sorbonne Université, INSERM, Institut Pierre Louis d'Epidémiologie et de Santé Publique, AP-HP, Hôpitaux Universitaires Pitié Salpêtrière - Charles Foix, Département Biostatistique Santé Publique et Information Médicale, F75013, Paris, France

Acknowledgements We thank Camila Olarte Parra for her help during the data management phase and her comments on this manuscript. We thank David Tovey, editor in chief of the Cochrane Library, for agreeing to share data from Cochrane reviews; Javier Mayoral Campos, system administrator; the Cochrane Central Executive for preparing files; and all Cochrane reviewers who collected data. We also thank Laura Smales for English revision of the manuscript.

Contributors LB was involved in the study conception, selection of trials, data extraction, data analysis, interpretation of results and drafting the manuscript. PB was involved in the study conception, data analysis, interpretation of results and drafting the manuscript. IA was involved in the study conception, data extraction and drafting the manuscript. PR was involved in the study conception and drafting the manuscript. AD was involved in the study conception, selection of trials, data extraction, data analysis, interpretation of results and drafting the manuscript.

Funding This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement no 676207.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Raw data and analyses are available on request from the authors.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

1. Cook DJ, Mulrow CD, Haynes RB. Systematic reviews: synthesis of best evidence for clinical decisions. *Ann Intern Med* 1997;126:376–80.
2. Hopewell S, Boutron I, Altman DG, et al. Incorporation of assessments of risk of bias of primary studies in systematic reviews of randomised trials: a cross-sectional study. *BMJ Open* 2013;3:e003342.
3. Jørgensen L, Paludan-Müller AS, Laursen DR, et al. Evaluation of the Cochrane tool for assessing risk of bias in randomized clinical trials: overview of published comments and analysis of user practice in Cochrane and non-Cochrane reviews. *Syst Rev* 2016;5:80.
4. Hrobjartsson A, Boutron I, Turner L, et al. Assessing risk of bias in randomised clinical trials included in Cochrane Reviews: the why is easy, the how is a challenge. *Cochrane Database Syst Rev* 2013;4:ED000058.
5. Higgins JP, Altman DG, Gotzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011;343:d5928.
6. Schulz KF, Chalmers I, Hayes RJ, et al. Empirical evidence of bias. Dimensions of methodological quality associated with estimates of treatment effects in controlled trials. *JAMA* 1995;273:408–12.
7. Moher D, Pham B, Jones A, et al. Does quality of reports of randomised trials affect estimates of intervention efficacy reported in meta-analyses? *Lancet* 1998;352:609–13.
8. Balk EM, Bonis PA, Moskowitz H, et al. Correlation of quality measures with estimates of treatment effect in meta-analyses of randomized controlled trials. *JAMA* 2002;287:2973–82.
9. Page MJ, Higgins JP, Clayton G, et al. Empirical Evidence of Study Design Biases in Randomized Trials: Systematic Review of Meta-Epidemiological Studies. *PLoS One* 2016;11:e0159267.
10. Dechartres A, Trinquart L, Faber T, et al. Empirical evaluation of which trial characteristics are associated with treatment effect estimates. *J Clin Epidemiol* 2016;77:24–37.
11. Higgins JPT, Altman DG, Sterne JAC. Chapter 8: Assessing risk of bias in included studies. In: Higgins JP, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions. Version 5.1.0 (updated March 2011)* ed: The Cochrane Collaboration, 2011. www.handbook.cochrane.org.
12. Hartling L, Ospina M, Liang Y, et al. Risk of bias versus quality assessment of randomised controlled trials: cross sectional study. *BMJ* 2009;339:b4012.
13. Hartling L, Bond K, Vandermeer B, et al. Applying the risk of bias tool in a systematic review of combination long-acting beta-agonists and inhaled corticosteroids for persistent asthma. *PLoS One* 2011;6:e17242.
14. Hartling L, Hamm MP, Milne A, et al. Testing the risk of bias tool showed low reliability between individual reviewers and across consensus assessments of reviewer pairs. *J Clin Epidemiol* 2013;66:973–81.
15. Armijo-Olivo S, Ospina M, da Costa BR, et al. Poor reliability between Cochrane reviewers and blinded external reviewers when applying the Cochrane risk of bias tool in physical therapy trials. *PLoS One* 2014;9:e96920.
16. Armijo-Olivo S, Stiles CR, Hagen NA, et al. Assessment of study quality for systematic reviews: A comparison of the cochrane collaboration risk of bias tool and the effective public health practice project quality assessment tool: Methodological research. *J Eval Clin Pract* 2012;18:12–18.
17. da Costa BR, Beckett B, Diaz A, et al. Effect of standardized training on the reliability of the Cochrane risk of bias assessment tool: a prospective study. *Syst Rev* 2017;6:44.
18. Jordan VM, Lensen SF, Farquhar CM. There were large discrepancies in risk of bias tool judgments when a randomized controlled trial appeared in more than one systematic review. *J Clin Epidemiol* 2017;81:72–6.
19. Wilkins AJ. Risk of bias in assessing Risk of Bias. *Ophthalmic Physiol Opt* 2017;37:107–9.
20. Review Manager (RevMan) [Computer program] [program]. Version 5.3 version. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.
21. R: A language and environment for statistical computing. [program]. Vien: R Foundation for Statistical Computing, 2013.
22. XML: Tools for Parsing and Generating XML Within R and S-Plus. [program]. 3.2.2 version, 2017.
23. Dechartres A, Trinquart L, Atal I, et al. Evolution of poor reporting and inadequate methods over time in 20920 randomised controlled trials included in Cochrane reviews: research on research study. *BMJ* 2017;357:j2490.
24. deMelo VV. *Conference: Advances in Logic Based Intelligent Systems*, 2005.
25. *Stata Statistical Software: Release 13* [program]. College Station, TX: StataCorp LP, 2013.
26. Savović J, Weeks L, Sterne JA, et al. Evaluation of the Cochrane Collaboration's tool for assessing the risk of bias in randomized trials: focus groups, online survey, proposed recommendations and their implementation. *Syst Rev* 2014;3:37.
27. Propadalo I, Tranfic M, Vuka I, et al. In Cochrane reviews, risk of bias assessments for allocation concealment were frequently not in line with Cochrane's Handbook guidance. *J Clin Epidemiol* 2019;106.
28. Shamseer L, Hopewell S, Altman DG, et al. Update on the endorsement of CONSORT by high impact factor journals: a survey of journal "Instructions to Authors" in 2014. *Trials* 2016;17:301.
29. Turner L, Shamseer L, Altman DG, et al. Does use of the CONSORT Statement impact the completeness of reporting of randomised controlled trials published in medical journals? A Cochrane review. *Syst Rev* 2012;1:60.
30. Turner L, Shamseer L, Altman DG, et al. Consolidated standards of reporting trials (CONSORT) and the completeness of reporting of randomised controlled trials (RCTs) published in medical journals. *Cochrane Database Syst Rev* 2012;11:MR000030.
31. Altman DG, Moher D, Schulz KF. Improving the reporting of randomised trials: the CONSORT Statement and beyond. *Stat Med* 2012;31:2985–97.
32. RoB 2.0 Tool. <https://sites.google.com/site/riskofbiastool/welcome/rob-2-0-tool> (Accessed 22 Aug 2018).
33. Higgins JP, Sterne JA, Savovic J, et al. A revised tool for assessing risk of bias in randomized trials. *Cochrane Database of Systematic Reviews* 2016.

Chapter 3: Influence of external factors on RoB assessment

3.1. Background

In the previous project (17), we evaluated the reasons for disagreements in risk-of-bias assessment of trials included in several Cochrane reviews. Our results confirmed suboptimal agreement. More than two thirds of disagreements could be related to differences in interpretation of the same information and these disagreements were mostly due to incomplete or unclear information in the published trial report (17).

Although an “unclear” judgement can be expected in case of incomplete or unclear information, authors may consider the information adequate to express a judgement of low or high RoB.

Erroneous decisions apart, the RoB tool clearly leaves ample space for personal interpretation and judgement of evidence from study reports.

We wondered whether contextual factors such as characteristics related to the review team or the review itself, might be associated with differences in RoB when evaluating the same incomplete or unclear information. We aimed to explore which factors, if any, are associated with high or low risk of bias judgement in case of incomplete or unclear information in reports of primary studies for which an “unclear” risk of bias judgment would have been expected.

3.2. Methods

This is a research-on-research study on RoB assessment using a large collection of published Cochrane Systematic Reviews.

Study Design: The study followed a matched case–control design with a sample of RCTs for which the risk of bias was assessed differently in two separate Cochrane reviews.

Data Sources: all Cochrane reviews published or updated between March 2011 and September 2014. The same set was used for the study reported in Chapter 2 (17).

Selection of eligible studies: trials for which we observed a disagreement between the two reviews and attributed this disagreement to a difference in interpretation related to incomplete or unclear information in the study report in our previous study.

Definition of cases and controls: A case was defined as a judgement of “high risk of bias” or “low risk of bias” for an RCT in one of the reviews. A matched control was defined as a judgement of “unclear risk of bias” for the same RCT, for the same item, in the second, different review. Cases and controls were identified for each item.

Data extraction: We collected information regarding characteristics of reviews, review authors and studies included in the review.

Statistical Analysis: We used a conditional logistic regression model, accounting for the matched design. The results are expressed as odds ratios (OR) with 95% confidence intervals (CI) for each factor, considering separately disagreements of an assessment of high versus unclear risk of bias and low versus unclear risk of bias. Quantitative variables were analysed as such, without dichotomization. We conducted a single joint analysis across different items, then subgroup analyses for each item separately. We did not adjust for multiple comparisons.

3.3. Results

Selection process: In our previous project (17) 402 disagreements were attributed to a different interpretation of the same incomplete or unclear information. These 402 disagreements related to 304 different RCTs, included in 376 different reviews.

Characteristics of included reviews: The 376 reviews were compiled by 49 different Cochrane review groups, with the largest number (45; 12%) from the “pregnancy” group. More than half of the reviews (n=204, 54%) evaluated only non-pharmacological treatments. The number of RCTs included in reviews varied from 1 to 171 (median 16); only one quarter of reviews had included more than 30 RCTs. Overall, 147 reviews (39%) reported a conflict of interest among authors of the review.

Characteristics of the disagreements: Across the different risk of bias items, 69% (n=278) of disagreements were between low and unclear judgements; 31% (n=124) were between high and unclear judgements. Among the 402 disagreements, there were 278 disagreements (69%) between low and unclear judgements and 124 (31%) between high and unclear. Most disagreements derived from two judgements expressed by the same Cochrane group. This occurred in two thirds of disagreements (n=261, 65%). In 376 of the 804 (47%) single judgements analysed, the most common judgement for that item in the review was “unclear” risk of bias. For only 15 judgements (2%) one of the authors of the systematic review was also among the authors of the RCT being evaluated.

Analysis of factors associated with disagreements: Among the low versus unclear disagreements, review groups and authors that had completed a lower number of reviews assigned significantly more often a low risk of bias where others reported “unclear” (OR 0.97 [95% CI 0.95-0.99] and OR 0.97 [95% CI 0.95-0.998], respectively).

Among the low risk of bias judgements, there were significantly a lower number of RCTs per review (OR 0.99 [95% CI 0.99-0.998]) and more reviews in which one of the authors had also co-authored a trial report compared to controls with unclear assignments (OR 11.71 [1.39-98.76]). There were also significantly fewer recent RCTs that were assigned a low risk of bias (OR 0.37 [0.15-0.96]). A similar difference, although not significant, was observed for including larger trials (OR 0.37 [0.13-1.02]).

Among the low risk of bias judgements there were significantly fewer reviews that had assigned the majority of RCTs in their review at an unclear risk of bias (OR 0.12 [0.07-0.19]). Similarly, among the high risk of bias judgements in the high vs unclear comparison, there were significantly fewer reviews that had evaluated the majority of RCTs at an unclear risk of bias (OR 0.11 [0.05-0.24]).

3.4. Discussion

We evaluated factors or characteristics that are potentially associated with differences in RoB assessments of RCTs included in systematic reviews, in case of incomplete or unclear information given in reports. The number of reviews conducted by the group and by the first author, as well as the publication year of the RCT relative to the other RCTs included in the review, were each associated with different risk-of-bias judgments.

We observed that judgements of unclear risk of bias more frequently came from reviews that judged most of the RCTs included in the review at unclear risk. One possible explanation is that when the risk-of-bias assessment of a study is not completely straightforward (hence resulting in a disagreement between two different reviews), the reviewers may tend to go back to the judgement they used for most of the RCTs they assessed.

An interesting point is that two thirds of the disagreements originated within the same Cochrane review group. Nevertheless, we also showed that a higher number of reviews conducted by review groups and by authors, thereby suggesting more experience with risk of bias assessment, seems to have influenced reviewers in their judgment. This finding could suggest that the level of training may differ across author groups, or it may be influenced by the experience of the reviewers.

Our results indicate that identifiable review and author factors may play a role in risk of bias judgements of RCTs in systematic reviews. Specific factors that influenced reviewers in this analysis were: the number of reviews they previously conducted, and the number authored by the review group to which they belong. Reviewers also tended to judge a study at unclear risk of bias if they already did so for most of the studies included in the same review. Awareness of contextual

characteristics that can affect risk of bias judgements may help researchers in conducting more reproducible and meaningful assessments of risk of bias.

3.5. Article

Bertizzolo L, Bossuyt PM, Atal I, Ravaud P, Dechartres A.

External factors may influence Cochrane reviewers when classifying the risk of bias of original reports.

J Clin Epidemiol. 2020 Jul;123:1-8.

doi: 10.1016/j.jclinepi.2020.03.008..

ORIGINAL ARTICLE

External factors may influence Cochrane reviewers when classifying the risk of bias of original reports

Lorenzo Bertizzolo^{a,b,c,*}, Patrick M. Bossuyt^c, Ignacio Atal^{a,b,d,e}, Philippe Ravaud^{a,b,d,f,g,h},
Agnès Dechartresⁱ

^aINSERM, U1153 Epidemiology and Biostatistics Sorbonne Paris Cité Research Center (CRESS), Methods of Therapeutic Evaluation of Chronic Diseases Team (METHODS), Paris F-75004, France

^bParis Descartes University, Sorbonne Paris Cité, France

^cDepartment of Clinical Epidemiology and Biostatistics, Academic University Medical Centers, University of Amsterdam, Amsterdam, the Netherlands

^dCochrane France, Paris, France

^eCentre de Recherches Interdisciplinaires (CRI), Université Paris Descartes, University of Paris, Paris, France

^fCentre d'Epidémiologie Clinique, Hôpital Hôtel Dieu AP-HP (Assistance Publique - Hôpitaux de Paris), Paris, France

^gFaculté de Médecine, Université Paris Descartes, Sorbonne Paris Cité, Paris, France

^hDepartment of Epidemiology, Columbia University, Mailman School of Public Health, New York, USA

ⁱSorbonne Université, Institut National de la Santé et de la Recherche Médicale (INSERM), Institut Pierre Louis d'Epidémiologie et de Santé Publique, AP-HP, Sorbonne Université, Hôpital Pitié Salpêtrière, Département de Santé Publique, Centre de Pharmacopépidémiologie de l'AP-HP (Cephepi), CIC-1422, F75013, Paris, France

Accepted 12 March 2020; Published online 19 March 2020

Abstract

Objective: The objective of the study was to explore contextual factors associated with high or low risk-of-bias judgment in case of incomplete or unclear information in study reports.

Study Design and Setting: Research-on-research study, using matched case-control design, with a sample of 304 randomized controlled trials (RCTs) included in two Cochrane reviews for which there was disagreement on the risk-of-bias judgment related to incomplete or unclear information in the study report. A case was defined as an RCT judged at high or low risk of bias; a control was the same RCT judged at unclear risk. We used a conditional logistic regression model for analysis.

Results: Review authors being also authors of the RCT were more likely to assess an item at low risk of bias than unclear (OR: 11.71; 95% CI: 1.39e98.76). Earlier trials in a review were more often assigned a low risk (OR: 0.37; [0.15e0.96]). Review groups and authors that had completed a lower number of reviews slightly more often assigned a low risk, whereas others reported "unclear" (OR: 0.97, [95% CI: 0.95e0.99] for groups) and 0.97 (95% CI: 0.95e0.998) for authors).

Conclusions: Risk-of-bias assessment of RCTs in case of incomplete or unclear information may be affected by contextual factors. © 2020 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: Risk of bias; Systematic reviews; Research on research; Reporting; Matched case-control

Conflict of interest: None.

Transparency declaration: The guarantor (Lorenzo Bertizzolo) affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

Ethics approval: Not applicable. This is a research on research study.

Lorenzo Bertizzolo is the guarantor. He had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Funding: This project received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 676207. The funding source had no involvement in any part of the study.

Data sharing: Raw data and analyses are available on request from the authors.

* Corresponding author. INSERM, U1153 - Centre d'Epidémiologie Clinique, Hôpital Hôtel-Dieu, 1, place du parvis Notre Dame, 75004 Paris, France. Tel.: p 33 01 42 34 78 25; fax: p 33 (0) 142 160 332.

E-mail address: lorenzo.bertizzolo@gmail.com (L. Bertizzolo).

<https://doi.org/10.1016/j.jclinepi.2020.03.008>

0895-4356/© 2020 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

What is new?**Key findings**

- If review authors are authors of the RCT, they are more likely to consider the item at low risk of bias than unclear.
- Within a review, authors may differently judge a study that is more recent and larger than the other included studies.

What this adds to what was known?

- This is the first study exploring review, author, and study characteristics that may affect risk-of-bias assessment of a randomized controlled trial in case of incomplete or unclear information in the trial report.

What is the implication and what should change now?

- Authors of systematic reviews should be aware that their risk-of-bias judgment can be affected by factors that are not specific to the study itself.

mostly due to incomplete or unclear information in the published trial report [17].

Although an “unclear” judgment can be expected in case of incomplete or unclear information, many authors may feel confident enough to consider the information adequate to express a judgment of low or high risk of bias. Erroneous decisions apart, the RoB tool clearly leaves ample space for personal interpretation and judgment of evidence from study reports.

We wondered whether contextual factors, in particular identifiable characteristics of the review team or the review itself, might be associated with differences in risk-of-bias judgments when evaluating the same incomplete or unclear information. These characteristics could be related to the reviewers, the review, or the study itself, when compared with the other studies included in the review. For example, authors may consider studies that are larger or newer differently than smaller or older studies, or different review groups may have different approaches for handling incompleteness of information.

We aimed to explore which factors, if any, are associated with high or low risk-of-bias judgment in case of incomplete or unclear information in reports of primary studies for which an “unclear” risk-of-bias judgment would have been expected. We evaluated these by using a sample of Cochrane reviews.

1. Introduction

Evaluation of risk of bias in individual studies is an essential part of the systematic review process [1–3], which allows researchers to determine how strongly to make conclusions [1–3]. Cochrane has developed a tool, the Risk of Bias (RoB) tool, for assessing risk of bias in randomized controlled trials (RCTs) [4]. The tool has become the reference method for evaluating risk of bias [5]. This tool examines trial characteristics that are organized in seven items: random sequence generation, allocation concealment, blinding of participants and personnel, blinding of outcome assessment, incomplete outcome data, selective reporting, and other sources of bias. Each trial is evaluated at “high,” “low,” or “unclear” risk of bias on each of these seven items.

Previous studies showed a lack of reproducibility of assessments with the RoB tool [6–14]. Some recent studies also suggested that the support for judgment, and therefore the assessments themselves, frequently do not agree with recommendations from the Cochrane Handbook [15,16]. In a previous project [17], we evaluated the reasons for disagreements in risk-of-bias assessment of trials included in several Cochrane reviews. Our results confirmed suboptimal agreement in risk-of-bias assessment, ranging from 81% agreement for random sequence generation to 57% for incomplete outcome data. More than two-thirds of disagreements could be related to differences in interpretation of the same information and these disagreements were

2. Methods

This was a research-on-research study on risk-of-bias assessment that used a large collection of published Cochrane systematic reviews.

2.1. Study design

The study followed a matched case–control design with a sample of RCTs for which the risk of bias was assessed differently in two separate Cochrane reviews.

2.2. Data sources

Data were collected from the set of all Cochrane reviews published or updated between March 2011 and September 2014. The same set was used for a previous study [17]. The process of selection, evaluation of agreement, and in-depth analysis of disagreements was described in detail elsewhere [17].

Briefly, for each review, we obtained all data entered by review authors in RevMan, the software used for managing Cochrane reviews [11], as xml files [18]. We identified RCTs assessed for risk of bias in two separate systematic reviews. When more than two different reviews included the same study, we randomly selected one of the possible pairs. For each of the five key risk-of-bias items (random sequence generation, allocation concealment, blinding of participants and personnel, blinding of outcome

assessment, and incomplete outcome data), we compared the risk-of-bias judgments between the two reviews to evaluate whether they agreed (e.g., both reviews assessing allocation concealment concluded “low risk”) or not. For each disagreement, we analyzed the support for the corresponding judgment in the two reviews, to evaluate whether it was the same or not. In case of a difference in the support for the judgment, we examined the study report to evaluate whether the reason for the disagreement could be related to having access to different information (e.g., the study protocol, additional published reports, or contact with authors) or to differences in interpretation of the same information.

We classified each case of a difference in interpretation according to the likely reason: “confusion with another item of the tool,” “incomplete or unclear information in the study report,” or other. This assessment was conducted independently by two researchers; disagreements were resolved by consensus. More details can be found in the report of the previous study [17].

2.3. Selection of eligible studies

For the purpose of this study, we focused on trials for which we observed a disagreement between the two reviews in one or more of the five key risk-of-bias items (random sequence generation, allocation concealment, blinding of participants and personnel, blinding of outcome assessment, and incomplete outcome data) and attributed this disagreement to a difference in interpretation related to incomplete or unclear information in the study report. In such situations, a judgment “unclear” would be expected.

2.4. Definition of cases and controls

A case was defined as a judgment of “high risk of bias” or “low risk of bias” for an RCT in one of the reviews. A matched control was defined as a judgment of “unclear risk of bias” for the same RCT, for the same item, in the second, different review. Cases and controls were identified for each item.

2.5. Data extraction

We collected the following information regarding characteristics of reviews and studies included in the review:

Data regarding the review:

- Year of publication of the review.
- Cochrane review group: There are 53 Cochrane review groups within Cochrane in charge of conducting Cochrane reviews. Each group is dedicated to a specific topic (e.g., breast cancer). We identified the Cochrane review group that authored the review and the number of Cochrane reviews conducted by this group during the period of interest, from March 2011 to

September 2014. We used the number of reviews conducted by the review group as a measure of the volume/experience of the Cochrane review group.

- Authors: We collected the name of the first author and searched the number of Cochrane reviews conducted by this author up to the end of the period of interest (September 2014). The first reviews included in the Cochrane Database of Systematic Reviews date from 1995; therefore, we have identified all Cochrane reviews from 1995 to 2014. We also evaluated whether authors of the review were also authors of included RCTs (shared authorship RCT/review).
- Size of the review (number of RCTs included in the review).
- Type of interventions analyzed in the review (only pharmacological/only nonpharmacological/both).
- Mention of any type of conflict of interest by review authors.

Data regarding the RCT in comparison to the other RCTs included in the review:

- Sample size of the RCT in comparison with the other studies included in the review: From the xml file containing all the information reported in the review, we extracted the sample size of all studies included in the review. We calculated the distribution of the sample size for the RCTs included and then calculated the percentile of the sample size for the RCT of interest, expressed in a decimal number from 0 to 1.
- Year of publication of the RCT in comparison with the other studies included in the review: We followed the same process explained previously for the year of publication of the RCTs included in the review, calculating the rank percentile for the RCT of interest.
- Risk-of-bias judgment of the RCT in comparison with the other judgments for the same item in the review: We evaluated whether most of the judgments for the item within the review equals “unclear risk of bias.”

2.6. Statistical analysis

We evaluated characteristics potentially associated with judgment of high vs. unclear risk of bias and low vs. unclear risk of bias by using a conditional logistic regression model, accounting for the matched design. We evaluated each factor separately in a univariate analysis. The factors analyzed derive from the data we extracted and are listed in the previous paragraph. The results are expressed as odds ratios (ORs) with 95% confidence intervals (CIs) for each factor, considering separately disagreements of an assessment of high vs. unclear risk of bias and low vs. unclear risk of bias. Quantitative variables were analyzed as such, without dichotomization. We conducted a single joint analysis across different items and then subgroup analyses for each item separately.

Fig. 1. Flow-chart of the selection process.

The analyses were conducted with Stata 13.1 [19]. $P < 0.05$ was considered statistically significant. We did not adjust for multiple comparisons in this exploratory study.

3. Results

3.1. Selection process

Fig. 1 shows the details of the selection process. Our previous project [17] showed 625 disagreements, with the same trial judged at low or high risk of bias in one review and unclear in a second review. Among these disagreements, 402 were attributed to a different interpretation of the same information, which was considered related to incomplete or unclear information. These 402 disagreements related to 304 different RCTs, included in 376 different reviews.

3.2. Characteristics of included reviews

Table 1 shows the characteristics of the Cochrane systematic reviews included in our sample. The 376 reviews were compiled by 49 different Cochrane review groups, with the largest number (45; 12%) from the “pregnancy” group (Supplementary Table 1 gives a complete list of the different Cochrane review groups, with names shortened for simplicity in Tables 1 and 2). More than half of the reviews ($n = 204$, 54%) evaluated only nonpharmacological treatments. The number of RCTs included in reviews varied from 1 to 171 (median 16); only one-quarter of reviews had included more than 30 RCTs. Overall, 147

reviews (39%) reported a conflict of interest among authors of the review.

3.3. Characteristics of the disagreements

Across the 402 disagreements in risk-of-bias items, 69% ($n = 278$) were between low and unclear judgments; 31% ($n = 124$) were between high and unclear judgments.

The most common disagreements in both the disagreements between low and unclear and high and unclear judgments concerned incomplete outcome data ($n = 89$, 32% for low vs. unclear and $n = 41$, 33% for high vs. unclear disagreement) and allocation concealment ($n = 74$, 27% and $n = 29$, 23%, respectively) (details in Fig. 1).

Most disagreements derived from two judgments expressed by the same Cochrane group. This occurred in two-thirds of disagreements ($n = 261$, 65%). In 376 of the 804 (47%) single judgments analyzed, the most common judgment for that item in the review was “unclear” risk of bias. For only 15 judgments (2%), one of the authors of the systematic review was also among the authors of the RCT being evaluated.

3.4. Analysis of factors associated with disagreements

Table 2 compares review and study characteristics for the different judgments: low vs. unclear and high vs. unclear disagreements. Among the low vs. unclear disagreements, review groups and authors that had completed a lower number of reviews assigned significantly more often a low risk of bias where others reported “unclear,” (OR: 0.97 [95% CI: 0.95–0.99] and OR: 0.97 [95% CI:

Table 1. Characteristics of the included reviews, *N* = 376

Characteristics	n (%) Except otherwise indicated
Most common Cochrane review groups	
Pregnancy	45 (12)
Airways	22 (6)
Tobacco	21 (6)
Pain	20 (5)
Anesthesia	15 (4)
Respiratory infections	13 (3)
Mental disorders	13 (3)
Gynecology	13 (3)
Neonatal	13 (3)
Stroke	13 (3)
Drug addiction	12 (3)
Cystic fibrosis	12 (3)
Number of reviews compiled by the authoring review group during 2011–2014 ^a	22 (16–37)
Year of publication of the reviews ^a	2012 (2012–2013)
Intervention evaluated	
Pharmacological	145 (39%)
Nonpharmacological	204 (54%)
Both	27 (7%)
Mention of conflict of interest by reviewer	147 (39%)
Number of RCTs included in the reviews ^a	16 (8–31)
Year of publication of RCTs ^a included	2003 (1998–2007)

Abbreviation: RCTs, randomized controlled trials.

^a Median (1st and 3rd quartiles).

0.95–0.998] for one additional Cochrane review conducted by the review group or by the author, respectively).

Among the low risk-of-bias judgments, there were significantly a lower number of RCTs per review (OR: 0.99 [95% CI: 0.99–0.998]), but the OR close to 1 shows in fact a minor effect. Among the low risk-of-bias judgments, there were significantly more reviews in which one of the authors had also coauthored a trial report compared with controls with unclear assignments (OR: 11.71 [1.39–98.76]). There were also significantly fewer recent RCTs that were assigned a low risk of bias (OR: 0.37 [0.15–0.96]). A similar difference, although not significant, was observed for including larger trials (OR: 0.37 [0.13–1.02]).

Among the low risk-of-bias judgments, there were significantly fewer reviews that had assigned most RCTs in their review at an unclear risk of bias (OR: 0.12 [0.07–0.19]). Similarly, among the high risk-of-bias judgments in the high vs. unclear comparison, there were significantly fewer reviews that had evaluated most RCTs at an unclear risk of bias (OR: 0.11 [0.05–0.24]).

In the [Supplementary Table 2](#), we present reviews and study characteristics for cases and controls, for each of the five key RoB items separately. The results for the

separate items were affected by small sample sizes. The number of disagreements varied from 9 (random sequence generation) to 41 (incomplete outcome data) for high/unclear disagreements and 16 (blinding of participants and personnel) to 89 (incomplete outcome data) for low/unclear disagreements. We observed a significant difference for only the items random sequence generation, allocation concealment, and incomplete outcome data, with similar findings as in the overall analysis.

4. Discussion

4.1. Main findings

In this study, we evaluated factors or characteristics that are potentially associated with differences in risk-of-bias assessments of RCTs included in systematic reviews, in case of incomplete or unclear information given in reports. If review authors are also authors of the RCT, they are more likely to consider a low risk of bias than an unclear risk. If publication year of the RCT is more recent relative to the other RCTs included in the review, reviewers are more likely to judge an item “low” risk of bias than “unclear.” The number of Cochrane reviews conducted by the review group and by the first author was statistically

Table 2. Comparison of review characteristics between the different judgments (high vs. unclear and low vs. unclear risk of bias) for the included trials for all items of the Cochrane Risk of Bias tool

All items	Low <i>N</i> = 278	Unclear <i>N</i> = 278	OR (95% CI)
Year of publication of the review ^b	2012 (2012–2013)	2012 (2012–2013)	0.99 (0.8–1.22)
Most common review Cochrane groups	Tobacco: 28 (10%)	Pregnancy: 28 (10%)	-
	Airways: 24 (9%)	Pain: 26 (9%)	
	Pregnancy: 22 (8%)	Tobacco: 25 (9%)	
	Pain: 17 (6%)	Airways: 22 (8%)	
	Stroke: 13 (5%)	Stroke: 18 (6%)	
Number of reviews compiled by the review group ^b	22 (15–29)	23 (18–42)	0.97 (0.95–0.99)
Number of reviews compiled by the first author ^b	3 (1–5)	3 (1–7)	0.97 (0.95–0.998)
Shared author between the review and the RCT	9 (3%)	1 (0.4%)	11.71 (1.39–98.76)
Mention of conflict of interest	118 (42%)	129 (46%)	0.77 (0.50–1.18)
Number of RCTs included in the review ^b	23 (10–45)	29 (13–55)	0.99 (0.99–0.998)
Percentile distribution of the sample size of RCTs ^a	0.58 (0.3)	0.61 (0.29)	0.37 (0.13–1.02)
Percentile distribution of the year of RCTs ^a	0.56 (0.29)	0.59 (0.28)	0.37 (0.15–0.96)
Majority of judgments are at “unclear” risk	73 (26%)	192 (69%)	0.12 (0.07–0.19)
All items	High <i>N</i> = 124	Unclear <i>N</i> = 124	OR (95% CI)
Year of publication of the review ^b	2012 (2012–2013)	2012 (2011–2013)	1.15 (0.8–1.68)
Most common review Cochrane groups	Pregnancy: 19 (15%)	Pregnancy: 20 (16%)	-
	Pain: 10 (8%)	Tobacco: 10 (8%)	
	Tobacco: 9 (7%)	Drugs Addiction: 7 (6%)	
	Gynaeco. Cancer: 7 (6%)	Anesthesia: 6 (5%)	
	Developmental: 6 (5%)	Pain: 6 (5%)	
Number of reviews compiled by the review group ^b	22 (16–42)	22 (16–39)	1.00 (0.98–1.02)
Number of reviews compiled by the first author ^b	2 (1–5)	3 (1–6)	1.01 (0.98–1.03)
Shared author between the review and the RCT	3 (2%)	2 (2%)	1.60 (0.23–10.87)
Mention of conflict of interest	64 (52%)	54 (44%)	1.51 (0.86–2.68)
Number of RCTs included in the reviews ^b	21 (12–45)	26 (10–47)	1.00 (0.99–1.01)
Percentile distribution of the sample size of RCTs ^a	0.53 (0.3)	0.54 (0.3)	0.65 (0.14–3.01)
Percentile distribution of the year of RCTs ^a	0.55 (0.28)	0.58 (0.29)	0.24 (0.05–1.20)
Majority of judgments are at “unclear” risk	30 (24%)	81 (65%)	0.11 (0.05–0.24)

Abbreviation: RCTs, randomized controlled trials.

n (%) except otherwise indicated.

^a Mean (standard deviation).^b Median (1st and 3rd quartiles).

associated with different risk-of-bias judgments. The CIs for these associations are close to the no-significance level, but the variables are quantitative. Therefore, the

interpretation of the ORs is for one additional review conducted either by the review group or author so the effect may not be negligible.

4.2. Strengths and limitations

The main strength of our study is that it relies on real-life data. We did not directly explore the intention or attitude of researchers in handling incomplete information but evaluated associations between features and judgments by the observation of high-quality data from systematic reviews. Our sample size of studies and our careful work in collecting and categorizing data are also strengths.

Our study has some limitations. Our analysis of disagreements originating from incomplete information is not fully objective even if the classification was conducted in duplicate by two authors and reconciled, following a precise methodology, to reduce the subjectivity. We did not correct for multiple testing, because this analysis was exploratory, so the results should be interpreted carefully. Moreover, because of the low number of disagreements, particularly for high vs. unclear disagreements ($n = 124$), our analysis may lack power. We did not perform a multivariate analysis on our sample because of the exploratory nature of this study and because the different variables may interact between each other in an unstudied way. We are aware that our sample of reviews is not up to date, and we cannot exclude that a sample of more recent reviews would have given different results regarding the agreement in the risk-of-bias assessment and the relative importance of different reasons of disagreement. We nevertheless believe that the general point of our study—contextual factors may influence judgment regarding risk of bias—is not affected by the age of the sample. We evaluated factors/characteristics that are not part of the risk-of-bias assessment and authors are not aware of them. Therefore, there is no reason why these factors would disappear in a more recent sample.

4.3. Comparison with other studies

To our knowledge, this is the first study to explore the potential influence of characteristics pertaining to review groups, studies, and reviewer experience on risk-of-bias assessment. The research of Jørgensen et al. [6] and Savović et al. [20] was based on surveys and user comments about the use of the tool. The authors reported on the difficulties reviewers experience in judging incomplete or unclear information, but they did not analyze how these difficulties translate into real practice. More recently, Babic et al. [16] and Propadalo et al. [15] analyzed the support for judgments provided by reviewers. Their results also show how the same information in the study report can lead to a different risk-of-bias judgment, but they focused on an incorrect use of the RoB tool and did not analyze which factors may be associated with a different attitude toward the same information.

4.4. Interpretation of results and implications

We observed that judgments of unclear risk of bias more frequently came from reviews that judged most of the RCTs

included in the review at unclear risk. One possible explanation is that when the risk-of-bias assessment of a study is not completely straightforward (hence resulting in a disagreement between two different reviews), the reviewers may tend to go back to the judgment they used for most of the RCTs they assessed. This situation could represent a “halo” effect of the assignments for the other trials in the review. We may even suppose that the factors we identified, or possibly other factors, have a role in the risk-of-bias assessment regardless of the clarity or completeness of information.

An interesting point is that two-thirds of the disagreements originated within the same Cochrane review group. Nevertheless, we also showed that a higher number of reviews conducted by review groups and by authors, thereby suggesting more experience with risk-of-bias assessment, seems to have influenced reviewers in their judgment. This finding could suggest that the level of training may differ across author groups or it may be influenced by the experience of the reviewers. Of note, the influence of “experience” was toward more frequent judgments of unclear risk, instead of low risk.

The ORs for high vs. unclear disagreements did not reach statistical significance as compared with those for low vs. unclear disagreements. This finding might be explained in part by a lack of power of the study owing to the smaller sample size for the former comparison, but other interpretations are possible.

4.5. Conclusions

Our results indicate that identifiable review and author factors may play a role in risk-of-bias judgments of RCTs in systematic reviews. Specific factors that influenced reviewers in this analysis were the number of reviews they previously conducted and the number authored by the review group to which they belong. Reviewers also tended to judge a study at unclear risk of bias if they already did so for most of the studies included in the same review and tend to judge an item at “low” risk if they were involved in the RCT they are assessing. Awareness of contextual characteristics that can affect risk-of-bias judgments may help researchers in conducting more reproducible and meaningful assessments of risk of bias.

CRedit authorship contribution statement

Lorenzo Bertizzolo: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Resources, Software, Visualization, Writing - original draft, Writing - review & editing. **Patrick M. Bossuyt:** Conceptualization, Funding acquisition, Methodology, Resources, Supervision, Validation, Writing - original draft, Writing - review & editing. **Ignacio Atal:** Data curation, Software, Writing - review & editing. **Philippe**

Ravaud: Funding acquisition, Investigation, Methodology, Project administration, Resources, Writing - review & editing. **Agnès Dechartres:** Conceptualization, Funding acquisition, Investigation, Methodology, Project administration, Resources, Supervision, Validation, Writing - original draft, Writing - review & editing.

Acknowledgments

The authors thank David Tovey, former editor in chief of the Cochrane Library, for agreeing to share data from Cochrane reviews; Javier Mayoral Campos, system administrator; the Cochrane Central Executive for preparing files; and all Cochrane reviewers who collected data. The authors also thank Laura Smales for English revision of the manuscript.

Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jclinepi.2020.03.008>.

References

- [1] Cook DJ, Mulrow CD, Haynes RB. Systematic reviews: synthesis of best evidence for clinical decisions. *Ann Intern Med* 1997;126:376–80.
- [2] Hopewell S, Boutron I, Altman DG, Ravaud P. Incorporation of assessments of risk of bias of primary studies in systematic reviews of randomised trials: a cross-sectional study. *BMJ Open* 2013;3(8):e003342.
- [3] Hrobjartsson A, Boutron I, Turner L, Altman DG, Moher D. Cochrane Bias Methods G. Assessing risk of bias in randomised clinical trials included in Cochrane Reviews: the why is easy, the how is a challenge. *Cochrane Database Syst Rev* 2013;4:ED000058.
- [4] Higgins JP, Altman DG, Gotzsche PC, Juni P, Moher D, Oxman AD, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011;343:d5928.
- [5] Guyatt GH, Tugwell PX, Feeny DH, Haynes RB, Drummond M. A framework for clinical evaluation of diagnostic technologies. *Can Med Assoc J* 1986;134(6):587–94.
- [6] Jorgensen L, Paludan-Muller AS, Laursen DR, Savovic J, Boutron I, Sterne JA, et al. Evaluation of the Cochrane tool for assessing risk of bias in randomized clinical trials: overview of published comments and analysis of user practice in Cochrane and non-Cochrane reviews. *Syst Rev* 2016;5:80.
- [7] Wilkins AJ. Risk of bias in assessing risk of bias. *Ophthalmic Physiol Opt* 2017;37(1):107–9.
- [8] Hartling L, Hamm MP, Milne A, Vandermeer B, Santaguida PL, Ansari M, et al. Testing the risk of bias tool showed low reliability between individual reviewers and across consensus assessments of reviewer pairs. *J Clin Epidemiol* 2013;66:973–81.
- [9] Armijo-Olivo S, Ospina M, da Costa BR, Egger M, Saltaji H, Fuentes J, et al. Poor reliability between Cochrane reviewers and blinded external reviewers when applying the Cochrane risk of bias tool in physical therapy trials. *PLoS One* 2014;9:e96920.
- [10] Hartling L, Bond K, Vandermeer B, Seida J, Dryden DM, Rowe BH. Applying the risk of bias tool in a systematic review of combination long-acting beta-agonists and inhaled corticosteroids for persistent asthma. *PLoS One* 2011;6:e17242.
- [11] Hartling L, Ospina M, Liang Y, Dryden DM, Hooton N, Krebs Seida J, et al. Risk of bias versus quality assessment of randomised controlled trials: cross sectional study. *BMJ* 2009;339:b4012.
- [12] da Costa BR, Beckett B, Diaz A, Resta NM, Johnston BC, Egger M, et al. Effect of standardized training on the reliability of the Cochrane risk of bias assessment tool: a prospective study. *Syst Rev* 2017;6(1):44.
- [13] Armijo-Olivo S, Stiles CR, Hagen NA, Biondo PD, Cummings GG. Assessment of study quality for systematic reviews: a comparison of the Cochrane Collaboration risk of bias tool and the effective public health practice project quality assessment tool: methodological research. *J Eval Clin Pract* 2012;18:12–8.
- [14] Jordan VM, Lensen SF, Farquhar CM. There were large discrepancies in risk of bias tool judgments when a randomized controlled trial appeared in more than one systematic review. *J Clin Epidemiol* 2017; 81:72–6.
- [15] Propadalo I, Tranfic M, Vuka I, Barcot O, Pericic TP, Puljak L. In Cochrane reviews risk of bias assessments for allocation concealment were frequently not in line with Cochrane's Handbook guidance. *J Clin Epidemiol* 2018;106:10–7.
- [16] Babic A, Tokalic R, Amilcar Silva Cunha J, Novak I, Suto J, Vidak M, et al. Assessments of attrition bias in Cochrane systematic reviews are highly inconsistent and thus hindering trial comparability. *BMC Med Res Methodol* 2019;19:76.
- [17] Bertizzolo L, Bossuyt P, Atal I, Ravaud P, Dechartres A. Disagreements in risk of bias assessment for randomised controlled trials included in more than one Cochrane systematic reviews: a research on research study using cross-sectional design. *BMJ Open* 2019; 9(4):e028382.
- [18] Review Manager (RevMan) [Computer program]. Version 5. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration; 2014.
- [19] StataCorp. Stata Statistical Software: Release 13. College Station, TX: StataCorp LP; 2013.
- [20] Savovic J, Weeks L, Sterne JA, Turner L, Altman DG, Moher D, et al. Evaluation of the Cochrane Collaboration's tool for assessing the risk of bias in randomized trials: focus groups, online survey, proposed recommendations and their implementation. *Syst Rev* 2014;3:37.

Chapter 4: Association between methodological characteristics and treatment effect: a case example on testosterone therapy in men

4.1. Background

Despite eight decades of clinical use of testosterone, considerable controversy exists regarding the risks and benefits of testosterone therapy in men [150, 151]. Findings from previous methodological research suggests that inconsistencies in treatment effect estimates may be driven by methodological differences related to study design, sample size or participant characteristics [17, 18]. The aim of this study is to investigate the association between several trial characteristics and reported treatment effect estimates in randomized trials of testosterone therapy in men.

4.2. Methods

This is a meta-epidemiological study on the association between trial characteristics and reported treatment effect estimates.

Search Strategy: Published systematic reviews with meta-analysis of randomized trials on testosterone therapy in men were identified from MEDLINE.

Eligibility criteria and study selection: We included systematic reviews with meta-analyses, regardless of the specific dose/delivery of testosterone therapy or the comparator investigated in the included trials. We excluded systematic reviews with meta-analyses of individual participant data and meta-analyses computed using non-standard statistical methods (e.g. Bayesian). Sets of meta-analyses with overlapping trials were identified and all duplicate trials were removed. This harmonisation process was used to generate a dataset without overlap between meta-analyses [152].

Data collection: For each meta-analysis, the following data were extracted: year of publication, medical specialty, number of randomized trials included in the meta-analysis, outcome domain (e.g. cardiovascular disease incidence), interventions compared (experimental and control group), meta-analysis model (fixed-effect or random-effects), and treatment effect estimates and 95% confidence intervals (CI) for each included randomized trial. For each randomized trial included in the meta-analysis, information on trial characteristics (including year of publication, sample size, trial registration status, centre status, whether the corresponding study author was US-based or not, funding source and conflicts of interest of study authors) were extracted. We also assessed RoB of the included RCTs using the version 1 of the Cochrane RoB tool [153].

Statistical analyses: Treatment effects for binary outcomes were estimated as odds ratios (ORs) and treatment effects for continuous outcomes as standardised mean differences (SMDs). The direction

of effect was standardized so that an OR <1 or a SMD <0 indicated a beneficial effect of testosterone therapy. We analysed the association between each trial characteristic and the magnitude of a treatment effect using the two-step “meta-meta-analytic” approach for meta-epidemiological analyses described by Sterne et al. [118].

4.3. Results

Selection Process: Our PubMed search yielded 123 records. After screening all titles/abstracts, we retrieved 28 full-text articles for review, of which 24 were initially considered eligible for inclusion. After removing duplicate trials across the meta-analyses, we were left with a total of 19 non-overlapping meta-analyses [154-172] including 132 trials (comprising data from 10,725 participants). All meta-analyses compared testosterone replacement therapy with placebo.

Characteristics of the included studies: most of the trials were registered (79%), conducted in the year 2000 or later (77%), and comprised a sample size below N = 50 (48%). Across the 5 risk of bias items analysed, the percentage of trials rated at “unclear risk of bias” was: Random Sequence Generation (42%), Allocation Concealment (62%), Blinding of Participants and Personnel (41%), Blinding of Outcome Assessment (20%), Incomplete Outcome Data (17%); overall risk of bias was also calculated; the percentage of trials at unclear risk was 56%.

Association between trial characteristics and treatment effect: None of the investigated 13 trial characteristics were statistically significantly associated with treatment effect estimates in randomized trials of testosterone therapy in men (Figure 5). The direction of the RORs suggested that treatment effects were larger in more recently published randomized trials (i.e. published in the year 2000 or later), smaller trials (N < 50), trials that were registered, trials with a corresponding author based outside of the US, industry-funded trials, trials with an author with a conflict of interest, trials with low risk of bias due to allocation concealment and blinding of participants and personnel, trials with high/unclear risk of bias due to blinding of outcome assessors and missing data and trials with high/unclear overall risk of bias. However, the 95% CIs of all ROR estimates encompassed the null and associations in the opposite direction.

4.4. Discussion

We found no clear evidence that any trial characteristic was associated with treatment effects in the trials; all associations were statistically non-significant, with wide confidence intervals making it

impossible to rule out a positive, negative or null association. Trials with high/unclear risk of bias due to blinding of outcome assessors had larger effect estimates than trials rated at low risk of bias, a finding similar to that observed by others [17, 173]. In our analyses, a substantial number of studies were rated at “high/unclear risk of bias” (ranging from 17% to 62% across the domains). This finding confirms the persistent high prevalence of incomplete reporting (previous research suggests that 89% of published randomized trials include at least one “unclear” risk of bias domain [147]) and stresses the need for improved reporting in trials.

A potential explanation for the statistically non-significant findings in our study is the relatively small number of included meta-analyses and randomized trials. Alternatively, the small magnitude of the investigated characteristics might also reflect their minor discipline- and topic-specific impact in the field of testosterone research.

None of the trial characteristic examined was associated with treatment effects in the trials. The present meta-epidemiological study underlines the necessity for complete reporting to assess the safety and efficacy of testosterone therapy in men.

Figure 5: Main results of associations between trial characteristics and treatment effect estimates.

4.5. Article

Haring R, Ghannad M, Bertizzolo L, Page MJ.

No evidence found for an association between trial characteristics and treatment effects in randomized trials of testosterone therapy in men: a meta-epidemiological study.

J Clin Epidemiol. 2020 Jun;122:12-19.

doi: 10.1016/j.jclinepi.2020.02.004.

ORIGINAL ARTICLE

No evidence found for an association between trial characteristics and treatment effects in randomized trials of testosterone therapy in men: a meta-epidemiological study

Robin Haring^{a,b,*}, Mona Ghannad^{c,d}, Lorenzo Bertizzolo^{c,d}, Matthew J. Page^a^aSchool of Public Health and Preventive Medicine, Monash University, Melbourne, Australia^bEuropean University of Applied Sciences, Faculty of Applied Public Health, Rostock, Germany^cDepartment of Clinical Epidemiology, Amsterdam UMC, University of Amsterdam, Biostatistics and Bioinformatics, Amsterdam Public Health Research Institute, Meibergdreef 9, 1105, AZ Amsterdam, The Netherlands^dUniversité de Paris, CRESS, INSERM, INRA, F-75004, Paris, France

Accepted 19 February 2020; Published online 24 February 2020

Abstract

Objective: The objective of this study was to identify potential trial characteristics associated with reported treatment effect estimates in randomized trials of testosterone therapy in adult men.

Study Design and Setting: This is a meta-epidemiological study. MEDLINE was searched for meta-analyses of randomized trials of testosterone therapy in men published between 2008 and 2018. Data on trial characteristics were extracted independently by two reviewers. The impact of trial characteristics on reported treatment effects was investigated using a two-step meta-analytic approach.

Results: We identified 132 randomized trials, included in 19 meta-analyses, comprising data from 10,725 participants. None of the investigated design characteristics, including year of publication, sample size, trial registration status, center status, regionality, funding source, and conflict of interest were statistically significantly associated with reported treatment effects of testosterone therapy in men. Although trials rated at high risk of bias overall reported treatment effects that were 21% larger compared with trials rated at low risk of bias overall, the 95% confidence interval included the null (ratio of odds ratio: 0.79, 95% confidence interval: 0.60 to 1.03).

Conclusion: The present study found no clear evidence that trial characteristics are associated with treatment effects in randomized trials of testosterone therapy in men. To establish stronger evidence about the treatment effects of testosterone therapy in men, future randomized trials should not only be adequately designed but also transparently reported.

Study Registration: osf.io/x9g6m. © 2020 Elsevier Inc. All rights reserved.

Keywords: Testosterone; Men; Randomized trial; Meta-research; Risk of bias

1. Introduction

The clinical effects of the sex hormone testosterone inspired the research and development since the time it was isolated and synthesized in a Nobel Prize-winning effort in 1935 [1]. But in spite of 8 decades of clinical use, considerable controversy exists regarding the risks and benefits of testosterone therapy in men [2,3]. From measurement and diagnosis of low testosterone, over

treatment formulations and duration, to treatment monitoring and goals, the safety, efficacy, and effects of testosterone therapy in men are still under debate.

Consequently, guidelines for testosterone therapy in men demand high quality evidence to strengthen recommendations for clinical decision-making about potential treatment [4]. Findings from previous methodological research suggest that inconsistencies in the estimates of treatment effect may be driven by methodological differences related to study design, sample size, or participant characteristics [5,6]. For example, systematic reviews of meta-epidemiological studies suggest that larger treatment effects are observed in randomized trials with inadequate sequence generation and allocation concealment, and in trials with smaller sample size. In spite of several trial characteristics being consistently found to affect the

Conflict of interest statement: None of the authors have any nonfinancial interests that could be relevant to the submitted work.

* Corresponding author. School of Public Health and Preventive Medicine, Monash University, 553 St Kilda Rd, Melbourne, Victoria 3004, Australia. Tel.: 0049 381 8087-264; fax: 0049 381 8087 105.

E-mail address: rtharing@gmail.com (R. Haring).

<https://doi.org/10.1016/j.jclinepi.2020.02.004>

0895-4356/© 2020 Elsevier Inc. All rights reserved.

What is new?**Key findings**

- The present meta-epidemiological study found no clear evidence that trial characteristics are associated with treatment effects in randomized trials of testosterone therapy in men.

What this adds to what was known?

- This is the first meta-epidemiological study systematically investigating the influence of trial characteristics in randomized trials of testosterone therapy in men.

What is the implication and what should change now?

- To establish stronger evidence about the treatment effects of testosterone therapy in men, future randomized trials should not only be adequately designed but also transparently reported.

magnitude of treatment effects, their impact on the results of testosterone research is unknown. In addition, several factors, such as funding and conflicts of interest of study authors, have been investigated in comparatively fewer meta-epidemiological studies than other factors (e.g., sequence generation, blinding) [6].

Therefore, the aim of this study is to investigate the association between several trial characteristics (both commonly investigated and underexplored) and reported treatment effect estimates in randomized trials of testosterone therapy in men. Knowledge of these factors may help improve the design and conduct of future clinical trials to establish stronger evidence about the treatment effects of testosterone therapy in men.

2. Methods

We conducted this study in accordance with a study protocol we uploaded to the Open Science Framework in April 2018 (osf.io/x9g6m).

2.1. Search strategy

Published systematic reviews with meta-analysis of randomized on testosterone therapy in men were identified from MEDLINE, via PubMed, using the following search strategy:

("testosterone" [All Fields] OR "TRT" [All Fields] OR "androgens" [All Fields] OR "sex hormone" [All Fields]) AND Meta-Analysis [ptyp] AND "2008/04/20" [PDAT]: "2018/04/17" [PDAT] AND "male" [MeSH Terms] AND English [lang].

2.2. Eligibility criteria and study selection

We included systematic reviews with meta-analyses of binary outcomes or meta-analyses of continuous outcomes, regardless of the specific dose/delivery of testosterone therapy or the comparator (placebo/control/standard of care) investigated in the included trials. We excluded systematic reviews with meta-analyses of individual participant data and meta-analyses computed using nonstandard statistical methods (e.g., Bayesian). All records yielded from the search were exported to Covidence software for screening. Screening of titles/abstracts and full-text articles retrieved against the eligibility criteria was performed independently by two reviewers (RH and MJP). Once all eligible systematic reviews were identified, the list of randomized trials included in the largest meta-analysis was extracted from each review, and PubMed IDs (PMIDs) were assigned to each trial reference to identify duplicate trials. Sets of meta-analyses with overlapping trials were identified, and all duplicate trials were removed, starting with the meta-analysis with the smallest number of trials and moving sequentially up to the meta-analysis with the largest number of trials. When this process led to only one or no unique trials being identified in the meta-analysis, the meta-analysis was excluded. This harmonization process was used to generate a dataset without overlap between meta-analyses [7].

2.3. Data collection

Two reviewers (MG and LB) independently extracted the data from all the meta-analyses, and the randomized trials included in each meta-analysis, by using a standardized data collection form. Disagreements were resolved via discussion with RH and MJP. From each meta-analysis, the following data were extracted: year of publication, medical specialty, number of randomized trials included in the meta-analysis, outcome domain (e.g., cardiovascular disease incidence), interventions compared (experimental and control group), meta-analysis model (fixed-effect or random-effects), and treatment effect estimates and 95% confidence intervals (CI) for each included randomized trial.

From each randomized trial included in the meta-analysis, information on trial characteristics (including the year of publication, sample size, trial registration status, center status, whether the corresponding study author was United States-based or not, funding source and conflicts of interest of study authors) were extracted. All information was extracted from the trial report, and we did not contact the trial authors for clarification.

In addition, version 1 of the Cochrane risk of bias tool [8] was applied. We assessed the following domains of the tool: random sequence generation, allocation concealment, blinding of participants and personnel, blinding of outcome assessors, incomplete outcome data, and overall risk of bias. Each domain was judged as "low risk," "unclear risk," or "high risk of bias." The overall risk of bias

was judged as “low risk” if all domains were considered “low risk,” “unclear risk” if at least one domain was considered “unclear risk” but none were considered “high risk,” if at least one domain was judged to be “high risk.” We omitted the selective reporting bias domain given the recent rethinking of the domain as a source of bias to be assessed at the level of meta-analyses rather than at the level of included studies [9].

2.4. Statistical analyses

We categorized all trial characteristics into the following prespecified binary categories:

- Year of publication: published before 2000 vs. published 2000 or later;
- Sample size: less than 50 participants vs. 50 or more participants;
- Trial registration status: registered vs. not registered (as declared in the trial report);
- Center status: single-center or not specified vs. multi-center;
- Region: United States-based corresponding author vs. non-United States-based corresponding author;
- Funding: trial funded by industry vs. trial not funded by industry;
- Conflicts of interest: trial of with vs. without an author with a conflict of interest declared in the trial report;
- Risk of bias: all domains classified as high/unclear vs. low risk of bias.

Fig. 1. Flow diagram of identification, screening, and inclusion of meta-analysis.

Treatment effects for binary outcomes were estimated as odds ratios (ORs) and treatment effects for continuous outcomes as standardized mean differences (SMDs). The direction of effect was standardized so that an OR < 1 or an SMD < 0 indicated a beneficial effect of testosterone therapy. We analyzed the association between each trial characteristic and the magnitude of a treatment effect using the two-step “meta-meta-analytic” approach for meta-epidemiological analyses described by Sterne et al. [10]. That is, we first estimated within each meta-analysis a ratio of odds ratio (ROR) for binary outcomes or a difference in standardized mean differences (dSMD) for continuous outcomes, including standard errors between the two subgroup categories (e.g., industry vs. no industry funding), by using a random-effects meta-regression model. Then, we estimated a combined ROR across meta-analyses and the 95% CI by using a random-effects meta-analysis model. To synthesize estimates for binary and continuous outcomes, we converted dSMDs to log RORs by multiplying by $\pi/\sqrt{3} = 1.814$ [11]. DerSimonian and Laird’s method of moments estimator was used to estimate the between-meta-analysis variance. The inconsistency across RORs

was quantified using the I^2 statistic and the between-meta-analysis variance estimated by τ^2 . All analyses were stratified by “type of outcome” given previous studies [12,13] showing a significant difference between the ROR for objectively measured vs. patient-reported subjective outcomes. We planned to control for potential confounding by other trial characteristics by adjusting the meta-regression models for all other characteristics investigated, but decided against doing so given the number of included trials was not considered sufficient to provide reliable estimates. All analyses were performed using the *metan* and *metareg* commands in the statistical software package Stata (version 15).

3. Results

Our PubMed search yielded 123 records (Figure 1). After screening all titles/abstracts, we retrieved 28 full-text articles for review, of which 24 were initially considered eligible for inclusion. After removing duplicate trials across the meta-analyses, we were left with a total of 19

Table 1. Characteristics of included meta-analyses

Meta-analysis ID	Specialty	No. trials	Intervention	Comparator	Outcome	Outcome category	Outcome type
Alexander 2017 [14]	Cardiology	5	TRT	Placebo	Composite outcome of myocardial infarction, stroke, and mortality	Objective	Binary
Amanatkar 2014 [23]	Psychiatry	8	TRT	Placebo	Mood	Patient-reported	Continuous
Borst 2014 [21]	Cardiology	3	TRT	Placebo	Cardiovascular events	Objective	Binary
Corona 2011a [29]	Cardiology	6	TRT	Placebo	Time to 1 mm ST segment depression	Objective	Continuous
Corona 2011b [30]	Endocrinology	4	TRT	Placebo	HbA1c	Objective	Continuous
Corona 2014 [24]	Endocrinology	16	TRT	Placebo	Overall erectile function	Patient-reported	Continuous
Corona 2016 [17]	Cardiology	14	TRT	Placebo	Body weight	Objective	Continuous
Cui 2013 [26]	Urology	5	TRT	Placebo	Prostate-specific antigen levels	Objective	Continuous
Cui 2014 [24]	Urology	3	TRT	Placebo	PSA levels’ change in the short-term treatment studies	Objective	Continuous
Fernandez-Balsells 2010 [31]	Cardiology	7	TRT	Placebo	Composite prostate endpoint	Patient-reported	Binary
Grossmann 2015 [20]	Endocrinology	4	TRT	Placebo	HbA1c	Objective	Continuous
Kang 2015 [19]	Urology	4	TRT	Placebo	PSA levels	Objective	Continuous
Kohn 2016 [16]	Urology	11	TRT	Placebo	International Prostate Symptom Score	Patient-reported	Continuous
Neto 2015 [18]	Endocrinology	8	TRT	Placebo	Lean body mass	Objective	Continuous
Nian 2017 [15]	Urology	5	TRT	Placebo	Psychological subscale score	Patient-reported	Continuous
Price 2012 [27]	Cardiology	3	TRT	Placebo	Subjective improvement in symptoms of intermittent claudication	Patient-reported	Binary
Toma 2012 [28]	Cardiology	4	TRT	Placebo	Exercise capacity	Objective	Continuous
Xu 2013 [25]	Cardiology	15	TRT	Placebo	Cardiovascular-related events	Objective	Binary
Zarrouf 2009 [32]	Psychiatry	7	TRT	Placebo	HAM-D response	Patient-reported	Binary

TRT = testosterone replacement therapy.

nonoverlapping meta-analyses [14–32], including 132 trials (comprising data from 10,725 participants).

Of the 19 included meta-analyses, the median year of publication was 2014 (interquartile range (IQR) 2012–2015), and they included a median of 5 trials (IQR 4 to 8) (Table 1). The medical specialties investigated by the meta-analyses included cardiology (8 [42%]), urology (5 [26%]), endocrinology (4 [21%]), and psychiatry (2 [10%]). All meta-analyses compared testosterone replacement therapy with placebo. The majority of meta-analyses examined an objective outcome (12 [63%]), such as prostate-specific antigen (PSA) levels, HbA1c, body weight, and cardiovascular-related events. Patient-reported outcomes (investigated in 7 [37%] meta-analyses) included

self-reported erectile dysfunction, mood, and subjective improvement in cardiovascular or prostate-related symptoms. The type of outcome for most of the meta-analyses examined was continuous (13 [68%]).

Characteristics of the 132 trials are summarized in Table 2. Briefly, most of the trials were registered (79%), conducted in the year 2000 or later (77%), and comprised a sample size below $N = 50$ (48%). Across the six risk of bias domains, the percentage of trials rated at “unclear risk of bias” was: random sequence generation (42%), allocation concealment (62%), blinding of participants and personnel (41%), blinding of outcome assessors (20%), incomplete outcome data (17%), and overall risk of bias (56%).

Table 2. Trial characteristics extracted from included randomized-controlled trials of testosterone therapy in men

Trial characteristic		Number of studies (N = 132)
Time effect	study conducted before year 2000	30 (23%)
	study conducted year 2000 or later	102 (77%)
Sample size	> 200	12 (9%)
	101–200	21 (16%)
	50–100	35 (27%)
	< 50	64 (48%)
Trial registration	Registered	104 (79%)
	not registered	28 (21%)
Center status	single center randomized trial	55 (42%)
	multi-center randomized trial	31 (23%)
	not reported	46 (35%)
Regionality	U.S. author	59 (45%)
	non-U.S. author	73 (55%)
Funding source	industry funding	34 (26%)
	nonindustry funding	50 (38%)
	both	20 (15%)
	no funding	1 (< 1%)
	not reported	27 (20%)
COI statement	at least one trialist declares having a COI	32 (24%)
	no trialist declares having a COI	21 (16%)
	COIs not declared	79 (60%)
Risk of bias (RoB) characteristics	random sequence generation	3 = high risk (2%) 73 = low risk (56%) 56 = unclear risk (42%)
	allocation concealment	3 (2%) 47 (36%) 82 (62%)
	blinding of participants and personnel	9 (7%) 69 (52%) 54 (41%)
	blinding of outcome assessors	2 (1%) 104 (79%) 26 (20%)
	incomplete outcome data	25 (19%) 84 (64%) 23 (17%)
	overall risk of bias	35 (27%) 23 (17%) 74 (56%)

Fig. 2. Main results of associations between trial characteristics and treatment effect estimates.

None of the investigated 13 trial characteristics were statistically significantly associated with treatment effect estimates in randomized trials of testosterone therapy in men (Figure 2; more detailed results are provided Figures S1-S13 in the Appendix). In addition, there was no statistically significant interaction between the type of outcome (objective vs. patient-reported) and the magnitude of the ROR for any trial characteristic (see Figure S1-S13 in the Appendix). The direction of the RORs suggested that treatment effects were larger in more recently published randomized trials (i.e., published in the year 2000 or later), smaller trials ($N < 50$), trials that were registered, trials with a corresponding author based outside of the US, industry-funded trials, trials with an author with a conflict of interest, trials with low risk of bias due to allocation concealment and blinding of participants and personnel, trials with high/unclear risk of bias due to blinding of outcome assessors and missing data and trials with high/unclear overall risk of bias. However, the 95% CIs of all ROR estimates encompassed the null and associations in the opposite direction.

For most of the meta-meta-analyses, there was no or a small amount of between-meta-analysis heterogeneity (see Figures S1-S13 in the Appendix). However, for three meta-meta-analyses, the between meta-analysis heterogeneity was high (for the associations with country of corresponding author [I-squared 63%, tau-squared 0.49], risk of bias due to blinding of participants and personnel [I-

squared 61%, tau-squared 0.48], and risk of bias due to missing outcome data [I-squared 65%, tau-squared 0.66]).

4. Discussion

To our knowledge, this is the first meta-epidemiological study systematically investigating the influence of trial characteristics in randomized trials of testosterone therapy in men. We found no clear evidence that any trial characteristic was associated with treatment effects in the trials; all of the associations were statistically nonsignificant, with wide confidence intervals making it impossible to rule out a positive, negative, or null association. Further, for the most part, we observed little or no between-meta-analysis heterogeneity in the ROR estimates, except in three cases, where the heterogeneity appeared to be driven by the results of a single meta-analysis (by Neto et al.; we were unable to determine anything unique about this meta-analysis that would explain why it had very different results).

The direction of several of the ROR associations was comparable to the direction observed in previous meta-epidemiological studies. For example, we observed larger treatment effect estimates in randomized trials with less than 50 participants compared with trials with 50 or more participants, which is in line with previous studies reporting stronger effect estimates in small to moderately sized trials [33].

Also, trials with high/unclear risk of bias due to blinding of outcome assessors had larger effect estimates than trials rated at low risk of bias, a finding similar to that observed by others [5,34]. Furthermore, our findings suggest but do not confirm support for a potential *industry bias* (i.e., larger treatment effects when at least one investigator reported a conflict of interest), a finding comparable with other meta-epidemiological studies of industry ties with outcomes [35].

Our findings differ from that observed previously in several ways. The postulated *decline effect*, an association between year of study publication and reported effect size, was not statistically significant in our analysis. Despite the majority of testosterone therapy trials performed after the year 2000 ($N = 102$ vs. 30) and the tripled testosterone use in the United States from 2001 through 2011 [36], we did not detect a field-specific *decline effect* over time. Similarly, we did not detect the previously suggested *U.S. effect* of overestimated effect sizes from authors working in the United States [37]. Also previously reported associations between treatment effect magnitude and trial registration [38] or center status [39] were not confirmed in the present analyses.

In our analyses, a substantial number of studies were rated at “high/unclear risk of bias” (ranging from 17% to 62% across the domains). This finding confirms the persistently high prevalence of incomplete reporting (previous research suggests that 89% of published randomized trials include at least one “unclear” risk of bias domain [40]) and stresses the need for improved reporting in trials.

A potential explanation for the statistically nonsignificant findings in our meta-epidemiological study is the relatively small number of included meta-analyses and randomized trials. In spite of the fact that we systematically included all available, nonoverlapping meta-analyses of testosterone therapy in men in the present study, the potential risk of being underpowered cannot be ruled out. Thus, it is possible that a larger meta-epidemiological study, including more testosterone therapy trials, will be able to detect different or stronger bias associations between trial characteristics and treatment effect estimates. Alternatively, the small magnitude of the investigated characteristics might also reflect their minor discipline- and topic-specific impact in the field of testosterone research.

A key strength of our study includes the use of systematic methods to minimize the error in the identification and selection of meta-analyses and the collection of data from meta-analyses and trials. In addition, unlike many previous meta-epidemiological studies, we investigated the influence of a large number of ($n = 13$) of trial characteristics, several of which have been underexplored. However, our findings must be considered in light of some limitations. We did not contact the trialists for clarification about any missing or unclear information in the trial reports. Therefore, the associations between trial characteristics and treatment effects reflect what was *reported* in the trial report, not necessarily what was *done* by the trialists.

5. Conclusion

The present meta-epidemiological study underlines the necessity for complete reporting to assess the safety and efficacy of testosterone therapy in men. Additionally, authors of systematic reviews and meta-analysis of testosterone trials should carefully consider potential characteristics that may bias the results of the included studies. Given the unquestionable importance of well-designed and -conducted randomized trials for the production of high-quality evidence, future trials on testosterone therapy, should not only be adequately performed but also transparently reported.

CRedit authorship contribution statement

Robin Haring: Conceptualization, Methodology, Project administration, Writing - original draft. **Mona Ghanad:** Data curation. **Lorenzo Bertizzolo:** Data curation. **Matthew J. Page:** Conceptualization, Methodology, Formal analysis, Data curation, Software, Visualization, Writing - review & editing, Writing - original draft.

Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jclinepi.2020.02.004>.

References

- [1] Nieschlag E, Nieschlag S. The history of discovery, synthesis and development of testosterone for clinical use. *Eur J Endocrinol* 2019;180(6):R201–12.
- [2] Snyder PJ, Bhasin S, Cunningham GR, Matsumoto AM, Stephens-Shields AJ, Cauley JA, et al. Effects of testosterone treatment in older men. *N Engl J Med* 2016;374:611–24.
- [3] Walther A, Breidenstein J, Miller R. Association of testosterone treatment with alleviation of depressive symptoms in men: a systematic review and meta-analysis. *JAMA Psychiatry* 2019;76(1):31–40.
- [4] Bhasin S, Brito JP, Cunningham GR, Hayes FJ, Hodis HN, Matsumoto AM, et al. Testosterone therapy in men with hypogonadism: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab* 2018;103(5):1715–44.
- [5] Page MJ, Higgins JP, Clayton G, Sterne JA, Hrobjartsson A, Savovic J. Empirical evidence of study design biases in randomized trials: systematic review of meta-epidemiological studies. *PLoS One* 2016;11:e0159267.
- [6] Dechartres A, Trinquart L, Faber T, Ravaud P. Empirical evaluation of which trial characteristics are associated with treatment effect estimates. *J Clin Epidemiol* 2016;77:24–37.
- [7] Savovic J, Harris RJ, Wood L, Beynon R, Altman D, Als-Nielsen B, et al. Development of a combined database for meta-epidemiological research. *Res Synth Methods* 2010;1(3–4):212–25.
- [8] Higgins JP, Altman DG, Gotzsche PC, Juni P, Moher D, Oxman AD, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011;343:d5928.
- [9] Page MJ, Higgins JP. Rethinking the assessment of risk of bias due to selective reporting: a cross-sectional study. *Syst Rev* 2016;5(1):108.
- [10] Sterne JA, Juni P, Schulz KF, Altman DG, Bartlett C, Egger M. Statistical methods for assessing the influence of study characteristics on

- treatment effects in 'meta-epidemiological' research. *Stat Med* 2002; 21:1513–24.
- [11] Chinn S. A simple method for converting an odds ratio to effect size for use in meta-analysis. *Stat Med* 2000;19:3127–31.
 - [12] Ferreira-Gonzalez I, Busse JW, Heels-Ansdell D, Montori VM, Akl EA, Bryant DM, et al. Problems with use of composite end points in cardiovascular trials: systematic review of randomised controlled trials. *BMJ* 2007;334:786.
 - [13] Ciani O, Buyse M, Garside R, Pavey T, Stein K, Sterne JA, et al. Comparison of treatment effect sizes associated with surrogate and final patient relevant outcomes in randomised controlled trials: meta-epidemiological study. *BMJ* 2013;346:f457.
 - [14] Alexander GC, Iyer G, Lucas E, Lin D, Singh S. Cardiovascular risks of exogenous testosterone use among men: a systematic review and meta-analysis. *Am J Med* 2017;130(3):293–305.
 - [15] Nian Y, Ding M, Hu S, He H, Cheng S, Yi L, et al. Testosterone replacement therapy improves health-related quality of life for patients with late-onset hypogonadism: a meta-analysis of randomized controlled trials. *Andrologia* 2017;49(4).
 - [16] Kohn TP, Mata DA, Ramasamy R, Lipshultz LI. Effects of testosterone replacement therapy on lower urinary tract symptoms: a systematic review and meta-analysis. *Eur Urol* 2016;69(6):1083–90.
 - [17] Corona G, Giagulli VA, Maseroli E, Vignozzi L, Aversa A, Zitzmann M, et al. Testosterone supplementation and body composition: results from a meta-analysis study. *Eur J Endocrinol* 2016; 174(3):R99–116.
 - [18] Neto WK, Gama EF, Rocha LY, Ramos CC, Taets W, Scapini KB, et al. Effects of testosterone on lean mass gain in elderly men: systematic review with meta-analysis of controlled and randomized studies. *Age (Dordr)* 2015;37(1):9742.
 - [19] Kang DY, Li HJ. The effect of testosterone replacement therapy on prostate-specific antigen (PSA) levels in men being treated for hypogonadism: a systematic review and meta-analysis. *Medicine* 2015; 94(3):e410.
 - [20] Grossmann M, Hoermann R, Wittert G, Yeap BB. Effects of testosterone treatment on glucose metabolism and symptoms in men with type 2 diabetes and the metabolic syndrome: a systematic review and meta-analysis of randomized controlled clinical trials. *Clin Endocrinol (Oxf)* 2015;83(3):344–51.
 - [21] Borst SE, Shuster JJ, Zou B, Ye F, Jia H, Wokhlu A, et al. Cardiovascular risks and elevation of serum DHT vary by route of testosterone administration: a systematic review and meta-analysis. *BMC Med* 2014;12:211.
 - [22] Corona G, Isidori AM, Buvat J, Aversa A, Rastrelli G, Hackett G, et al. Testosterone supplementation and sexual function: a meta-analysis study. *J Sex Med* 2014;11(6):1577–92.
 - [23] Amanatkar HR, Chibnall JT, Seo BW, Manepalli JN, Grossberg GT. Impact of exogenous testosterone on mood: a systematic review and meta-analysis of randomized placebo-controlled trials. *Ann Clin Psychiatry* 2014;26(1):19–32.
 - [24] Cui Y, Zong H, Yan H, Zhang Y. The effect of testosterone replacement therapy on prostate cancer: a systematic review and meta-analysis. *Prostate Cancer Prostatic Dis* 2014;17:132–43.
 - [25] Xu L, Freeman G, Cowling BJ, Schooling CM. Testosterone therapy and cardiovascular events among men: a systematic review and meta-analysis of placebo-controlled randomized trials. *BMC Med* 2013;11:108.
 - [26] Cui Y, Zhang Y. The effect of androgen-replacement therapy on prostate growth: a systematic review and meta-analysis. *Eur Urol* 2013; 64(5):811–22.
 - [27] Price J, Leng GC. Steroid sex hormones for lower limb atherosclerosis. *Cochrane Database Syst Rev* 2012;10:CD000188.
 - [28] Toma M, McAlister FA, Coglianese EE, Vidi V, Vasaiwala S, Bakal JA, et al. Testosterone supplementation in heart failure: a meta-analysis. *Circ Heart Fail* 2012;5(3):315–21.
 - [29] Corona G, Rastrelli G, Monami M, Guay A, Buvat J, Sforza A, et al. Hypogonadism as a risk factor for cardiovascular mortality in men: a meta-analytic study. *Eur J Endocrinol* 2011;165(5):687–701.
 - [30] Corona G, Monami M, Rastrelli G, Aversa A, Sforza A, Lenzi A, et al. Type 2 diabetes mellitus and testosterone: a meta-analysis study. *Int J Androl* 2011;34(6 Pt 1):528–40.
 - [31] Fernandez-Balsells MM, Murad MH, Lane M, Lampropoulos JF, Albuquerque F, Mullan RJ, et al. Clinical review 1: adverse effects of testosterone therapy in adult men: a systematic review and meta-analysis. *J Clin Endocrinol Metab* 2010;95(6):2560–75.
 - [32] Zarrouf FA, Artz S, Griffith J, Sirbu C, Kommor M. Testosterone and depression: systematic review and meta-analysis. *J Psychiatr Pract* 2009;15(4):289–305.
 - [33] Dechartres A, Trinquart L, Boutron I, Ravaud P. Influence of trial sample size on treatment effect estimates: meta-epidemiological study. *BMJ* 2013;346:f2304.
 - [34] Hrobjartsson A, Thomsen AS, Emanuelsson F, Tendal B, Hilden J, Boutron I, et al. Observer bias in randomised clinical trials with binary outcomes: systematic review of trials with both blinded and non-blinded outcome assessors. *BMJ* 2012;344:e1119.
 - [35] Chartres N, Fabbri A, McDonald S, Turton J, Allman-Farinelli M, McKenzie J, et al. Association of industry ties with outcomes of studies examining the effect of wholegrain foods on cardiovascular disease and mortality: systematic review and meta-analysis. *BMJ Open* 2019;9(5):e022912.
 - [36] Baillargeon J, Kuo YF, Westra JR, Urban RJ, Goodwin JS. Testosterone prescribing in the United States, 2002–2016. *JAMA* 2018; 320:200–2.
 - [37] Baillargeon J, Urban RJ, Ottenbacher KJ, Pierson KS, Goodwin JS. Trends in androgen prescribing in the United States, 2001 to 2011. *JAMA Intern Med* 2013;173(15):1465–6.
 - [38] Odutayo A, Emdin CA, Hsiao AJ, Shakir M, Copsey B, Dutton S, et al. Association between trial registration and positive study findings: cross sectional study (Epidemiological Study of Randomized Trials-ESORT). *BMJ* 2017;356:j917.
 - [39] Unverzagt S, Prondzinsky R, Peinemann F. Single-center trials tend to provide larger treatment effects than multicenter trials: a systematic review. *J Clin Epidemiol* 2013;66:1271–80.
 - [40] Jorgensen L, Paludan-Muller AS, Laursen DR, Savovic J, Boutron I, Sterne JA, et al. Evaluation of the Cochrane tool for assessing risk of bias in randomized clinical trials: overview of published comments and analysis of user practice in Cochrane and non-Cochrane reviews. *Syst Rev* 2016;5:80.

Chapter 5: A survey on the use and knowledge of STROBE

5.1. Background

Authors of biomedical manuscripts are often unaware of the existence or utility of RG and those responding to peer reviewers often have problems adhering to the methodological standards proposed [2–4]. Many journals do not require a relevant RG checklist to be submitted with a manuscript, therefore, there is often no incentive for authors to complete one [5].

This study was designed to explore how researchers view and interact with one reporting guideline dedicated to observational research, the STrengthening the Reporting of Observational studies in Epidemiology (STROBE) Statement. We aimed to identify the personal and environmental facilitators, barriers, and motivators to using the STROBE Statement through an online survey.

5.2. Methods

Survey Design: We followed the CHERRIES guideline for online surveys [14]. After consenting to participate, adaptive questioning branched the survey based on participant's level of awareness and use of STROBE (*i.e.*, *never heard of*, *never used*; *heard of*, *never used*; *heard of*, *have used*). All participants were presented 25 questions informed by the Unified Theory of Acceptance and Use of Technology (UTAUT) scale [17,18].

Recruitment: Eligible participants were researchers involved in manuscript writing (within the past 10 years) reporting the results of observational studies. The survey was distributed from March 5 to August 31, 2018. Survey recruitment used several snowball and purposive sampling routes. Firstly, Authors used their professional network. Next, the survey was promoted through social media, primarily Twitter. We then emailed the editors of 257 biomedical journals identified in another study [9,20] and asked them to invite their authors to participate (e.g., via e-mail list-servs, Twitter, LinkedIn, etc.). When initial recruitment methods failed to provide sufficient respondents, we used Python to scrape emails of corresponding authors from an observational study corpus which examined endorsement of seven STROBE extensions [9,20,21]. To broaden the scope, we also included other journals primarily focused in Epidemiology.

Statistical Analyses: General information on demographics, STROBE extension awareness, research stage usage, and awareness referral mechanisms are presented as counts and percentages in the aggregate and per subgroup. Likert scale responses are reported as means and standard deviations. Completion/dropout rates were calculated overall and for the different groups (users of STROBE, never heard of STROBE, know it but never used).

5.3. Results

Of the 257 editorial offices contacted, 65 (25.3%) responded after three attempts. Of those who responded, 20 (30.8%) reported that they would invite their authors to participate (via Twitter, LinkedIn, listserv, blog, etc.), 42 (64.6%) declined to participate, and 3 (4.6%) reported individual-level participation.

After accounting for invalid addresses, 14,621 e-mails were sent to authors. Over 100 authors ($n = 109$) informed us of participation, 23 declined (giving no reason or stating no time/ interest), and 23 reported ineligibility (i.e., did not work in observational research). Another 145 were ineligible as they were unreachable during the recruitment period (e.g., family, sick, sabbatical leaves) or unreachable permanently (e.g., left job, retired, died).

As the survey was anonymous and recruitment methods used network snowballing, social media, and mass mailings, we cannot estimate the total number of people that read the survey invitation. However, we know that 1293 visitors read the informed consent page and 1265 (97.8%) agreed to participate. After evaluating free-text responses, seven indicated ineligibility (e.g., “*I do not do observational research*”). Of the 1258 eligible participants, 1015 (80.7%) completed the survey. Nearly 20% ($n = 195$) indicated they never heard of STROBE nor used it prior to the study (*group 1*), 18% ($n = 185$) had heard of it but never used it (*group 2*), and over half (63%, $n = 635$) had heard of and used it (*group 3*). The completion rates were 67% for group 1, 81% for group 2, and 97% for group 3 (Figure 3).

Motivators of Use: When asked about what factors would or have motivated use of STROBE, the journal submission process and mandatory reporting guideline use were the most frequently chosen options. After journal policies, self-motivation was among the top-ranked reported influences. Around half (53%) of those who were aware of STROBE but had not used it (*group 2*) reported that this was because journals did not require it. The next most frequently reported reason was that their co-authors did not use it (24.3%). (Table 2).

Usage Timing and Frequency: Participants who used STROBE (*group three*) most commonly did so during the manuscript writing process ($n = 451$) or after completing their draft to check that all relevant information had been reported ($n = 439$). Participants who had not used STROBE before (*groups one and two*; $n = 380$), most frequently reported that they would most likely use it during the manuscript writing process (62.6%; 68.1%) or during the protocol/study design stage (64.6%; 59.5%).

Awareness: Of those who were aware of STROBE prior to the survey (*groups 2 and 3*, $n = 820$), the most frequently reported route that made them aware of STROBE was a journal requiring or

recommending it [*group 2: n = 55, 29.7%; group 3: n = 234, 36.9%*]. The other options (peers, superiors, courses, or online) ranged from 12.6 to 16.7%. Most participants (70.7%, *n = 718*) indicated that they were not aware of any STROBE extension.

5.4. Discussion

Our project used a broad and multi-faceted sampling strategy which created a diverse sample of observational study authors. We also engaged nearly 200 participants who previously had never heard of STROBE, making our survey an awareness intervention itself.

Participants reported overall positive views towards STROBE, considering it useful, clear, and relatively easy to use. They also thought it would increase manuscript quality and the chances of getting published. However, they were not as positive regarding time requirements, reporting effects on productivity, speed and ease of writing. Our results should be reassuring to journal editors who fear losing authors to other journals with less strict requirements for publication [3].

These fears may be unfounded as participants indicated that, despite time costs, there are benefits of using an RG such as increasing the quality of their manuscripts and the chances of being published. Furthermore, they thought that the publishing environment (i.e., journals) would or do support its use. Widespread interventions are needed to improve RG adherence. Efforts to target research clusters, not just individuals, in order to foster broader support are needed. With increased uptake amongst co-authors completing reporting checklists, the time required may be reduced further, thus making using STROBE more appealing. When reporting guidelines become an expected part of the research process, self-regulation can occur and formal journal and institutional policies can be more fruitful as well [38]. Targeted and widespread promotion of reporting guidelines is needed to improve the completeness of reporting and reduce research waste [39].

5.5. Article

Sharp MK, Bertizzolo L, Rius R, Wager E, Gómez G, Hren D.

Using the STROBE statement: survey findings emphasized the role of journals in enforcing reporting guidelines.

J Clin Epidemiol. 2019 Dec;116:26-35.

doi: 10.1016/j.jclinepi.2019.07.019.

ORIGINAL ARTICLE

Using the STROBE statement: survey findings emphasized the role of journals in enforcing reporting guidelines

Melissa K. Sharp^{a,b,*}, Lorenzo Bertizzolo^b, Roser Rius^c, Elizabeth Wager^{d,e},
Guadalupe Gómez^c, Darko Hren^a

^aFaculty of Humanities and Social Sciences, Department of Psychology, University of Split, Split, Croatia

^bUniversité de Paris, CRESS, INSERM, INRA, F-75004 Paris, France

^cDepartament d'Estadística i Investigació Operativa, Universitat Politècnica de Catalunya-BarcelonaTech, Barcelona, Spain

^dSchool of Medicine, University of Split, Split, Croatia

^eSideview, Princes Risborough, UK

Accepted 30 July 2019; Published online 6 August 2019

Abstract

Objectives: The objective of the study was to identify factors affecting the use of the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement, specifically authors' attitudes toward and experiences with it.

Study Design and Setting: An online survey was distributed to authors of observational studies recruited via social media, personal network snowballing, and mass mailings using targeted search strategies. Data on demographics, awareness, motivators, and usage were collected in conjunction with a modified Unified Theory of Acceptance and Use of Technology (UTAUT) scale on which confirmatory factor analysis (CFA) was performed.

Results: One thousand fifteen participants completed the survey. Of these, 185 (18.2%) indicated they had never heard of STROBE nor used it previously, 195 (19.2%) had heard of it but never used it, and 635 (62.6%) had used it. Journals promoting STROBE were both key motivators and awareness mechanisms; peers and educational workshops were also important influencing factors to a lesser degree. The internal consistency of the modified UTAUT scale was strong (Cronbach's alpha 5 0.94). CFA supported a four-factor model with 23 questions.

Conclusion: The endorsement of STROBE by journals is key to authors' awareness and use of the guideline. We tested and validated our scale which can guide future research on reporting guidelines. © 2019 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: Observational studies; Guidelines as topic; Epidemiologic research design; Information dissemination/methods; STROBE; Online survey; Scientific writing

Authors' contributions: All authors have made substantive intellectual contributions to the development of the protocol and this manuscript. M.K.S. conceptualized the study and led the writing of the manuscript. D.H. led the supervision of the manuscript preparation. M.K.S. and L.B. managed survey recruitment. M.K.S., G.G., and R.R. assisted with all analyses. M.K.S. performed analyses. All authors read and approved the final manuscript.

Source of funding: This work was supported by the European Union's Horizon 2020 Research and Innovation Programme under the Marie Skłodowska-Curie grant agreement No 676207. Coauthor G.G. was supported by work from Ministerio de Economía y Competitividad, Grant/Award Numbers: MTM2015-64465-G2-1-R, MDM-2014-0445.

Ethics approval and consent to participate: Ethical approval was granted by the University of Split (2181-198-03-04-18-0010).

Availability of data and material: The final R Markdown code used for the current study will be made available on the Open Science Framework (<https://osf.io/2f9ny/>) and in the Zenodo repository in the Methods in Research on Research (MiRoR) community (<https://zenodo.org/communities/mirror/>).

Conflict of interest: M.K.S. works with the STROBE statement as a part of her doctoral studies. D.H., G.G., and E.W. provide support and mentoring as a part of the Methods in Research on Research (MiRoR) project. E.W. was a Fellow of the UK EQUATOR Centre which promotes the use of reporting guidelines (this was an unpaid position).

* Corresponding author: University of Split, Université de Paris, Poljicka cesta 35, Split 21000, Croatia. Tel.: p 1 248 701 8109.

E-mail addresses: msharp@unist.hr; melissaasharp@etu.parisdescartes.fr (M.K. Sharp).

<https://doi.org/10.1016/j.jclinepi.2019.07.019>

0895-4356/© 2019 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

What is new?**Key findings**

- We used a health technology assessment framework to investigate authors' attitudes and experiences of the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) reporting guideline. Our survey captured over 1,000 authors' views on the topic.
- Respondents reported STROBE to be useful, easy to use, clear, and understandable. However, they were concerned about the time needed to use the tool and expressed apprehension about how supportive peers and the research environment were toward using STROBE. Nearly 200 respondents were introduced to STROBE for the first time, making the survey an educational intervention itself.

What this adds to what was known?

- We have tested and validated a scale to assess author interactions with and views toward STROBE. This scale can be used to inform the promotion and evaluation of other reporting guidelines.

What is the implication and what should change now?

- Journals are key to raising awareness of reporting guidelines and enforcing their use. The research climate surrounding authors (i.e., peers and educational workshops) is also an important secondary influencing factor. Interventions should focus on establishing incentive systems and a culture change with these actors.

1. Introduction

Reporting guidelines (RGs) provide a protective “cognitive net” against the fallibility of human memory and support the skills of expert professionals [1]. Authors of biomedical manuscripts are generally unaware of the existence or utility of RGs and those responding to peer reviewers often have problems adhering to the methodological standards proposed [2–4]. Many journals do not require a relevant RG checklist to be submitted with a manuscript therefore, there is often no incentive for authors to complete one [5].

Some authors reject RGs, claiming that RGs can be condescending and rigid [6,7]. It is unclear what maintains these attitudes. Therefore, it would be useful to understand factors affecting use [8]. This study was designed to explore how researchers view and interact with one RG, the Strengthening the Reporting of Observational studies in Epidemiology (STROBE) Statement. STROBE was

created in 2007 to improve the reporting of observational studies (e.g., cross-sectional, cohort, case-control).

Many journals promote STROBE by requiring or recommending its use during the manuscript preparation process. However, endorsement rates are relatively low [9–13], and there is a diffusion of responsibility among journal editors, authors, and peer reviewers for RG compliance [3]. To better understand the current situation facing authors, we aimed to identify the personal and environmental facilitators, barriers, and motivators to using the STROBE statement. With this information, we hoped to extend the practical value of STROBE and perhaps other RGs.

2. Methods**2.1. Survey design**

We followed the CHERRIES guideline for online surveys (Supplemental File 1) [14]. Before distribution, we piloted the survey within the Methods in Research on Research (MiRoR) network [15], allowing collaborators to give feedback on content and functionality [16]. The University of Split School of Medicine Ethical Review Committee granted ethical approval.

The survey flow is presented in Figure 1, and the survey is in Supplemental File 2. All questions were forced response except for one optional open-ended question and mistakenly, the question asking about the respondent's country. After consenting to participate, adaptive questioning branched the survey based on participant's level of awareness and use of STROBE (i.e., never heard of, never used; heard of, never used; heard of, have used). After branching, participants were presented with questions about their interactions with STROBE (e.g., real or theoretical timing of use: writing a grant or peer-reviewing an article).

Next, all participants were presented 25 questions informed by the Unified Theory of Acceptance and Use of Technology (UTAUT) scale [17,18]. UTAUT is an amalgamation of eight dominant psychological and health technology assessment (HTA) theories and models that attempts to explain one's intention to use a piece of technology and their subsequent use behavior. The scale aims to explain information system usage behavior by measuring: Performance Expectancy (PE), Effort Expectancy (EE), attitude toward using technology, Social Influence (SI), Facilitating Conditions (FCs), self-efficacy, anxiety, and behavioral intention to use the tool [17]. HTA systematically evaluates direct and indirect consequences of using a piece of health technology. It can tap into whether the technology works, for whom, and at what cost [19].

We rephrased questions to be relevant to STROBE and kept the scale's four core constructs (PE, EE, SI, and FCs) (Fig. 2). Each subscale contained several items to ensure reliability and validity. The final version contained nine Likert scale items from PE, six from EE, five from SI, four from FCs, and one assessing the intention to use STROBE. Respondents rated statements on seven-point

Fig. 1. Survey flow. STROBE, Strengthening the Reporting of Observational Studies in Epidemiology.

Likert-type scales from “strongly disagree” to “strongly agree” (Supplemental File 2).

2.2. Recruitment

Eligible participants were researchers involved in manuscript writing (within the past 10 years) reporting the results of observational studies. The survey was distributed from March 5 to August 31, 2018.

Survey recruitment used several snowball and purposive sampling routes. First, M.K.S. invited her professional network and those involved in the MiRoR consortium [15] to participate. Next, the survey was promoted through social media, primarily Twitter. We then emailed the editors of 257 biomedical journals identified in another study [9,20] and asked them to invite their authors to participate (e.g., via e-mail list-servs, Twitter, LinkedIn, and so forth). Up to three e-mails were sent if they did not respond. When initial recruitment methods failed to provide sufficient respondents, we used Python to scrape emails of corresponding authors from an observational study corpus which examined endorsement of seven STROBE extensions [9,20,21]. To broaden the scope, we also included other journals primarily focused in Epidemiology. We identified 75 English language journals from the “Epidemiology” Broad Subject Term in the National Library of Medicine [22], 122 endorsing journals from the STROBE statement website [23], and 98 top-ranked journals in the Scimago Journal and Country 2017 “Medicine” ranking [24]. We

ran an Ovid MEDLINE observational study search filter from the same previous study [20] on all journals, deleted nonrelevant publication types (e.g., case summaries, editorials), and restricted the search to English language articles published within the past year (to reduce bounced emails). Supplemental File 3 details search strategies and journals searched. We deduplicated e-mails and sent up to two emails to each author.

2.3. Statistical analyses

General information on demographics, STROBE extension awareness, research stage usage, and awareness referral mechanisms is presented as counts and percentages

Fig. 2. Model and domain definitions. STROBE, Strengthening the Reporting of Observational Studies in Epidemiology.

Table 1. Sample demographics

Demographics	Total sample N (%) 1,015 (100)	Never heard of STROBE, never used (group 1) N (%) 195 (19)	Heard of STROBE, never used (group 2) N (%) 185 (18)	Heard of STROBE, have used (group 3) N (%) 635 (62)
Time spent in research				
1–10 years	332 (33)	57 (29)	65 (35)	210 (33)
11–30	362 (36)	107 (55)	95 (51)	372 (59)
31+	86 (10)	30 (15)	25 (14)	48 (8)
I do not work in research	3 (<1)	1 (0)	0 (0)	2 (<1)
Prefer not to say	3 (<1)	0 (0)	0 (0)	3 (<1)
Age				
18–34	185 (18)	36 (19)	38 (21)	111 (1)
35–54	589 (58)	101 (52)	83 (45)	405 (64)
55+	235 (23)	58 (30)	64 (35)	113 (18)
Prefer not to say	6 (<1)	0 (0)	0 (0)	6 (<1)
Gender				
Woman	469 (46)	97 (50)	82 (44)	289 (46)
Man	525 (52)	94 (48)	101 (55)	329 (52)
Trans	3 (<1)	0 (0)	0 (0)	3 (<1)
Prefer not to say	20 (2)	4 (2)	2 (1)	14 (2)
Region				
Africa	22 (2)	5 (3)	2 (1)	15 (2)
Asiatic region	31 (3)	7 (4)	4 (2)	20 (3)
Eastern Europe	33 (3)	12 (6)	5 (3)	16 (3)
Latin America	54 (5)	14 (7)	10 (5)	30 (5)
Middle East	26 (3)	11 (6)	6 (3)	9 (1)
Northern America	283 (28)	58 (30)	57 (31)	168 (27)
Pacific Region	54 (5)	4 (2)	10 (5)	40 (6)
Western Europe	465 (46)	69 (35)	83 (45)	313 (49)
Not reported	47 (5)	15 (8)	8 (4)	24 (4)

Abbreviation: STROBE, Strengthening the Reporting of Observational Studies in Epidemiology.

in the aggregate and per subgroup. Likert scale responses are reported as means and standard deviations. Completion/dropout rates were calculated overall and per group based on completion of the final forced-response question.

As we used a modified UTAUT scale (Table 3), we had a priori assumptions about our model and its latent factors (Fig. 1). Essentially, we were testing the HTA theory in our setting. Thus, confirmatory factor analysis (CFA) was used to test a four-factor model of intention to use STROBE in the overall sample and subgroups (Fig. 1). Rather than simply comparing average attitudes between groups, CFA allows us to test a theory and whether we captured relevant indicators and how they relate to each other (e.g., that we captured the key influencing factors that affect one's likelihood to use STROBE and furthermore, that we are comprehensive with our questioning and not redundant). All questions were scored from 1 to 7 and treated as continuous variables (Supplemental File 2). Three negatively worded questions from the EE scale were reverse-coded before calculating Cronbach's alpha and conducting CFA (see Table 4). For judging internal consistency, or the estimate of the reliability indicating the degree to which items measure different

aspects of the same concept, we used Cronbach's alpha and considered ≥ 0.7 an acceptable value [25,26].

All analyses were performed in R, version 3.4.0. The R Markdown file, containing code and output, is available on Open Science Framework [27]. The model was fit using lavaan, version 0.6–3 [28]. Maximum likelihood estimation (MLM specification) with robust standard errors was used to account for non-normality sample variance-covariance matrices and provide scaled test statistics. Latent factors were standardized, allowing for free estimation of all factor loadings. As suggested by Hu and Bentler [29], we considered Comparative Fit Index (CFI) and Tucker-Lewis Index (TLI) values ≥ 0.90 for acceptable and ≥ 0.95 for good fit, root mean squared error of approximation (RMSEA) values ≤ 0.06 (poor fit > 0.10), and standardized root mean squared residual (SRMR) values ≤ 0.08 to indicate a good fit between the model and data. When conducting multiple-group CFA, convergence issues are common [30]. When they occurred, we investigated the model within subgroups to detect issues with modification indices, individual factor loadings, and covariances between latent factors.

Table 2. Motivators, usage, and awareness descriptives

Question	Never heard of STROBE, never used; group 1, <i>n</i> = 195	Heard of STROBE, never used; group 2, <i>n</i> = 185	Heard of STROBE, used; group 3, <i>n</i> = 635
Motivator of use (past/theoretical)^{a,b}			
Self	128 (66)	55 (30)	308 (49)
Coauthors	57 (29)	72 (39)	116 (18)
Mentor/supervisor encouraged	40 (21)	-	-
Social norm	44 (23)	-	-
Journal submission process	104 (53)	134 (72)	376 (59)
Journal peer review	82 (42)	90 (49)	77 (12)
Incentivized in workplace	-	28 (15)	-
Immediate feedback	-	35 (19)	-
Free text	8 (4)	8 (4)	44 (7)
Reasons for not using^b			
My writing would not benefit	-	26 (14)	-
Do not want strict rules	-	25 (14)	-
Hard to understand	-	11 (6)	-
Word count	-	20 (11)	-
Format is difficult	-	16 (9)	-
Coauthors do not use it	-	45 (24)	-
May result in more negative review	-	1 (<1)	-
Journals do not require it	-	98 (53)	-
Not applicable for study type	-	25 (14)	-
Other	-	35 (19)	-
Frequency of current use			
Do not currently use	-	-	5 (<1)
Less than a quarter of manuscripts	-	-	305 (48)
Roughly half of manuscripts	-	-	134 (21)
Roughly 75% of manuscripts	-	-	118 (19)
All applicable manuscripts	-	-	73 (12)
Research stage of use (past/theoretical)^b			
Did not consider	22 (11)	10 (5)	NA
Protocol/design stage	126 (65)	110 (60)	239 (38)
Grant	76 (39)	61 (33)	89 (14)
Manuscript	122 (63)	126 (68)	451 (71)
After completing the article to check	98 (50)	80 (43)	439 (69)
Evaluating the article	70 (36)	69 (37)	243 (38)
Awareness mechanism			
Peer/colleague	-	31 (17)	97 (15)
Boss/mentor/supervisor	-	16 (9)	90 (14)
Journal	-	55 (30)	234 (37)
Course/workshop	-	32 (17)	105 (17)
Online	-	37 (20)	66 (10)
Other	-	14 (8)	43 (7)

^a Columns/items are blank as not all questions were presented to all branches.

^b Question allowed for multiple responses to be selected.

3. Results

Of the 257 editorial offices contacted, 65 (25.3%) responded after three attempts. Of those who responded, 20 (30.8%) reported that they would invite their authors to participate (via Twitter, LinkedIn, listserv, blog, etc.), 42 (64.6%) declined to participate, and 3 (4.6%) reported

individual-level participation. Reasons for declining included no access to a list, no time, a desire to remain neutral, the inability to contact authors due to General Data Protection Regulation restrictions [31], a belief that the journal did not publish observational studies (although we contend that it did), and a belief that the survey was flawed.

Table 3. Comparisons of model fit

Model	Group (n)	χ^2	df	RMSEA (90% CI) ^a	SRMR ^b	TLI ^{c,d}	CFI ^{c,d}	AIC
Four-factor model	Overall (1,015)	—	—	—	—	—	—	—
	1 (195)	776.900	266	0.109 (0.100, 0.118)	0.087	0.818	0.838	12593.900
	2 (185)	730.552	266	0.108 (0.099, 0.117)	0.085	0.797	0.802	12305.731
	3 (635)	1,582.699	266	0.102 (0.097, 0.107)	0.077^b	0.813	0.834	42,959.805
Final model: Four-factor model, No FC3, Method effects ^e	Overall (1,015)	1,931.539	717	0.078 (0.074, 0.082)	0.072^b	0.895	0.909^d	64,488.934
	1 (195)	489.527	239	0.079 (0.069, 0.089)	0.077^b	0.904^d	0.917^d	11,915.867
	2 (185)	496.303	239	0.084 (0.074, 0.095)	0.075^b	0.877	0.894	11,734.401
	3 (635)	927.172	239	0.076 (0.071, 0.081)	0.070^b	0.897	0.911^d	40,838.666

Abbreviations: χ^2 , chi-squared; df, degree of freedom; RMSEA, root mean square error of approximation with 90% confidence intervals; SRMR, square root mean residual; TLI, Tucker-Lewis Index; CFI, Comparative Fit Index; AIC, Akaike Information Criterion.

Bold = reached pre-established cutpoint threshold.

^a Within the range (≤ 0.06), indicating a good fit between the model and the data.

^b Within the range (≤ 0.08), indicating a good fit between the model and the data.

^c Within the range (≥ 0.95), indicating a good fit between the model and the data.

^d Within the range ($0.90 \leq x \leq .95$), indicating an acceptable fit between the model and the data.

^e Method effects address the reverse-coded items EE4, EE5, and EE6 and the high covariance between PE4 and PE5.

After accounting for 2,304 invalid addresses, 14,621 e-mails were sent to authors; we sent a second reminder e-mail to nonrespondents. Over 100 authors ($n = 109$) informed us of participation, 23 declined (giving no reason or stating no time/interest), and 23 reported ineligibility (i.e., did not work in observational research). Another 145 were ineligible as they were unreachable during the recruitment period (e.g., family, sick, sabbatical leaves) or unreachable permanently (e.g., left job, retired, died).

As the survey was anonymous and recruitment methods used network snowballing, social media, and mass mailings, we cannot estimate the total number of people that read the survey invitation. However, we know that 1,293 visitors read the informed consent page and 1,265 (97.8%) agreed to participate. After evaluating free-text responses, seven indicated ineligibility (e.g., “I do not do observational research”). Of the 1,258 eligible participants, 1,015 (80.7%) completed the survey. Nearly 20% ($n = 195$) indicated they never heard of STROBE nor used it before the study (group 1), 18% ($n = 185$) had heard of it but never used it (group 2), and over half (63%, $n = 635$) had heard of and used it (group 3) (Fig. 3, Table 1). The completion rates were 67% for group 1, 81% for group 2, and 97% for group 3 (Fig. 3).

We found fairly equal distributions for demographic categories across groups (Table 1, Supplemental File 4). The top five countries responding were the United States (21.6%), United Kingdom (9.8%), Italy (6.8%), Canada (6.4%), and Australia (4.9%). To account for the multidisciplinary nature of research, we allowed up to three selections for area of work. Participants working in public health and epidemiology were well represented with 470 (46%) and 247 (24%), respectively, choosing those options as one of their primary fields of work.

3.1. Motivators of use

When asked about what factors would or have motivated use of STROBE, the journal submission process and mandatory RG use were the most frequently chosen options. After journal policies, self-motivation was among the top-ranked reported influences. Around half (53%) of those who were aware of STROBE but had not used it (group 2) reported that this was because journals did not require it. The next most frequently reported reason was that their coauthors did not use it (24.3%) (Table 2).

3.2. Usage timing and frequency

Participants who used STROBE (group three) most commonly did so during the manuscript writing process ($n = 451$) or after completing their draft to check that all relevant information had been reported ($n = 439$). Participants who had not used STROBE before (groups one and two; $n = 380$) most frequently reported that they would most likely use it during the manuscript writing process (62.6%; 68.1%) or during the protocol/study design stage (64.6%; 59.5%). For those who previously used STROBE (group 3), 48% used it for less than a quarter of their manuscripts, whereas 11.5% used it for all of their manuscripts (Table 2).

3.3. Awareness

Of those who were aware of STROBE before the survey (groups 2 and 3, $n = 820$), the most frequently reported route that made them aware of STROBE was a journal requiring or recommending it (group 2: $n = 55$, 29.7%; group 3: $n = 234$, 36.9%). The other options (peers, superiors, courses, or online) ranged from 12.6 to 16.7%. A

Table 4. Factor loadings of final model ($n = 1,015$)

Items	Performance Expectancy (PE)			Effort Expectancy (EE)			Social Influence (SI)			Facilitating Conditions (FC)		
	1 ^a	2 ^b	3 ^c	1 ^a	2 ^b	3 ^c	1 ^a	2 ^b	3 ^c	1 ^a	2 ^b	3 ^c
PE1 STROBE will be/is useful in my job	0.728	0.710	0.665									
PE2 Using STROBE will enable/enables me to write papers more quickly	0.868	0.821	0.818									
PE3 STROBE will increase/increases my productivity	0.865	0.817	0.775									
PE4 If I use STROBE, I (will) increase my chances of getting published	0.766	0.699	0.511									
PE5 If I use STROBE, I will get a more positive peer review of my paper	0.737	0.670	0.553									
PE6 Using STROBE will make/makes it easier for me to write papers	0.903	0.861	0.864									
PE7 Using STROBE will improve/improves the quality of my manuscripts	0.803	0.675	0.767									
PE8 Using STROBE will make/makes my manuscript writing more efficient	0.849	0.867	0.870									
PE9 Using STROBE increases the quality of my output for the same amount of effort	0.850	0.831	0.804									
EE1 I think STROBE will be/is easy to use				0.841	0.771	0.887						
EE2 I think STROBE's content is clear and understandable				0.869	0.833	0.866						
EE3 I think that it will be/is easy for me to become skillful at using STROBE				0.793	0.797	0.693						
EE4 Using STROBE will take/takes too much time compared with my normal writing process*				0.437	0.464	0.604						
EE5 STROBE is so complicated, it will be/is difficult to understand what to do*				0.622	0.579	0.671						
EE6 Will take/takes too long to learn how to properly use STROBE to make it worth the effort*				0.569	0.542	0.598						
SI1 My peers will think/think that I should use STROBE							0.848	0.909	0.870			
SI2 My superiors will think/think that I should use it							0.621	0.562	0.639			
SI3 The research climate is helpful in promoting the use of reporting guidelines such as STROBE							0.887	0.890	0.831			
SI4 In general, I think that journals will support/support the use of STROBE							0.649	0.461	0.473			
SI5 I will use STROBE because a lot of scientists in my field are using it							0.531	0.553	0.549			
FC1 I have the knowledge necessary to use STROBE										0.599	0.490	0.567
FC2 STROBE is compatible with my current workflow										0.785	0.777	0.817
FC4 Using STROBE fits well with the way I like to work										0.852	0.878	0.843

Abbreviation: STROBE, Strengthening the Reporting of Observational Studies in Epidemiology.

1^a Subgroup one: Never heard of STROBE and never used it ($n = 195$).

2^b Subgroup two: Heard of STROBE but never used it ($n = 185$).

3^c Subgroup three: Heard of STROBE and have used it ($n = 635$).

Fig. 3. Participant flow diagram.

majority of participants (70.7%, $n = 718$) indicated that they were not aware of any STROBE extension.

3.4. Confirmatory factor analysis on modified UTAUT scale

We attempted CFA on the overall sample, but it would not converge. Therefore, we investigated the model within subgroups to identify convergence issues; it converged in all subgroups (Table 3). There were three recurring issues across groups: (1) the third FC item (FC3) appeared to not belong to the FC scale; (2) the FC and EE covariance was very high (0.88–0.91); and (3) two pairs of items (EE4:EE5 and PE4:PE5) had significant shared variance, with the highest modification indices across all subgroups.

Model fit statistics and accompanying step-by-step descriptions are in Supplemental File 4, Table 6. Succinctly, the FC3 item phrasing was redundant with EE1. When FC3 was deleted, the model converged. This also reduced the high covariance between the FC and EE factors. The EE4 and EE5 items, along with EE6, were negatively worded, so we allowed them to covary to account for method effects [32]. Items PE4 and PE5 were also allowed to covary as they were both related to academic publishing,

suggesting that they could covary for reasons other than the shared influence of the latent factor.

The four-factor model addressing these issues was the best fit model for our data (Table 3). The CFI (0.91) and TLI (0.90) reached the “acceptable” cut point of 0.90. The SRMR (0.07) was below its cut point of 0.08. The RMSEA (0.08) was not less than 0.06, however. All factor loadings were statistically significant (all p s ≤ 0.001) and salient (0.437 to 0.909) (Table 4). The internal consistency reliability of all four subscales was strong (Cronbach’s $\alpha \geq 0.94$ for all). Our items were parsimonious, functional, and internally consistent.

An overall pattern between groups was seen where those who had used STROBE before (group 3) had the highest scores, those who had never heard of STROBE before the survey (group 1) had second highest scores, and those who had heard of STROBE but never used it (group 2) most often had the lowest scores (Additional File 4, Table 4; Fig. 3).

4. Discussion

To the best of our knowledge, this is the first project to ask authors about their attitudes toward and experiences with STROBE, especially using HTA framework. Our project used a broad and multifaceted sampling strategy which created a diverse sample of observational study authors. We also engaged nearly 200 participants who previously had never heard of STROBE, making our survey an awareness intervention itself.

The large sample enabled us to test a modified UTAUT scale on our entire sample and within subgroups. With an acceptable fit between our model and the data, we expect that this instrument may be useful for evaluating interactions with other RGs. Our results confirm the applicability of an HTA approach to RGs, reveal important factors impacting STROBE use, and highlight a unique additional aspect of use, which may separate it from other pieces of technology—the academic publishing environment. Because our model-data fit was only “acceptable” and we needed to address shared error variance of two publishing-related items (PE4 and PE5), we believe that these two PE items might signal an unaccounted latent factor related to publishing.

Our CFA should be considered complementary to the descriptive results which emphasize the key role that journals have in raising awareness, motivating, and enforcing use. Journals were the most typical medium by which participants originally became aware of STROBE. Moreover, journals not requiring STROBE were the top reason why authors did not use it. We recommend that future work explores this concept more deeply. We suggest building on the most parsimonious model (Table 3), not including FC3, accounting for method effects on the EE scale, and addressing the shared error variance of PE4 and PE5.

With regard to limitations, estimating a sample size was not tenable as there is no clearly defined participant pool. In

addition, we used mass mailings, thus, potentially, some e-mails were likely blocked by spam filters [33]. In addition, we had differential dropout rates between groups which is expected as is conceptually harder to think in theoretical terms (e.g., when would you consider using STROBE which you were just introduced to vs. when have you used it). The differences in participation rates between groups and nonresponse and self-selection biases also could have skewed our responses to be more positive toward STROBE. In addition, the introduction to STROBE may have not been detailed enough and/or the participants may have not spent enough time on it.

Despite these limitations, overall, participants reported positive views toward STROBE, considering it useful, clear, and relatively easy to use. They also thought it would increase manuscript quality and the chances of getting published. However, they were not as positive regarding time requirements, reporting effects on productivity and speed and ease of writing. Our results should be reassuring to journal editors who fear losing authors to other journals with less-strict requirements for publication [3]. These fears may be unfounded as participants indicated that, despite time costs, there are benefits of using an RG such as increasing the quality of their manuscripts and the chances of being published. Furthermore, they thought that the publishing environment (i.e., journals) would or do support its use.

Despite this perceived benefit of an increase in quality, we caution that, empirically speaking, the research in this regard is mixed. Recent work (2019) demonstrated that having a methodological reviewer dedicated to looking for missing RG items (not only STROBE) increased the number of citations that an article received by 43% [34]. This could be perceived as a proxy for higher quality or impact. Conversely, other authors have found no effect on the reporting of confounding [35] or insufficient evidence to determine an impact on overall completeness of reporting [4,36]. To further assuage editors' (and authors') concerns, more research is needed in this area which focuses on a broad range of journals (i.e., not only high impact) and which takes endorsement type (i.e., requiring vs. recommending use) into account.

A 2019 scoping review complements our results, highlighting the complexity of RG adherence and highlighting the need to implement interventions with different stakeholders throughout the research process [37]. Their review showed that most of the evaluated interventions to improve RG adherence have been conducted in journals. There have been mixed results but promising ones for more active implementation efforts (i.e., requiring a checklist with submission), including editorial assistants trained on reporting issues, and automatic peer review tools.

Widespread interventions are needed to improve RG adherence. Efforts to target research clusters, not just individuals, to foster broader support are needed. With increased uptake among coauthors completing reporting

checklists, the time required may be reduced further, thus making using STROBE more appealing. When RGs become an expected part of the research process, self-regulation can occur and formal journal and institutional policies can be more fruitful as well [38]. Targeted and widespread promotion of RGs is needed to improve the completeness of reporting and reduce research waste [39].

CRediT authorship contribution statement

Melissa K. Sharp: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Visualization, Writing - original draft, Writing - review & editing. **Lorenzo Bertizzolo:** Data curation, Methodology, Writing - review & editing. **Roser Rius:** Formal analysis, Methodology, Writing - review & editing, Supervision. **Elizabeth Wager:** Writing - review & editing. **Guadalupe Gómez:** Formal analysis, Methodology, Writing - review & editing, Supervision, Resources. **Darko Hren:** Supervision, Methodology, Formal analysis, Visualization, Writing - review & editing, Resources.

Acknowledgments

The authors would like to acknowledge the Methods in Research on Research (MiRoR) network for their support and guidance. The authors would also like to thank Dr. Michail Kovanis for assistance gathering author emails and Ms. Alice Biggane for her assistance with the recruitment strategy.

Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.jclinepi.2019.07.019>.

References

- [1] Gawande A. The checklist manifesto: How to get things right. New York, NY: Metropolitan Books; 2010.
- [2] Costa BR da, Cevallos M, Altman DG, Rutjes AWS, Egger M. Uses and misuses of the STROBE statement: bibliographic study. *BMJ Open* 2011;1:e000048.
- [3] Grindlay DJ, Dean RS, Christopher MM, Brennan ML. A survey of the awareness, knowledge, policies and views of veterinary journal Editors-in-Chief on reporting guidelines for publication of research. *BMC Vet Res* 2014;10:10.
- [4] Cobo E, Cortés J, Ribera JM, Cardellach F, Selva-O'Callaghan A, Kostov B, et al. Effect of using reporting guidelines during peer review on quality of final manuscripts submitted to a biomedical journal: masked randomised trial. *BMJ* 2011;343:d6783.
- [5] Larson EL, Cortazar M. Publication guidelines need widespread adoption. *J Clin Epidemiol* 2012;65:239–46.
- [6] MacMahon B, Weiss NS. Is there a dark phase of this STROBE? *Epidemiology* 2007;18:791.
- [7] Rothman KJ, Poole C. Some guidelines on guidelines: they should come with expiration dates. *Epidemiology* 2007;18:794.

- [8] Hopewell S, Boutron I, Altman DG, Barbour G, Moher D, Montori V, et al. Impact of a web-based tool (WebCONSORT) to improve the reporting of randomised trials: results of a randomised controlled trial. *BMC Med* 2016;14:199.
- [9] Sharp MK, Tokalić R, Gómez G, Wager E, Altman DG, Hren D. A cross-sectional bibliometric study showed suboptimal journal endorsement rates of STROBE and its extensions. *J Clin Epidemiol* 2019;107:42–50.
- [10] Hua F, Walsh T, Glenny A-M, Worthington H. Surveys on reporting guideline usage in dental journals. *J Dent Res* 2016;95:1207–13.
- [11] Kunath F, Grobe HR, Rücker G, Engehausen D, Antes G, Wullich B, et al. Do journals publishing in the field of urology endorse reporting guidelines? A survey of author instructions. *Urol Int* 2011;88:54–9.
- [12] Meerpohl JJ, Wolff RF, Niemeyer CM, Antes G, von Elm E. Editorial policies of pediatric journals: survey of instructions for authors. *Arch Pediatr Adolesc Med* 2010;164:268–72.
- [13] Toews I, Binder N, Wolff RF, Toprak G, von Elm E, Meerpohl JJ. Guidance in author instructions of hematology and oncology journals: a cross sectional and longitudinal study. *PLoS One* 2017;12:e0176489.
- [14] Eysenbach G. Improving the quality of web surveys: the checklist for reporting results of internet E-surveys (CHERRIES). *J Med Internet Res* 2004;6:e34.
- [15] Projet MiRoR | An innovative and ambitious joint doctoral training programme. Available at <http://miro-ejd.eu/>. Accessed July 16, 2018.
- [16] Welcome to SurveyMonkey!. Available at <https://www.surveymonkey.com/home/>. Accessed March 17, 2017.
- [17] Venkatesh V, Morris MG, Davis GB, Davis FD. User acceptance of information technology: toward a unified view. *MIS Q* 2003;27:425–78.
- [18] Williams MD, Rana NP, Dwivedi YK. The unified theory of acceptance and use of technology (UTAUT): a literature review. *J Enterp Inf Manag* 2015;28:443–88.
- [19] Technology (U.S.) NIC on HSR& HC. HTA 101: Introduction to Health Technology Assessment. Available at <https://www.nlm.nih.gov/nichsr/hta101/ta10103.html>. Accessed July 2, 2019.
- [20] Sharp MK, Utrobičić A, Gómez G, Cobo E, Wager E, Hren D. The STROBE extensions: protocol for a qualitative assessment of content and a survey of endorsement. *BMJ Open* 2017;7:e019043.
- [21] Sharp MK. Methods in research on research (MiRoR). STROBE guidelines (ESR9). 2017. Available at osf.io/u75gb. Accessed July 1, 2019.
- [22] Broad subject terms for indexed journals. Available at <https://wwwcf.nlm.nih.gov/serials/journals/index.cfm>. Accessed July 16, 2018.
- [23] STROBE statement: endorsement. Available at <https://www.strobe-statement.org/index.php?id=strobe-endorsement>. Accessed July 16, 2018.
- [24] Journal rankings on agricultural and biological sciences. Available at <https://www.scimagojr.com/journalrank.php?area=2700>. Accessed July 16, 2018.
- [25] Cronbach LJ. Coefficient alpha and the internal structure of tests. *Psychometrika* 1951;16:297–334.
- [26] Bland JM, Altman DG. Statistics notes: Cronbach's alpha. *BMJ* 1997;314:572.
- [27] Sharp MK. Use of reporting guidelines as an educational intervention for teaching research methods and writing. *Open Sci Framew* <https://doi.org/10.17605/OSF.IO/HP623>. Accessed November 16, 2017.
- [28] Rosseel Y. Lavaan: an R package for structural equation modeling. *J Stat Softw* 2012;1(2):2012.
- [29] Hu L, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis: conventional criteria versus new alternatives. *Struct Equ Modeling Multidiscip J* 1999;6:1–55.
- [30] Brown TA. Confirmatory factor analysis for applied research. 2nd ed. New York, NY: The Guilford Press; 2015.
- [31] EU GDPR Information Portal. EU GDPR portal. Available at <http://eugdpr.org/eugdpr.org-1.html>. Accessed August 10, 2018.
- [32] Maul A. Method effects and the meaning of measurement. *Front Psychol* 2013;4:169.
- [33] Fan W, Yan Z. Factors affecting response rates of the web survey: a systematic review. *Comput Human Behav* 2010;26:132–9.
- [34] Vilaró M, Cortés J, Selva-O'Callaghan A, Urrutia A, Ribera J-M, Cardellach F, et al. Adherence to reporting guidelines increases the number of citations: the argument for including a methodologist in the editorial process and peer-review. *BMC Med Res Methodol* 2019;19:112.
- [35] Pouwels KB, Widyakusuma NN, Groenwold RH, Hak E. Quality of reporting of confounding remained suboptimal after the STROBE guideline. *J Clin Epidemiol* 2016;69:217–24.
- [36] Stevens A, Shamseer L, Weinstein E, Yazdi F, Turner L, Thielman J, et al. Relation of completeness of reporting of health research to journals' endorsement of reporting guidelines: systematic review. *BMJ* 2014;348:g3804.
- [37] Blanco D, Altman D, Moher D, Boutron I, Kirkham JJ, Cobo E. Scoping review on interventions to improve adherence to reporting guidelines in health research. *BMJ Open* 2019;9:e026589.
- [38] Fuller T, Pearson M, Peters J, Anderson R. What affects authors' and editors' use of reporting guidelines? Findings from an online survey and qualitative interviews. *PLoS One* 2015;10:e0121585.
- [39] Glasziou P, Altman DG, Bossuyt P, Boutron I, Clarke M, Julious S, et al. Reducing waste from incomplete or unusable reports of biomedical research. *Lancet* 2014;383:267–76.

Chapter 6: Use of causal claims in observational studies

6.1. Background

Researchers may be tempted to draw causal conclusions from observational data despite acknowledging that mere association is not causation [50, 51].

To find out whether and how statements in study reports present confusing use of causal language (or lack thereof), we examined research articles reporting an observational study published in The BMJ in 2018. Our focus was on the causal message The BMJ readers receive from these papers, particularly from the abstract. We evaluated the consistency of causal statements in the published abstracts of observational studies, whether what was reported in the abstract was consistent with the full text and if any changes had been made as a result of the peer review process.

6.2. Methods

Sampling and inclusion criteria: we identified all original research articles published in The BMJ in 2018 described as either cohort or longitudinal studies in the study design section of the abstract. Studies identified as “observational” in the abstract were included if they suggested a period of follow-up rather than being cross-sectional. Articles described as case-cohorts were excluded as their interpretation and analysis differs from other studies with follow-up assessing the exposure-outcome relationship.

Assessment of published abstracts: for each abstract included, two reviewers independently assessed the text included under each of the subheadings (objective, design, setting, participants, outcome, results, conclusion). They determined whether there was an (implicit) causal (cl)aim using a yes/no/unclear response. Finally, each reviewer gave an overall assessment of the main claims in the abstract as either ‘consistently causal’, ‘inconsistent’ or ‘consistently not causal’. The overall rating of the abstract was compared between both reviewers; where there was disagreement, a third reviewer (EG) was consulted.

Assessment of published full text: We further evaluated the full published text of all eligible articles to identify the statistical methods applied and any further causal claims, looking for statements that would support or undermine a causal aim.

Assessment of initially submitted abstract version: we further assessed articles judged as ‘inconsistent’ to check if there were changes introduced to the manuscript between submission and publication, leading to this inconsistent use of causal language. As before, the assessments were

compared and, in cases of disagreement, a third reviewer (EG) was consulted and consensus reached.

6.3. Results

Selection and assessment of published abstracts: 60 research papers met our inclusion criteria. We identified 29 studies (48%) reporting causal language consistently. A further twelve (20%) studies were considered inconsistent. Finally, 19 papers (32%) were considered consistently not causal.

Assessment of published full text: We found that all articles classified as ‘consistently causal’ based on the abstract, also used causal language and contained causal statements in the full text. Conversely, more than half (11/19) of the abstracts classified as ‘consistently not causal’, contained causal language in the full text. In such articles, the authors applied causal methods, discussed residual confounding, biological plausibility or a dose-response relationship suggesting a causal aim. Looking at the methods section in the full text of the studies whose abstract was classified as ‘inconsistent’, we found that 11 of the 12 provided adjusted estimates. Most of the studies (8/12, 67%) used outcome regression models, mainly Cox proportional hazard models, or (propensity score) matching (3/12, 25%).

Assessment of submitted abstract version: Of the 12 published abstracts classified as ‘inconsistent’, we further classified 11/12 (92%) as also inconsistent on submission. There was only one study where the submitted version of the abstract described a different type of association. In this case, in the published version the term “impact” in the results was replaced by “association”.

6.4. Discussion

Statement of principal findings We found that the majority (80%) of the abstracts of observational studies we examined had a consistent use of causal language in the abstract. We think that beyond a deeper attention on the wording, readers can learn much about the aim of a paper, after the interpretation from described statistical methods, and assumptions made explicit in the manuscript. For authors, we think it is better to be transparent about an aim to draw a causal conclusion from a study and to acknowledge to fall short of that, than to generate confusion.

Implications: As observational data resources abound, methods for causal inference from observational data have surged in tandem with the call for real world evidence. The new

opportunities bring new challenges and the responsibility for clear and well supported statements on the evidence. Further research is needed to explore how causal claims presented in the abstract and full text are supported by the design and methods applied, which entails assessing the methods used and evaluating whether the underlying assumptions were met.

A prerequisite for good causal language practice includes awareness of which language implies a causal statement and which does not. Words like “effect”, “impact”, “determinant of”..., inevitably point in the causal direction and their use should come with the requirement of at least stating and ideally critically evaluating the necessary assumptions[48]. Uncertainty on the causal nature of the conclusion should tone down any suggestion for intervening on the studied exposure. Our suggestion is to be specifying the corresponding level of evidence, rather than hiding the ultimate causal aim of a study. In the conclusion the margin of error typical of any empirical study should be acknowledged.

6.5. Article

PREPRINT

Camila Olarte Parra, Lorenzo Bertizzolo, Sara Schroter, Agnès Dechartres, Els Goetghebeur

Use of causal claims in observational studies: a research on research study

medRxiv 2020.09.17.20194530

doi: <https://doi.org/10.1101/2020.09.17.20194530>

Use of causal claims in observational studies: a research on research study

Camila Olarte Parra^{1,2}, Lorenzo Bertizzolo^{2,3}, Sara Schroter⁴, Agnès Dechartres⁵, Els Goetghebeur¹

1. Department of Applied Mathematics, Computer Science and Statistics, Ghent University, Ghent, Belgium
2. Department of Clinical Epidemiology, Biostatistics and Bioinformatics, Academisch Medisch Centrum, Amsterdam, The Netherlands
3. INSERM, U1153 Epidemiology and Biostatistics Sorbonne Paris Cité Research Center (CRESS), Methods of therapeutic evaluation of chronic diseases Team (METHODS), Paris, F-75004 France; Paris Descartes University, Sorbonne Paris Cité, France.
4. The BMJ, London, United Kingdom
5. Sorbonne Université, Institut National de la Santé et de la Recherche Médicale (INSERM), Institut Pierre Louis d'Epidémiologie et de Santé Publique, AP-HP. Sorbonne Université, Hôpital Pitié Salpêtrière, Département de Santé Publique, F75013, Paris, France

Correspondence to: Camila Olarte Parra
Department of Applied Mathematics, Computer Science and Statistics
Ghent University, Ghent, Belgium
Email: camila.olarteparra@ugent.be, colartep@gmail.com

Abstract

Objective: To evaluate the consistency of causal statements in the abstracts of observational studies published in The BMJ.

Design: Research on research study.

Data source: All cohort or longitudinal studies describing an exposure-outcome relationship published in The BMJ during 2018. We also had access to the submitted papers and reviewer reports.

Main outcome measures: Proportion of published research papers with 'inconsistent' use of causal language in the abstract. Papers where language was consistently causal or non-causal were classified as 'consistently causal' or 'consistently not causal', respectively; those where causality may be inferred were classified as 'suggests causal'. For the 'inconsistent' papers, we then compared the published and submitted version.

Results: Of 151 published research papers, 60 described eligible studies. Of these 60, we classified the causal language used as 'consistently causal' (13%), 'suggests causal' (35%), 'inconsistent' (20%) and 'consistently not causal' (32%). The majority of the 'Inconsistent' papers (92%) were already inconsistent on submission. The inconsistencies found in both submitted and published versions was mainly due to mismatches between objectives and conclusions. One section might be carefully phrased in terms of association while the other presented causal language. When identifying only an association, some authors jumped to recommending acting on the findings as if motivated by the evidence presented.

Conclusion: Further guidance is necessary for authors on what constitutes a causal statement and how to justify or discuss assumptions involved. Based on screening these abstracts, we provide a list of expressions beyond the obvious 'cause' word which may inspire a useful more comprehensive compendium on causal language.

Strengths and limitations of this study

- We present examples of ambiguous causal statements in published abstracts of observational studies in a high impact journal
- We focused on the abstract where clear messages are especially important, as many readers just read the abstract of a study
- The focus on the abstract may miss further discussion on the validity of underlying assumptions justifying causal inference in the setting studied.
- The prevalence and nature of the problems found is a call for better instruction on and consideration of causal language throughout the editorial process in clinical and epidemiological research.
- We provide a list of words and study elements that could point in the direction of causality or otherwise, which may inspire a more comprehensive compendium.

Introduction

Many researchers remain tempted to draw causal conclusions from observational data despite acknowledging that mere association is not causation because causal inference is the ultimate goal of most clinical and public health research (1, 2). Gold-standard answers are typically sought through randomized controlled trials (RCTs). The unique ability of RCTs to avoid confounding bias (3) has led to demands that empirical research must be drawn from randomized studies to justify causal statements (4-6). RCTs are mainly used to assess the effect of a treatment or intervention but are not easily adapted to evaluate prognostic or risk factors rather than interventions.

There are however good reasons to look beyond RCTs for evidence on treatment effects. In many settings, RCTs are not feasible, ethical or timely and thus observational data are all that is available for some time, as in the recent COVID-19 crisis. Furthermore, observational studies typically involve broader real-world contexts than RCTs, where the costs and risks of experimentation suggest studying high risk patients without major comorbidities (7). This selection challenges generalization to the target population. Trials further suffer from treatment non-compliance which complicates analysis, as treatment-specific populations lose the benefit of randomization. Recent ICH9 guidelines therefore emphasize the importance of causal estimands beyond intention-to-treat, such as per-protocol and as-treated analysis (8, 9).

Deliberately avoiding causal statements on a hoped-for causal answer brings ambiguity and contrived reporting (10, 11). Instead, authors should openly discuss the likely distance in meaning and magnitude between the data based measure they are able to estimate and the desired targeted causal effect. Arguments would consider study design with additional assumptions in context (12). Owing to decades of progress in statistical science (involving potential outcomes, directed acyclic graphs, propensity scores and more) (13), this allows for results, often unreachable by randomized trials, with a justified causal interpretation (14).

In 2010, Cofield et al (5) assessed the use of causal language in observational studies in nutrition but deemed causal language inappropriate for all observational studies. From a different angle, Haber et al (15) examined whether the tone and strength of causal claims made in a given paper matched the language describing the findings in social media. Not surprisingly they found stronger causal statements in the media in half of the cases, emphasizing the importance of clear scientific messages.

To promote this, Lederer et al (16) recently published a guide for authors and editors on how to report causal studies in Respiratory, Sleep and Critical Care Journals. Rather than circumventing the problem by asking to avoid causal language, they provide key elements that ensure valid causal claims (17). Besides briefly explaining causal inference, they provide a definition of a confounder, outline how to identify confounding through so-called directed acyclic graphs and discuss how p-values are often misinterpreted and how their value does not reflect the magnitude, direction or clinical importance of

a given association. All these elements empower their target audience to critically assess observational studies.

To find out whether and how statements in study reports present confusing use of causal language (or lack thereof), we examined abstracts of research papers concerned with exposures and outcomes published in The BMJ in 2018. Our focus was on the causal message The BMJ readers receive from the abstracts. We evaluate the consistency of causal statements in the abstracts of observational studies published abstract and if any a priori changes had been made as a result of the peer review process.

Methods

Sampling and inclusion criteria

COP identified all original research articles published in The BMJ in 2018 described as either cohort or longitudinal studies in the study design of the abstract. The eligible studies were identified by statements in this section of the abstract such as “cohort”, “longitudinal” or “registry-based”. Those identified as “observational” were included if they suggested a period of follow-up rather than being cross-sectional. Articles described as case-cohorts were excluded as their interpretation and analysis differs from other studies with follow-up assessing the exposure-outcome relationship.

Assessment of published abstracts

Two reviewers (COP, LB) independently screened all the abstracts of the eligible papers. For the text included under each of the subheadings in the abstract (objective, design, setting, participants, outcome, results, conclusion), the reviewers assessed whether there was an (implicit) causal claim using a yes/no/unclear response. After assessing each separate subheading, each reviewer then gave an overall assessment of the main claims in the paper’s abstract as either ‘consistently causal’, ‘suggests causal’, ‘inconsistent’ or ‘consistently not causal’. After the independent assessments, the overall rating of the abstract was compared between both reviewers; where there was disagreement, a third reviewer (EG) was consulted and a consensus reached.

Assessment of submitted versions

As the focus of this paper is the avoidance of misleading and ambiguous messages, we further assessed those articles judged as ‘Inconsistent’ to see if there were changes introduced to the manuscript between submission and publication. For this subset, we obtained the submitted version of the manuscripts and the associated peer reviewers’ comments from The BMJ’s manuscript tracking system. We then compared the published version with the first submitted version to identify whether

the wording related to causal claims appeared in the submitted version of the abstract and whether changes occurred as a result of comments from peer reviewers and editors.

The same reviewers (COP, LB) independently evaluated the submitted versions of the papers. The reviewers assessed whether the content under each subheading of the submitted abstract differed from the published version. Where there were discrepancies between versions, each reviewer indicated the presence of a causal claim as yes/no/unclear for each abstract subheading (title, objective, design, setting, participants, outcome, results, conclusion) and made an overall assessment of the submitted abstract as either 'consistently causal', 'suggests causal', 'inconsistent' or 'consistently not causal'. As before, the assessments were compared and, in cases of disagreement, a third reviewer (EG) was consulted and consensus reached.

Assessment of full text

For the published papers classified as 'inconsistent', we further evaluated the full published text to identify the statistical method applied. We look for statements that would support a causal aim, including confounding adjustment, sources of bias and issues of generalizability.

Ethics and consent

This study used routinely collected data. When authors and reviewers submit manuscripts and reviews to The BMJ, they are notified that their paper or review may be entered into research projects for quality improvement purposes. COP was given access to The BMJ's data under a confidentiality agreement.

Patient and public involvement

Patients were not involved in the design, analysis or interpretation of the study. Patients were not participants in this study; it was a methodological study (research on research). Patients' opinions of causal statements and the use of ambiguous language in research papers is important and further work in this area partnered by patients is important.

Results

Assessment of published abstracts

In 2018, 151 research papers were published in The BMJ, of which 60 (40%) were eligible for inclusion in our study (Figure 1). We identified 29 studies (48%) using causal language ('consistently causal' and 'suggests causal'). A further twelve (20%) abstracts were considered inconsistent mainly because the objective stated evaluating an association while the conclusion presented a causal finding (9/12) or the opposite (3/12). Finally, there were abstracts that described studies that aim for prediction or reported associations without (implicitly) suggesting that they had a causal nature that were considered consistently not causal (n=19, 32%).

Tables

Table 1 shows excerpts from the abstracts that were evaluated. Each row corresponds to statements from the same study. The first column indicates the assigned category, based on the type of association it describes. The last column explains why a given abstract was considered to belong to the assigned category. As the assessment pertains to causal claims in general, the words referring to the particular topic of the corresponding study were removed from the statements. The examples shown are not an exhaustive list, but were chosen to illustrate the different phrasing of statements belonging to the different categories. It is worth noting that the statements presented correspond to the objective and conclusion subheading of the abstract. When assessing the abstracts, we identified that these were the subheadings under which the information to classify the abstract was mainly found. Other subheadings like Design, Setting and Participants were not as relevant for this purpose.

To further illustrate how statements in these two sections can be misleading, we tabulated a few examples in a 2 by 2 table showing mismatches between what was reported in the objectives and conclusion resulting in the paper being categorised as either 'consistently (not) causal' or 'inconsistent' (Table 2).

Assessment of submitted versions

After evaluating the first submitted version of the 12 abstracts classified as 'inconsistent', we classified 11/12 (92%) as also inconsistent on submission. There was only one study where the submitted version described a different type of association. In this case, the conclusion of both the submitted and published versions was rather conservative by stating that the intervention was "independently associated" with the outcome. The submission expressed a causal objective, stating the aim of evaluating the "impact" of a particular intervention with corresponding methods: providing adjusted estimated effects and including sensitivity analysis using propensity score matching. However, in the published version the term "impact" was replaced by "association" making the abstract less clear

about a causal aim because both the objectives and the conclusion described an association but the authors still provided adjusted hazard ratios and resorted to propensity score matching.

Assessment of full text

Looking at the methods used in the papers classified as 'inconsistent', we found that 11 of the 12 provided adjusted estimates. Most of the studies (8/12, 67%) used outcome regression models, mainly Cox proportional hazard models, or (propensity score) matching (3/12, 25%). Table 3 presents statements found in both the published abstract and full text of each of these papers regarding the method used and considerations suggesting a causal aim.

Discussion

Statement of principal findings

We found that the majority of the published research paper abstracts of observational studies had a consistent use of causal language. Still 20% of them contained inconsistent messages on the causal nature of the key "effect". Inconsistencies showed up in two directions: an intentional quest for causality ending in uncriticized non-causal conclusions or carefully phrased mere associations ending with recommendations to act and intervene based on the exposure outcome association.

Beyond the wording in the abstract, readers can learn much about the sought, after interpretation from described statistical methods, and assumptions made explicit in the paper. On a case by case basis, one could then assess whether additional assumptions, e.g. involving 'no-unmeasured confounders', would justify the causal assessment derived from these approaches. Identifying key elements like the ones presented in Table 3 would help to assess if causal inference is possible. If in doubt, a sensitivity analysis may be in order. It seems better to be transparent about the ultimate aim to draw a causal conclusion and to acknowledge to fall short of that, than to generate confusion.

Comparison with other studies

This is not the first study to evaluate the use of causal language in the medical literature. Cofield et al (5) assessed the use of causal language in observational studies in nutrition. However, they reduced the problem to assessing whether authors included causal language or not, as it was deemed inappropriate due to the observational nature of the study. We have made the case that merely avoiding explicit causal terms is not a real solution. Even without them, a causal conclusion is implicit when the take home message encourages interventions based on the presented findings. Avoiding inconsistency is important but equally one should be able to trust that the use of consistent causal

language is not in vain. This requires a more in depth look at methods and assumptions validating the casual claims.

Strengths and weaknesses of the study

Accurate abstracts are important. In just a few brief paragraphs, the author summarizes key elements of design, methods and results, and comes to a conclusion. Many readers only read the abstract. However, a powerful abstract opens the door to readers and sets the scene for any study. It serves the different roles of informing the audience about its main findings while motivating the reader to further explore the full text, all within the constraints of brevity. This demands from authors special attention to ensure that every word in the abstract is required. All of the above makes the assessment of the abstract relevant but also challenging.

Further research is needed to explore how causal claims presented in the abstract are supported by the full article, which entails assessing the methods used and evaluating whether the underlying assumptions were met (18). The ultimate conclusion should not simply label a study as black or white in causal terms. In the present study we used a convenient limited number of classifications for short statements. In practice a continuous degree of confidence in a potential causal relationship is likely to emerge based on observed association.

We are aware that by limiting our assessment to the abstract, we may have missed the discussion of the extent to which the underlying assumptions that enable causal inference were met. Indeed, when there was a clear causal aim but the authors considered that these assumptions were not fulfilled, they may have decided that a causal claim was inappropriate and phrased their conclusion in terms of association rather than causation. If this is the case, the apparent inconsistency found in the abstract would no longer hold. On the contrary, any undue causal claims can be viewed as a form of spin (19, 20).

Conclusions and policy implications

As observational data resources abound, methods for causal inference from observational data have surged in tandem with the call for real world evidence. The new opportunities bring new challenges and the responsibility for clear and well supported statements on the evidence. In this spirit and motivated by novel guidelines as proposed by ICH9 and FDA, Miguel Hernan and collaborators have embarked on a project entitled “Developing Guidelines for the Analysis of Randomized Controlled Trials in Real-World Settings”(21). The importance of such initiatives, supports a shift towards being explicit and discussing assumptions underlying causal methods that allow for causal interpretations in context, with or without an RCT (13). In the meantime, uncritical ambiguous phrasing in observational studies remains prevalent (14). Those searching for the best possible evidence supporting future treatment decisions, are best served by transparent reports on observational studies.

Faced with uncertainty when concluding on the nature of the observed exposure outcome relationship, a justifiable balance between the type I and II error rate is a natural guide for action. The cost of errors must be weighed in context, for instance as in clinical trials emphasizing control of the type I error to avoid introducing new unhelpful drugs at a potentially large cost. Alternative weights are typical in screening programs where false positives will be caught in follow-up examinations, but false negatives are lost forever. In a crisis, such as the current COVID-19 pandemic, we must act before long term randomised trials have materialised. It becomes undeniably important to learn as much as we can from observational data, be aware of the types of risk when acting or not, as displayed in the schematic Table 4.

A prerequisite for good causal language practice includes awareness of which language implies a causal statement and which does not. To support correct phrasing and raise awareness, we have compiled a short list of words and expressions with dedicated (non) causal meaning (Box). The list draws on phrases found in our study and in the references cited, particularly Hernan et al (10) and Thapa et al (6). We consider that a definition of causal language that is generally recognized by the research community is needed (22, 23).

Words like “effect”, “impact”, “determinant of”..., inevitably point in the causal direction and their use should come with the requirement of at least stating and ideally critically evaluating the necessary assumptions (6). Uncertainty on the causal nature of the conclusion should tone down any suggestion for intervening on the studied exposure. Specifying the corresponding level of evidence rather than hiding the ultimate causal aim of a study is what we recommend (19), while acknowledging a margin of error in any empirical study (20).

In summary, we have found that causal messages are embedded in studies otherwise carefully phrased in terms of association. Further guidance for authors appears needed on what constitutes a causal statement, similar to the one published by Lederer et al (13) for Respiratory, Sleep and Critical Care Journals. We look forward to similar guidance for other disease groups. From the screened BMJ abstracts, we provided a list of expressions with clear interpretation which may inspire a useful more comprehensive compendium. We argue that such awareness and special attention amongst authors and reviewers would serve our communication on the best available evidence for conceived interventions.

Box. Examples of words and study elements that could point to causality or otherwise.

Words expressing a causal relationship	<ul style="list-style-type: none"> • Affect • Attributable • Benefit • Cause • Contribute • Determinant • Efficacy • Impact • Improve • Leads to • Responsible for • Results in • Safety
Words that could suggest causality in a given context	<ul style="list-style-type: none"> • Effect • Induce • Higher (lower) probability • Modify • Risk (factor)
Specific expressions avoiding suggestions of casual effects	<ul style="list-style-type: none"> • Association • Correlation • Less (more) likely link • Predict • Pattern
Key aspects suggesting causal aim	<ul style="list-style-type: none"> • Adjusting for confounders • Discussing 'unmeasured confounders' assumption • Suggesting/recommending intervention • Using instrumental variables • Propensity score adjustment

Contributorship statement

- Conceptualization: COP, LB, AD, SS, EG
- Methodology: COP, EG
- Data curation: COP, LB
- Formal analysis: COP, LB, EG
- Writing original draft: COP
- Writing-review and editing: LB, SS, AD, EG
- Approved final version: COP, LB, AD, SS, EG

Competing interests declaration

SS is a full-time employee of the BMJ Publishing Group. No other conflicts of interest to declare.

Funding

This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 676207.

Patient and public involvement

Patients were not involved in the design, analysis or interpretation of the study. Patients were not participants in this study; it was a methodological study (research on research). Patients' opinions of causal statements and the use of ambiguous language in research papers is important and further work in this area partnered by patients is important.

Ethics Approval

This study used routinely collected data. When authors and reviewers submit manuscripts and reviews to The BMJ, they are notified that their paper or review may be entered into research projects for quality improvement purposes.

Transparency declaration

COP confirms that this manuscript is an honest, accurate, and transparent account of the study being reported and that no important aspects of the study have been omitted.

Data sharing statement

The reviews and published versions of the papers included in the study are publicly available at BMJ.com. No further data will be made available as it is confidential submission data.

References

1. Begg MD, March D. Cause and association: Missing the forest for the trees. American Journal of Public Health. 2018;108(5):620-.
2. Hernán MA. A definition of causal effect for epidemiological research. Journal of Epidemiology and Community Health. 2004;58(4):265-71.
3. Kunz R, Oxman aD. The unpredictability paradox: review of empirical comparisons of randomised and non-randomised clinical trials. Bmj. 1998;317(October):1185-90.
4. Ruich P. The use of cause-and-effect language in the JAMA Network journals. 2017.
5. Cofield SS, Corona RV, Allison DB. Use of causal language in observational studies of obesity and nutrition. Obesity Facts. 2010;3(6):353-6.
6. Thapa DK, Visentin DC, Hunt GE, Watson R, Cleary M. Being honest with causal language in writing for publication. Journal of Advanced Nursing. 2020;76(6):1285-8.
7. Pinsky PF, Miller A, Kramer BS, Church T, Reding D, Prorok P, et al. Evidence of a healthy volunteer effect in the prostate, lung, colorectal, and ovarian cancer screening trial. American journal of epidemiology. 2007;165:874-81.
8. Food Drug Administration (FDA), Permutt T, Scott J. E9(R1) Statistical Principles for Clinical Trials: Addendum: Estimands and Sensitivity Analysis in Clinical Trials. 2017;9.
9. European Medicine Agency (EMA). ICH E9 (R1) addendum on estimands and sensitivity analysis in clinical trials to the guideline on statistical principles for clinical trials 2020 17 February 2020
10. Hernan MA. The C-Word : Scientific Euphemisms Do Not Improve Causal Inference From Observational Data. 2018;108(5):616-9.
11. Glymour MM, Hamad R. Causal thinking as a critical tool for eliminating social inequalities in health. American Journal of Public Health. 2018;108(5):623-.
12. Green MJ. Calculating Versus Estimating Causal Effects. American Journal of Public Health. 2018;108(8):e4-e5.

13. Hernán M, Robins J. Causal Inference: What If: Boca Raton: Chapman & Hall/CRC; 2020.
14. Goetghebuer E, Cessie SI, De Stavola B, Moodie E, Waernbaum I. Formulating causal questions and principled statistical answers. 2019.
15. Haber N, Smith ER, Moscoe E, Andrews K, Audy R, Bell W, et al. Causal language and strength of inference in academic and media articles shared in social media (CLAIMS): A systematic review. PLoS ONE. 2018;13(5):1-21.
16. Lederer DJ, Bell SC, Branson RD, Chalmers JD, Marshall R, Maslove DM, et al. Control of Confounding and Reporting of Results in Causal Inference Studies. Guidance for Authors from Editors of Respiratory, Sleep, and Critical Care Journals. Annals of the American Thoracic Society. 2019;16(1):22-8.
17. Harhay MO, Au DH, Dell SD, Gould MK, Redline S, Ryerson CJ, et al. Methodologic Guidance and Expectations for the Development and Reporting of Prediction Models and Causal Inference Studies. Annals of the American Thoracic Society. 2020;17(6):679-82.
18. Hernán MA, Robins JM. Estimating causal effects from epidemiological data. Journal of Epidemiology and Community Health. 2006;60(7):578-86.
19. Boutron I. Reporting and Interpretation of Randomized Controlled Trials With Statistically Nonsignificant Results for Primary Outcomes. JAMA. 2010;303(20):2058-.
20. Ghannad M, Olsen M, Boutron I, Bossuyt PM. A systematic review finds that spin or interpretation bias is abundant in evaluations of ovarian cancer biomarkers. Journal of Clinical Epidemiology. 2019;116:9-17.
21. Developing Guidelines for the Analysis of Randomized Controlled Trials in Real-World Settings [Available from: <https://www.pcori.org/research-results/2015/developing-guidelines-analysis-randomized-controlled-trials-real-world>].
22. Doan S, Yang EW, Tilak SS, Li PW, Zisook DS, Torii M. Extracting health-related causality from twitter messages using natural language processing. BMC Medical Informatics and Decision Making. 2019;19(Suppl 3).
23. Chiolero A. Causality in public health: One word is not enough. American Journal of Public Health. 2019;109(10):1319-20.

Figures

Figure 1. Adapted Prisma flow diagram showing study selection and main study findings.

Tables

Table 1. Examples of statements found in the objectives and conclusions of abstracts of observational studies published in The BMJ in 2018 and their corresponding assigned category.

Assigned category	Objective	Conclusion	Comment
Consistently causal	"...assess the effectiveness of..."	"Little evidence was found of a direct impact of..."	When discussing associations, words like effect, contribution or role are similar to cause and then (direct) impact and effect will be their consequence
	"To determine the effect of ... in ..."	"...has led to risk..."	
	"To describe the contributions of..."	"... an important role in ..."	
	"To evaluate the impact of ..."	"...impacts are..."	

	"To investigate whether improving adherence to"	"...the beneficial effect of improved..."	Evaluates taking an action 'improving adherence' and concludes that the effect is beneficial
	"...benefit of ... in reducing ... risk"	"... is an overlooked risk factor for ..."	Evaluates how a given intervention can reduce the risk of an outcome and then labels it as an 'overlooked risk factor'
	"To determine outcomes and safety of..."	"... is at least as effective and safe as ..."	Evaluates and determines that a certain intervention is as safe as the comparator
	"to quantitatively decompose this joint association to ... only, to ... only, and to their interaction."	"...excess risk of...These findings suggest that most cases of ... could be prevented by ..."	Suggests interest in direct and indirect effect i.e. mediation analysis, and concludes consequently
Consistently not causal (Associations)	"...is associated with ...compared with..."	"...is associated with ...compared with..."	Describes associations without labelling them as causal or prediction

	"To describe trends in..."	"...rates were high during the study period of ... with the highest rates in ... versus ..."	Limits to describe frequency
	"To assess how often ..."	"One in ... adults ... were ..."	
	"To examine the association between..."	"...could increase ... confirmation of these findings are warranted, preferably in an intervention setting."	Suggests further research to determine the nature of the association
	"...compared with...is associated with..."	"Additional studies, with long term follow-up, are needed to investigate the effects of..."	
Consistently not causal (Prediction)	"To develop and validate a set of practical prediction tools that reliably estimate the outcome of..."	"...prediction models reliably estimate the outcome..."	Describes developing and validating prediction models

	"To prospectively validate the ... algorithm to ..."	"...accurately classified..."	
Inconsistent	"...evaluate safety of..."	"...associated with..."	Phrasing the objective as causal and limiting to describing an association in the conclusion
	"...analyse the effect of..."		
	"...critical determinant..."		
	"...association with..."	"...is safe..."	Phrasing the objective as just to explore an association and presenting a causal claim in the conclusion
		"... had no substantial effect on long term survival..."	
		"... was determined by... may be largely explained by..."	

		"... was found to be the safest drug, with reduced risks of..."	
		"These results emphasise the benefit of..."	
	"...association with..."	"...tackling all these risk factors might substantially..."	Phrasing the objective and conclusion as if just to assess an association but then suggesting to take action given the findings
		"...Targeting ... prevention strategies among these patients should be considered."	
		"Systematically addressing ... may be an important public health strategy to reduce the incidence of"	

		"...present findings encourage the downward revision of such guidelines ..."	
--	--	--	--

Table 2. Examples of (mis)matching causal and non-causal statements found respectively in the objectives and conclusions of abstracts of observational studies published in The BMJ in 2018.

		Conclusion	
		Causal	Not causal
Objective	Causal	<p>Consistent</p> <p>"...assess the effectiveness of..." and "Little evidence was found of a direct impact of..."</p> <p>"...benefit of ... in reducing ... risk" and "... is an overlooked risk factor for ..."</p>	<p>Inconsistent</p> <p>"...evaluate safety of..." and "...associated with..."</p> <p>"...analyse the effect of..." and "...associated with..."</p> <p>"...critical determinant..." and "...associated with..."</p>
	Not causal	<p>Inconsistent</p> <p>"...association with..." and "...is safe..."</p> <p>"...association with..." and "... had no substantial effect on long term survival..."</p>	<p>Consistent</p> <p>"To describe trends in..." and "...rates were high during the study period of ... with the highest rates in ... versus ..."</p> <p>"To assess how often ..." and "One in ... adults ... were ..."</p>

		<p>"...association with..." and "...tackling all these risk factors might substantially..."</p> <p>"...association with..." and "Systematically addressing ... may be an important public health strategy to reduce the incidence of"</p>	<p>"To develop and validate a set of practical prediction tools that reliably estimate the outcome of..." and "...prediction models reliably estimate the outcome..."</p>
--	--	---	---

Table 3. Statements in each of the 12 published observational studies classified as “inconsistent”

Abstract		Full text			
Objective	Conclusion	Method	Confounder adjustment	Estimates provided	Considerations
“To evaluate the relation between [...] and development of [...]”	“ [...] was associated with an increased risk of [...] that was mediated by [...]. Systematically addressing [...] may be an important public health strategy to reduce the incidence of [...] among patients with [...].”	“We calculated the hazard ratios for the relation of [...] to the risk of MRSA using Cox proportional hazard models. [...] We also calculated the absolute risk difference. [...] We performed mediation analyses to examine the extent to which the effect of [...] on the risk of [...] was through [...]. Using marginal structural models we then estimated the natural direct effect [...] and the natural indirect effect [...] while	“We performed a matched cohort study [...] matched on age (one year either way), sex, and study entry time (within one year either way). Such comparators were chosen to further ensure the comparability [...] In the multivariable Cox model we adjusted for [...].”	“The matched and multivariable adjusted hazard ratios for patients with [...] were 1.69 (1.51 to 1.90) for [...] and 1.26 (1.12 to 1.40) for [...].”	“Our GP practice based dataset could have missed the detection of some inpatient cases of [...]; however, these potential non-differential misclassifications would have biased our results towards the null, rendering our findings conservative”

		adjusting for the same confounding variables"			
"To quantify the effects of varying [...]"	"[...] is associated with a large increase in [...] among [...] patients. The data from this study suggest that [...] rather than [...] is more strongly associated with [...]"	"For adjusted analysis of time until [...] we used Cox proportional hazards models."	"Adjusted models included [...]"	"Each additional [...] increased the rate of [...] by 70.7% (95% confidence interval 54.6% to 88.4%) before adjustment and increased the hazard of [...] by 44.0% (40.8% to 47.2%, $P < 0.001$) after adjusting for covariates."	"To determine the extent to which strong unobserved confounding might explain the observed association, we included this synthetic confounder in a Cox model. [...] As part of a sensitivity analysis, we constructed models that removed potential confounders. [...]"
"To evaluate the long term association between [...]"	"Widespread utilisation of [...] may be contributing to long term increased risk of [...]. The potential for [...] should be considered when [...]"	"We conducted Poisson regression analyses using person years as observations."	"We included several variables as known confounders or effect modifiers in the relation between. [...]The final fully adjusted model adjusted for [...]"	"After adjustment for covariates, the rate ratio was [...] indicating that during the entire period of follow-up the risk of [...] was 21%"	"The registered active [...] population is generally representative of the UK population in terms of age, sex, and regional distribution"

				higher during [...] than at other times.”	
“To investigate the association of [...]”	“The shape of the association between [...] and [...] was determined by [...]. This finding suggests that the [...] may be largely explained by [...]”	“We used Cox proportional hazards models to estimate hazard ratios and 95% confidence intervals. We stratified the analysis by age in months and calendar year of the questionnaire cycle.”	“For the main analysis, we used [...] measured at baseline to minimize the effect of underlying diseases on mortality [...]. In multivariable models, we adjusted for potential confounders including [...]”	“A multivariable adjusted model showed a positive association between [...] and all cause mortality, whereas [...] showed a U shaped association with all cause mortality. In a mutually adjusted model including both [...] and [...] we consistently observed a strong positive association between [...] and all cause mortality.”	“Our findings remained robust in several sensitivity analyses [...] we cannot entirely rule out the possibility of unmeasured or unknown confounding factors that may account for the associations observed in this study.”
“ To examine the associations of [...]”	“This association could be explained by the finding that [...] These results emphasise the importance of revisiting [...] or establishing specific guidelines for	“We performed Cox models with penalised splines”	“In final Cox models with penalised splines, we made adjustments for: [...]”	“[A]fter adjustment for confounding factors, the U shaped association with [...]”	

	management among [...]"				
" To estimate long term survival, health, and educational/social functioning in patients with [...]"	"[...] had no substantial effect on [...]"	"We calculated mortality rate ratios and incidence rate ratios as measures of relative risk."	"For each [...] patient, we used the Danish Civil Registration System and the DNPR to identify all Danish residents with the same sex and date of birth as the patient who had not tested positive [...] and who met the study's inclusion and exclusion criteria [...]. From this population, we extracted 10 people at random for each patient. People in the population comparison cohort were assigned the same date of study inclusion as [...] patients to whom they were matched."	Patients and members of the comparison cohort were well matched with respect to [...] Mortality was not higher among patients in the [...] cohort"	

<p>"To compare the risk of [...]"</p>	<p>" Although residual confounding cannot be excluded, this finding deserves consideration when [...] is used for [...]"</p>	<p>"We estimated the crude hazard ratio of [...] using Cox proportional hazard regression, and the adjusted hazard ratio was obtained using propensity score matching"</p>	<p>" We identified potential confounders that were plausibly associated with both [...]based on clinical knowledge [...]. In the context of this study, the propensity score is the probability of receiving [...] as opposed to [...], given the baseline characteristics. Patients who received [...] were matched to patients who [...] using a 1:1 nearest neighbor matching algorithm with a caliper of 0.2 of the standard deviation of the propensity score on the logit scale. Covariate balance between the two groups was assessed after matching, and we considered an absolute standardized difference less than 0.1 as evidence of balance"</p>	<p>"The crude hazard ratio of death in the unmatched cohort was 1.51 (95% confidence interval 1.22 to 1.85) and the adjusted hazard ratio in the matched cohort was 1.50 (1.14 to 1.96)"</p>	<p>"Comparison of the baseline characteristics in the unmatched cohort provided little evidence of confounding [...]. It is unlikely that a few additional unmeasured variables can explain a 50% increase in the risk independent of all other confounder and proxies of confounders that are adjusted for in our study."</p>
---------------------------------------	--	--	--	--	--

<p>“To determine whether [...] is associated with [...]”</p>	<p>“[...] was independently associated</p> <p>With [...]”</p>	<p>“We fitted both a mixed effect logistic regression model (in which the outcome was defined as dichotomous [...] and the Prentice, Williams, and Peterson (PWP) model”</p>	<p>“We examined the relation between [...] and [...] adjusted for ...”</p>	<p>“The rate of distinct criteria met per year increased by 24% if a patient had been admitted to hospital (hazard ratio 1.24, 95% confidence interval 1.20 to 1.28) when controlled for the other covariates”</p>	<p>“We did a sensitivity analysis using propensity score matching to assess whether the association between [...] and [...] could be due to unmeasured confounders [...] Although we adjusted for a range of characteristics of patients, as with any observational study potential exists for unmeasured confounding, which may partly or fully explain the relation between [...]”</p>
<p>“To investigate associations between [...] and to analyse the effect of changes [...]”</p>	<p>“Risks of [...] are inversely associated with [...]”</p>	<p>“We used multivariable Cox regression analysis to compare the rates of [...] and [...].”</p>	<p>“Confounders included in the final models were based on the literature or statistical significance</p>	<p>“Compared with [...], [...] had increased hazard ratios of [...]”</p>	<p>“We believe that our findings are widely applicable and provide justification</p>

			(P<0.10). The full model included [...]"		for [...] and continuing [...]"
"To estimate the rates of [...]"	"In cases of [...], approximately [...] will become [...], of which a third will have [...]."	"We present denominators where data for the secondary outcome are missing. We defined the population attributable fraction as $(Re-Run)/(Re-(RR-1)/RR)$, calculated using Stata. To test the robustness of our findings, we did a sensitivity analysis."	"We compared the demographic and clinical variables of [...]. We used the binomial Wilson score to calculate confidence intervals of single proportions and the Pearson exact method to calculate confidence intervals of risk ratios and medians."	"[...] had a higher risk of [...]. The population attributable fraction of [...] was 47% for [...] and 61% for [...]"	"Considering these results when counselling potentially exposed [...] seems reasonable"
To perform an expedited assessment of [...] risk associated with exposure to [...]"	"The results do not imply a markedly increased short term overall risk of [...] in [...]."	"We used Cox regression to estimate the hazard ratio with 95% confidence intervals for [...] associated with [...], both for ever use and for the predefined	"Analyses were, however, performed as crude comparisons adjusted only for [...] as well as adjusted for [...] and the potential confounding factors."	"Overall, exposure to [...] showed no association with [...] compared with exposure to [...] (adjusted hazard ratio 1.09, 95% confidence interval 0.85 to 1.41)	"This ensured that the estimates were not affected by immortal time bias [...]. As all comparisons were performed within users of [...], the exposure to [...]"

		categories of cumulative use"		and no evidence of a dose-response relation"	can reasonably be expected to be a random event, and confounding is thus expected to be limited."
"To investigate the risks of [...] in [...]"	"No increased risk of [...] was detected in [...], but increased risks of [...] were found in this study. Our results suggest that [...] risks could be due to [...], rather than [...]."	"To calculate expected [...], we multiplied the person years at risk by corresponding national incidence rates (by 5 year age band and individual calendar year) for the general female population of England and Wales. Standardised incidence ratios were calculated by the comparison of observed values with expected values."	"We obtained data relating to potential confounding factors such as [...]"	"There was no overall increased risk of [...] (2578 observed v 2641.2 expected [...]; standardised incidence ratio 0.98 (95% confidence interval 0.94 to 1.01); absolute excess risk -2.8 cases per 100 000 person years (95% confidence interval -7.1 to 1.8); table 2)".	"Given previous inconsistent results, small study size, and lack of information on potential confounders, we undertook a population based linkage study in [...]"

Table 4. Impact of the errors of causal effect assignment.

		True nature of the main exposure effect	
		Causal	Not causal
Reported nature of the studied exposure effect	Causal	A true causal effect has been discovered. Recommendation to act on this should be considered. Language in the context of a study intended for causal inference.	Type I error: there is no causal effect, but it is claimed. Causal language used or suggestion to take action made when the purpose was to find associations
	Not causal	Type II error: hiding the true causal objective by avoiding use of causal language	No causal language when the objective is prediction or to explore associations

Chapter 7: Discussion and Implications

7.1. Discussion

7.1.1. Multidisciplinary approach

The number of scientific articles published has been steadily increasing in the recent years; in parallel we have unfortunately witnessed a rise in the research waste. Research waste can be created in every step of the research process. In this thesis we studied possible areas of research waste, happening at two different moments of such process: the conduct of the study and the reporting and worked on RoB and its assessment and the incomplete or inadequate reporting of findings.

The thesis reported five interconnected projects on these two topics. The two core projects dealt with the RoB assessment of RCTs. A third project, using a meta-epidemiological design, evaluated for a specific topic the association between RoB and other trial characteristics with treatment effect estimates. The two remaining projects focused on issues related to the reporting of findings in observational studies. In the first project on reporting, we worked on RGs. Through a survey we examined the knowledge and attitude of researchers regarding a specific RG dedicated to observational studies, STROBE. The last project of this thesis explored the use of inappropriate causal language in the reporting of observational studies.

This work in its thesis can be characterized by its multidisciplinary nature and approach. We dealt with different moments and aspects of the complex research process and this allowed us to both involve different specialists - such as methodologists, specialists of randomized trials and systematic reviews and meta-analyses and specialists of epidemiological studies - and partners such as Cochrane and journal editors. Risk of bias and quality of reporting are highly connected as shown in our first project where we found that the most common reason for discrepancies between reviewers was related to an incomplete or unclear reporting in the study report.

The projects on RoB assessment were centred on research methodologists and researchers in the field of evidence synthesis. In our findings we highlighted areas of intervention to improve the reproducibility of RoB assessment, which may be useful for conducting future systematic reviews. Moreover, the critical points in the RoB assessment that need attention by reviewers easily correspond to points that need to be improved in the reporting of RCTs. This probably makes our results also useful for researchers conducting RCTs.

The project on the association between a series of trial characteristics and the effect estimates in studies of hormone replacement therapy, used a meta-epidemiological approach and its results were aimed to researchers in the field of evidence synthesis, but also for specialists in the topic of interest.

The project on the use of STROBE involved researchers with experience in observational studies and journal editors. The project used a completely different design than previous works in this thesis, consisting in an online survey. The results of the STROBE project are immediately useful for the researchers involved in the survey, who are experts in the field of RGs, but they are also important for editors of scientific journals. We identified factors that may be very useful to improve the acceptance and use of RGs in the everyday routine of authors and editors.

The final project on the use of causal language directly involved journal editors. We believe the results are relevant for a wide audience: from one side researchers involved in observational studies should be aware of how their manuscripts may be read by the public. On the other side, the public, meaning all the readers of observational studies, have a direct interest in being able to recognise inappropriate causal claims in what they read. The results are also of interest for research methodologists, because there is need for research on how to correctly use causal language when reporting such study. Finally, the results are aimed at journal editors, because their review process should possibly also include the appraisal of proper use of causal claims.

7.1.2. Novelty and main results

The first two projects on RoB assessment differed from most previous literature on the topic. Firstly, our sample size was more than ten times larger than the single similar study on the topic. More importantly, we assessed the agreement in the RoB assessment in a real-life scenario by evaluating how the same RCTs were assessed for RoB in two different Cochrane reviews. Most of the studies on reproducibility of RoB published up to the time of our project consisted of comparisons of the independent RoB assessment conducted by two or more researchers in an experimental context. Such studies may not be representative of how the assessment works in real-life. Despite this important difference in the design, our results regarding agreement in RoB assessment did not differ too much from previous literature on the topic. We confirmed that agreement between different reviewers is suboptimal (overall agreement in our sample was about 70%), with differences across the different items of the RoB tool.

We believe that our study is unique in its in-depth exploration of reasons for disagreements. We systematically evaluated the support for judgement (“the justification”) for each judgement that led to a disagreement and we strengthened our analysis with the information we could gather from the full-text publications of individual studies and from the review including information about use of other sources such as protocol or contact with authors. This careful work allowed us to highlight that the most common cause of disagreement in RoB assessment is the difference in interpretation

of incomplete or unclear information in the study report. Disagreements caused by access to different or additional information were relatively infrequent, being in fact limited to certain items of the RoB tool. The results of our study showed the importance of providing further guidance to assess RoB in case of common situations of incomplete reporting ; at the same time our results underline the need to improve the quality and completeness of reporting especially regarding certain aspects of the study conduct of RCTs.

The second study on RoB assessment was directly inspired by the results of the first study. Here we decided to look at the issue of the RoB assessment agreement from a different angle, trying to explore the potential influence of “contextual factors”, which were defined as characteristics that are not part of the RoB assessment process, on the result of such assessment. To our knowledge, no other study evaluated the possible influence of such contextual factors on RoB assessment. We showed that such factors may exist and have an influence. We highlighted the importance of factors that may seem obvious, such as being among the authors of the study under scrutiny, but also less obvious aspects, such as the experience of authors in having conducted more SRs in the past. We think it was very important to draw the attention of researchers to the existence of such factors, which are not part of the RoB tool itself. A better knowledge of such factors may improve reproducibility of the RoB assessment in the future.

The meta-epidemiological study on testosterone replacement therapy used this approach to evaluate whether some factors may explain discrepancies in the effect estimates presented in the literature on this specific topic. In such a narrow topic, we found no evidence that RoB was associated with treatment effect estimates. This was probably due to the small sample size, but we also noticed a low prevalence of studies at low RoB in our sample.

In the last two studies we decided to shift from issues in research conduct to issues in the reporting of research. The study on STROBE had a direct approach, surveying authors of observational studies. The participants to the study were identified through a comprehensive strategy involving a series of consecutive steps and with the direct involvement of journal editors. Our results were interesting because we highlighted the key role of scientific journals in improving the use and acceptance of RGs. Journals could consider recommending the use of RG or require their use with a jointed submission of the filled checklist with the main manuscript.

The final project reported in this thesis, evaluating the inappropriate use of causal language when reporting observational studies, concerned a less-often explored topic. We decided to use the abstracts and then delve into the full-text manuscripts of a well-known journal. Our approach was quite explorative in its nature and we decided to provide readers with plenty of examples that will serve as good or bad examples of how to report such in observational studies and material for

further research on the issue. In our conclusion, we again suggested that a more active role of editors and reviewers could help avoiding the misuse of causal claims in observational studies.

7.2. Implications

The projects reported in this thesis, although they differed in topic, design and implications, share a common theme. This common message across different projects is the importance of conducting research on methods to improve the quality of research methodology and how such research should address every phase of the research process.

Throughout our projects, we identified several areas for intervention. The different areas concern specific aspects and elements of the RoB assessment, of the dissemination and acceptance/use of RGs and the need for more research and guidance on how to avoid improper causal claiming while reporting observational research.

In the first project on RoB assessment we found that the main cause of disagreement was dealing with incomplete or unclear information from the primary study report. The implication of this finding was dual. From one side there is the necessity to improve the reporting, especially in regard of certain aspects of the study conduct, such as the randomization process and how researchers dealt with missing data. From the other side, the process of RoB assessment could be improved and more guidance could be given to assessors on how to deal with common cases of incomplete information. Such cases were well outlined in the results.

Recently an important change has been introduced in the RoB assessment and some of its implication may move in the direction we pointed at through our project. Cochrane in 2019 launched a new version of the tool [55], which was again recently updated (November 2020)[56]. The new tool is characterized by a different approach to the RoB assessment process, adopting a step-by-step approach using “signalling questions”. Such approach was accompanied with decision algorithms, which could provide the increased guidance that was lacking before. The use of signalling questions and algorithms could therefore improve the reproducibility of the RoB assessment. Nevertheless, the new tool does not answer all the questions we raised, for example, more guidance on assessment of how researchers dealt with missing data or what “quota” of missing data could be considered acceptable for RoB purposes. The new RoB tool has been just recently released and researchers involved in evidence synthesis are starting using it only in recent time. Future research is necessary to evaluate whether the important changes introduced by new tool will improve the reproducibility of RoB assessment.

The second project shared light on new aspects and subsequent implications of the RoB assessment process. In our results we hinted at the existence of factors that are not part of the RoB tool and the RoB assessment, but which may nevertheless influence reviewers in their judgement. The project opened to a new area of research on factors influencing RoB assessment, while also giving

important clues for reviewers. Reviewers should be aware that their judgement, even if methodologically sound, could be affected by factors that are not part of the RoB assessment itself. Such factors in most cases may be out of control for the reviewers, as for example their experience with previous SRs. On other cases, factors such as being an author among the study under assessment, are amenable to intervention. A simple suggestion that also links this project with the previous one is to focus on the training of “less experienced” reviewers, possibly using a more practical approach which should encompass the critical issues we highlighted in our projects. The importance of formation, but also of constant guidance, is the common theme emerging from our studies on reporting. Our project on STROBE showed that authors of observational studies are generally aware of its existence and its importance, but it seems that this knowledge did not necessarily translate into action. Similarly, although we are confident that authors unanimously agree that “association is not causation”, we noticed very frequent slips into making inappropriate causal claims in observational studies.

The implications of our RoB projects mainly apply to researchers in the field of evidence synthesis and research methodologists. The projects on reporting, on the other hand, put editors of scientific journals at the centre of the stage. These two last projects made us realize the important role that editors may have, together with methodologists, in limiting the research waste. The project on STROBE showed that scientific journals are key factors in promoting the use of RGs. In our view this role should expand to become even more active. We realise that such suggestion could raise major feasibility issues. The improvements in the editorial approach that we suggested would be very time consuming and expensive for journals trying to adopt it. We are aware that many journals do not have the resources to verify the submission of the corresponding RG checklist with a manuscript. Similar issues will apply even more if editors must check the finer aspects of semantics relative to causal language in observational studies, given also the controversial nature of some aspects regarding wording. We think that this message would probably concern the journals with more resources. Journals with high impact factors seem to already publish studies with better overall quality of reporting[54]. This may be due to having more resources or to being more engaged in quality improvement, and to the ability to attract “better” research. Nevertheless, we are confident in suggesting that, with additional resources, journals could move towards demanding the use of RGs, while also engaging in a more proactive role, by offering tools to form and guide authors in the use of relevant RGs. We also think that strong editorial efforts should be made to avoid the use of unsupported and inappropriate causal claims in observational studies. Editors could focus on the appropriate dissemination of science, by supporting and hosting initiatives to

encourage the use of the right wording as well as fair discussions of the limits of different study designs.

7.3. Conclusions

We hope this thesis contributed to the existing literature on avoidable research waste and how to prevent it. We handled the issue of research waste from different directions, addressing different causes and different phases in the research process that each can generate waste, such as design deficiencies with a high risk of bias and incomplete or inadequate reporting. We were able to address these issues with multiple disciplines and various stakeholders in the research environment, including methodologists, researchers conducting systematic reviews and observational studies, and Cochrane and journal editors.

We identified several aspects of the RoB assessment process that can be improved, including external factors, while also indicating opportunities for more complete reporting of RCTs by authors.

Our project on reporting guidelines highlighted the role of scientific journals in improving completeness and fairness in reporting. Fairness in reporting also applies to the wording of causal claims in observational research. Here also, editors and journals can play a central role.

Highlighting shortcomings in the research process and suggesting interventions to overcome these may help in reducing avoidable waste in research, to the benefit of the research community and society as a whole.

Bibliography

1. Sackett, D.L., et al., *Evidence based medicine: what it is and what it isn't*. BMJ, 1996. **312**(7023): p. 71-2.
2. Eddy, D.M., *Clinical decision making: from theory to practice. Practice policies--guidelines for methods*. JAMA, 1990. **263**(13): p. 1839-41.
3. Bastian, H., P. Glasziou, and I. Chalmers, *Seventy-five trials and eleven systematic reviews a day: how will we ever keep up?* PLoS Med, 2010. **7**(9): p. e1000326.
4. Chalmers, I. and P. Glasziou, *Avoidable waste in the production and reporting of research evidence*. Lancet, 2009. **374**(9683): p. 86-9.
5. Macleod, M.R., et al., *Biomedical research: increasing value, reducing waste*. Lancet, 2014. **383**(9912): p. 101-4.
6. Ioannidis, J.P.A., et al., *Increasing value and reducing waste in research design, conduct, and analysis*. The Lancet, 2014. **383**(9912): p. 166-175.
7. Chalmers, I., et al., *How to increase value and reduce waste when research priorities are set*. The Lancet, 2014. **383**(9912): p. 156-165.
8. Chan, A.W., et al., *Increasing value and reducing waste: addressing inaccessible research*. Lancet, 2014. **383**(9913): p. 257-66.
9. Al-Shahi Salman, R., et al., *Increasing value and reducing waste in biomedical research regulation and management*. Lancet, 2014. **383**(9912): p. 176-85.
10. Hartling, L., et al., *Applying the risk of bias tool in a systematic review of combination long-acting beta-agonists and inhaled corticosteroids for persistent asthma*. PLoS One, 2011. **6**(2): p. e17242.
11. Hrobjartsson, A., et al., *Assessing risk of bias in randomised clinical trials included in Cochrane Reviews: the why is easy, the how is a challenge*. Cochrane Database Syst Rev, 2013(4): p. ED000058.
12. Hrobjartsson, A., et al., *Assessing Risk of Bias in Randomised Clinical Trials Included in Cochrane Reviews: The why is Easy, the how is a Challenge [editorial]*. Cochrane Database Syst Rev, 2013. **5**: p. ED000058.
13. Higgins JPT, Altman DG, Sterne JAC (editors). Chapter 8: Assessing risk of bias in included studies., in *Cochrane Handbook for Systematic Reviews of Interventions*, J.P. Higgins and S. Green, Editors., The Cochrane Collaboration, 2011. Available from www.handbook.cochrane.org.
14. Schulz, K.F., et al., *Empirical evidence of bias. Dimensions of methodological quality associated with estimates of treatment effects in controlled trials*. JAMA, 1995. **273**(5): p. 408-12.
15. Moher, D., et al., *Does quality of reports of randomised trials affect estimates of intervention efficacy reported in meta-analyses?* The Lancet, 1998. **352**(9128): p. 609-613.
16. Balk, E.M., et al., *Correlation of quality measures with estimates of treatment effect in meta-analyses of randomized controlled trials*. JAMA, 2002. **287**(22): p. 2973-82.
17. Page, M.J., et al., *Empirical Evidence of Study Design Biases in Randomized Trials: Systematic Review of Meta-Epidemiological Studies*. PLoS One, 2016. **11**(7): p. e0159267.
18. Dechartres, A., et al., *Empirical evaluation of which trial characteristics are associated with treatment effect estimates*. J Clin Epidemiol, 2016. **77**: p. 24-37.
19. Higgins, J.P.T., S. Green, and editors, *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]*. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org
20. Hartling, L., et al., *Testing the risk of bias tool showed low reliability between individual reviewers and across consensus assessments of reviewer pairs*. J Clin Epidemiol, 2013. **66**(9): p. 973-81.
21. Hartling, L., et al., *Risk of bias versus quality assessment of randomised controlled trials: cross sectional study*. BMJ, 2009. **339**(oct19 1): p. b4012.
22. Armijo-Olivo, S., et al., *Poor reliability between Cochrane reviewers and blinded external reviewers when applying the Cochrane risk of bias tool in physical therapy trials*. PLoS One, 2014. **9**(5): p. e96920.

23. Armijo-Olivo, S., et al., *Assessment of study quality for systematic reviews: a comparison of the Cochrane Collaboration Risk of Bias Tool and the Effective Public Health Practice Project Quality Assessment Tool: methodological research*. J Eval Clin Pract, 2012. **18**(1): p. 12-8.
24. Jordan, V.M., S.F. Lensen, and C.M. Farquhar, *There were large discrepancies in risk of bias tool judgments when a randomized controlled trial appeared in more than one systematic review*. J Clin Epidemiol, 2016.
25. da Costa, B.R., et al., *Effect of standardized training on the reliability of the Cochrane risk of bias assessment tool: a prospective study*. Syst Rev, 2017. **6**(1): p. 44.
26. Altman, D.G., D. Moher, and K.F. Schulz, *Improving the reporting of randomised trials: the CONSORT Statement and beyond*. Stat Med, 2012. **31**(25): p. 2985-97.
27. Begley, C.G. and J.P. Ioannidis, *Reproducibility in science: improving the standard for basic and preclinical research*. Circ Res, 2015. **116**(1): p. 116-26.
28. Glasziou, P., et al., *Reducing waste from incomplete or unusable reports of biomedical research*. The Lancet, 2014. **383**(9913): p. 267-276.
29. Mhaskar, R., et al., *Published methodological quality of randomized controlled trials does not reflect the actual quality assessed in protocols*. J Clin Epidemiol, 2012. **65**(6): p. 602-9.
30. Soares, H.P., et al., *Bad reporting does not mean bad methods for randomised trials: observational study of randomised controlled trials performed by the Radiation Therapy Oncology Group*. BMJ, 2004. **328**(7430): p. 22-4.
31. Altman, D.G., *Making research articles fit for purpose: structured reporting of key methods and findings*. Trials, 2015. **16**: p. 53.
32. Dechartres, A., et al., *Reviews assessing the quality or the reporting of randomized controlled trials are increasing over time but raised questions about how quality is assessed*. J Clin Epidemiol, 2011. **64**(2): p. 136-44.
33. Agha, R.A., et al., *Randomised controlled trials in plastic surgery: a systematic review of reporting quality*. Eur J Plast Surg, 2014. **37**: p. 55-62.
34. Bryant, J., et al., *A systematic review of the quality of reporting in published smoking cessation trials for pregnant women: an explanation for the evidence-practice gap?* Implement Sci, 2014. **9**: p. 94.
35. Chen, B., et al., *A retrospective survey of quality of reporting on randomized controlled trials of metformin for polycystic ovary syndrome*. Trials, 2014. **15**: p. 128.
36. Altman, D.G. and I. Simera, *A history of the evolution of guidelines for reporting medical research: the long road to the EQUATOR Network*. J R Soc Med, 2016. **109**(2): p. 67-77.
37. Chan, A.W., et al., *Empirical evidence for selective reporting of outcomes in randomized trials: comparison of protocols to published articles*. JAMA, 2004. **291**(20): p. 2457-65.
38. *A proposal for structured reporting of randomized controlled trials. The Standards of Reporting Trials Group*. JAMA, 1994. **272**(24): p. 1926-31.
39. Kendall, J.M., *Designing a research project: randomised controlled trials and their principles*. Emerg Med J, 2003. **20**(2): p. 164-8.
40. Begg, C., et al., *Improving the quality of reporting of randomized controlled trials. The CONSORT statement*. JAMA, 1996. **276**(8): p. 637-9.
41. *Call for comments on a proposal to improve reporting of clinical trials in the biomedical literature. Working Group on Recommendations for Reporting of Clinical Trials in the Biomedical Literature*. Ann Intern Med, 1994. **121**(11): p. 894-5.
42. Schulz, K.F., D.G. Altman, and D. Moher, *CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials*. PLoS Med, 2010. **7**(3): p. e1000251.
43. *Consort Extensions*. 20/11/2020]; Available from: <http://www.consort-statement.org/extensions>.
44. von Elm, E., et al., *The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies*. Epidemiology, 2007. **18**(6): p. 800-4.
45. Vandembroucke, J.P., et al., *Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): explanation and elaboration*. Epidemiology, 2007. **18**(6): p. 805-35.

46. da Costa, B.R., et al., *Uses and misuses of the STROBE statement: bibliographic study*. BMJ Open, 2011. **1**(1): p. e000048.
47. Cofield, S.S., R.V. Corona, and D.B. Allison, *Use of causal language in observational studies of obesity and nutrition*. Obes Facts, 2010. **3**(6): p. 353-6.
48. Thapa, D.K., et al., *Being honest with causal language in writing for publication*. J Adv Nurs, 2020. **76**(6): p. 1285-1288.
49. Kunz, R. and A.D. Oxman, *The unpredictability paradox: review of empirical comparisons of randomised and non-randomised clinical trials*. BMJ, 1998. **317**(7167): p. 1185-90.
50. Begg, M.D. and D. March, *Cause and Association: Missing the Forest for the Trees*. Am J Public Health, 2018. **108**(5): p. 620.
51. Hernan, M.A., *A definition of causal effect for epidemiological research*. J Epidemiol Community Health, 2004. **58**(4): p. 265-71.
52. Green, M.J., *Calculating Versus Estimating Causal Effects*. Am J Public Health, 2018. **108**(8): p. e4-e5.
53. Goetghebuer, E., et al., *Formulating causal questions and principled statistical answers*. Stat Med, 2020.
54. Dechartres, A., et al., *Evolution of poor reporting and inadequate methods over time in 20 920 randomised controlled trials included in Cochrane reviews: research on research study*. BMJ, 2017. **357**: p. j2490.
55. Sterne, J.A.C., et al., *RoB 2: a revised tool for assessing risk of bias in randomised trials*. BMJ, 2019. **366**: p. l4898.
56. *RoB 2.0 Tool*. 22/08/2018]; Available from: <https://sites.google.com/site/riskofbiastool/welcome/rob-2-0-tool>.
57. Sackett, D.L. and W.M. Rosenberg, *The need for evidence-based medicine*. J R Soc Med, 1995. **88**(11): p. 620-4.
58. Bordley, D.R., M. Fagan, and D. Theige, *Evidence-based medicine: a powerful educational tool for clerkship education*. Am J Med, 1997. **102**(5): p. 427-32.
59. Group, E.-B.M.W., *Evidence-based medicine. A new approach to teaching the practice of medicine*. JAMA, 1992. **268**(17): p. 2420-5.
60. Sackett, D.L., et al., *Evidence-Based Medicine: How to Practice and Teach EBM*. 2nd edition ed. 2000: Churchill Livingstone.
61. Ridsdale, L., *Evidence-Based Practice in Primary Care*. 1999.
62. *Evidence-based medicine, in its place*. Lancet, 1995. **346**(8978): p. 785.
63. Eddy, D.M., *Practice policies: where do they come from?* JAMA, 1990. **263**(9): p. 1265, 1269, 1272 passim.
64. WA, S., *Where is the evidence? Debates in modern medicine*. 1998, Oxford.
65. Haynes, B., et al., *Clinical Epidemiology: How to Do Clinical Practice Research*. 2006.
66. Atkins, D., et al., *Grading quality of evidence and strength of recommendations*. BMJ, 2004. **328**(7454): p. 1490.
67. Tugwell, P. and J.A. Kottnerus, *Is the 'Evidence-Pyramid' now dead?* J Clin Epidemiol, 2015. **68**(11): p. 1247-50.
68. Burns, P.B., R.J. Rohrich, and K.C. Chung, *The levels of evidence and their role in evidence-based medicine*. Plastic and reconstructive surgery, 2011. **128**(1): p. 305-310.
69. Reddy, K.R., *Evidence Based Medicine: A Paradigm for Clinical Practice*. Journal of Gandaki Medical College-Nepal, 2018. **11**(02): p. 74-81.
70. <https://pubmed.ncbi.nlm.nih.gov/?term=all%5Bsb%5D&sort=date>. 16/01/2021]; Available from: <https://pubmed.ncbi.nlm.nih.gov/?term=all%5Bsb>.
71. Hecht, L., S. Buhse, and G. Meyer, *Effectiveness of training in evidence-based medicine skills for healthcare professionals: a systematic review*. BMC Med Educ, 2016. **16**: p. 103.
72. Masic, I., M. Miokovic, and B. Muhamedagic, *Evidence based medicine - new approaches and challenges*. Acta Inform Med, 2008. **16**(4): p. 219-25.

73. Glasziou, P., J.P. Vandenbroucke, and I. Chalmers, *Assessing the quality of research*. BMJ, 2004. **328**(7430): p. 39-41.
74. Moher, D., et al., *Epidemiology and reporting characteristics of systematic reviews*. PLoS Med, 2007. **4**(3): p. e78.
75. Moher, D., et al., *Increasing value and reducing waste in biomedical research: who's listening?* The Lancet, 2016. **387**(10027): p. 1573-1586.
76. Clarke, M. and I. Chalmers, *Discussion sections in reports of controlled trials published in general medical journals: islands in search of continents?* JAMA, 1998. **280**(3): p. 280-2.
77. Clarke, M., S. Hopewell, and I. Chalmers, *Clinical trials should begin and end with systematic reviews of relevant evidence: 12 years and waiting*. Lancet, 2010. **376**(9734): p. 20-1.
78. Dickersin, K., *Health-care policy. To reform U.S. health care, start with systematic reviews*. Science, 2010. **329**(5991): p. 516-7.
79. Higgins, J.P., et al., *Cochrane Handbook for Systematic Reviews of Interventions*. 6.0 ed. 2019: The Cochrane Collaboration.
80. Kuehn, B.M., *Institute of Medicine outlines priorities for comparative effectiveness research*. JAMA, 2009. **302**(9): p. 936-7.
81. Luce, B.R., et al., *Rethinking randomized clinical trials for comparative effectiveness research: the need for transformational change*. Ann Intern Med, 2009. **151**(3): p. 206-9.
82. Sox, H.C. and S. Greenfield, *Comparative effectiveness research: a report from the Institute of Medicine*. Ann Intern Med, 2009. **151**(3): p. 203-5.
83. Sullivan, P. and D. Goldmann, *The promise of comparative effectiveness research*. JAMA, 2011. **305**(4): p. 400-1.
84. Straus, S., et al., *Evidence based medicine: How to practice and teach EBM*. 3rd ed. 2005.
85. Egger, M. and G.D. Smith, *Meta-Analysis. Potentials and promise*. BMJ, 1997. **315**(7119): p. 1371-4.
86. Egger, M., G.D. Smith, and A.N. Phillips, *Meta-analysis: principles and procedures*. BMJ, 1997. **315**(7121): p. 1533-7.
87. *Cochrane, About us*. 09/01/2021]; Available from: <https://www.cochrane.org/about-us>.
88. *Cochrane.org*. 09/01/2021]; Available from: <https://www.cochrane.org/>.
89. Pedersen, T., A.M. Moller, and J. Cracknell, *The mission of the cochrane anesthesia review group: preparing and disseminating systematic reviews of the effect of health care in anesthesiology*. Anesth Analg, 2002. **95**(4): p. 1012-8, table of contents.
90. Chalmers, I., *The Cochrane collaboration: preparing, maintaining, and disseminating systematic reviews of the effects of health care*. Ann N Y Acad Sci, 1993. **703**: p. 156-63; discussion 163-5.
91. *Review Manager (RevMan) [Computer program]*. 2014, The Nordic Cochrane Centre, The Cochrane Collaboration: Copenhagen.
92. *Cochrane Library*. 09/01/2021]; Available from: <https://www.cochranelibrary.com/>.
93. <https://www.cochrane.org/news/2019-journal-impact-factor-cochrane-database-systematic-reviews-7890>. Available from: <https://www.cochrane.org/news/2019-journal-impact-factor-cochrane-database-systematic-reviews-7890>.
94. Henderson, L.K., et al., *How to write a Cochrane systematic review*. Nephrology (Carlton), 2010. **15**(6): p. 617-24.
95. Begg, C.B. and J.A. Berlin, *Publication bias and dissemination of clinical research*. J Natl Cancer Inst, 1989. **81**(2): p. 107-15.
96. Berlin, J.A., C.B. Begg, and T.A. Louis, *An assessment of publication bias using a sample of published clinical trials*. Journal of the American Statistical Association, 1989. **84**(406): p. 381-392.
97. Dickersin, K., Y.I. Min, and C.L. Meinert, *Factors influencing publication of research results. Follow-up of applications submitted to two institutional review boards*. JAMA, 1992. **267**(3): p. 374-8.
98. Dickersin, K. and Y.I. Min, *Publication bias: the problem that won't go away*. Ann N Y Acad Sci, 1993. **703**: p. 135-46; discussion 146-8.
99. Hopewell, S., et al., *Publication bias in clinical trials due to statistical significance or direction of trial results*. Cochrane Database Syst Rev, 2009(1): p. MR000006.

100. Sterne, J.A., et al., *Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials*. BMJ, 2011. **343**: p. d4002.
101. Ioannidis, J.P., *Effect of the statistical significance of results on the time to completion and publication of randomized efficacy trials*. JAMA, 1998. **279**(4): p. 281-6.
102. Stern, J.M. and R.J. Simes, *Publication bias: evidence of delayed publication in a cohort study of clinical research projects*. BMJ, 1997. **315**(7109): p. 640-5.
103. Gotzsche, P.C., *Multiple publication of reports of drug trials*. Eur J Clin Pharmacol, 1989. **36**(5): p. 429-32.
104. Tramer, M.R., et al., *Impact of covert duplicate publication on meta-analysis: a case study*. BMJ, 1997. **315**(7109): p. 635-40.
105. Chan, A.W. and D.G. Altman, *Identifying outcome reporting bias in randomised trials on PubMed: review of publications and survey of authors*. BMJ, 2005. **330**(7494): p. 753.
106. Kirkham, J.J., et al., *The impact of outcome reporting bias in randomised controlled trials on a cohort of systematic reviews*. BMJ, 2010. **340**: p. c365.
107. Moher, D., et al., *Assessing the quality of randomized controlled trials: an annotated bibliography of scales and checklists*. Control Clin Trials, 1995. **16**(1): p. 62-73.
108. Juni, P., et al., *The hazards of scoring the quality of clinical trials for meta-analysis*. JAMA, 1999. **282**(11): p. 1054-60.
109. Viswanathan, M., et al., *Assessing the Risk of Bias of Individual Studies in Systematic Reviews of Health Care Interventions*, A.f.H.R.a.Q.M.G.f.C.E. Reviews., Editor. March 2012, AHRQ Publication.
110. Viswanathan, M., et al., *Assessing the Risk of Bias of Individual Studies in Systematic Reviews of Health Care Interventions*, in *Methods Guide for Effectiveness and Comparative Effectiveness Reviews*. 2008: Rockville MD. p. 198-.
111. Hartling, L., et al., *Developing and Testing a Tool for the Classification of Study Designs in Systematic Reviews of Interventions and Exposures*. 2010, Rockville MD.
112. Olivo, S.A., et al., *Scales to assess the quality of randomized controlled trials: a systematic review*. Phys Ther, 2008. **88**(2): p. 156-75.
113. Higgins, J.P., et al., *The Cochrane Collaboration's tool for assessing risk of bias in randomised trials*. BMJ, 2011. **343**: p. d5928.
114. Wood, L., et al., *Empirical evidence of bias in treatment effect estimates in controlled trials with different interventions and outcomes: meta-epidemiological study*. BMJ, 2008. **336**(7644): p. 601-5.
115. Sanderson, S., I.D. Tatt, and J.P. Higgins, *Tools for assessing quality and susceptibility to bias in observational studies in epidemiology: a systematic review and annotated bibliography*. Int J Epidemiol, 2007. **36**(3): p. 666-76.
116. Hopewell, S., et al., *Incorporation of assessments of risk of bias of primary studies in systematic reviews of randomised trials: a cross-sectional study*. BMJ Open, 2013. **3**(8): p. e003342.
117. Naylor, C.D., *Meta-analysis and the meta-epidemiology of clinical research*. BMJ, 1997. **315**(7109): p. 617-9.
118. Sterne, J.A., et al., *Statistical methods for assessing the influence of study characteristics on treatment effects in 'meta-epidemiological' research*. Stat Med, 2002. **21**(11): p. 1513-24.
119. Hill, A.B., *Suspended judgment. Memories of the British Streptomycin Trial in Tuberculosis. The first randomized clinical trial*. Control Clin Trials, 1990. **11**(2): p. 77-9.
120. Kjaergard, L.L., J. Villumsen, and C. Gluud, *Reported methodologic quality and discrepancies between large and small randomized trials in meta-analyses*. Ann Intern Med, 2001. **135**(11): p. 982-9.
121. Siersma, V., et al., *Multivariable modelling for meta-epidemiological assessment of the association between trial quality and treatment effects estimated in randomized clinical trials*. Stat Med, 2007. **26**(14): p. 2745-58.
122. Als-Nielsen, B., et al. *Are trial size and reported methodological quality associated with treatment effects? Observational study of 523 randomised trials [abstract]*. 12th Cochrane Colloquium: Bridging the Gaps; 2004 Oct 2-6; Ottawa, Ontario, Canada, 2004. 102-103.

123. Pildal, J., et al., *Impact of allocation concealment on conclusions drawn from meta-analyses of randomized trials*. Int J Epidemiol, 2007. **36**(4): p. 847-57.
124. Savovic, J., et al., *Influence of reported study design characteristics on intervention effect estimates from randomized, controlled trials*. Ann Intern Med, 2012. **157**(6): p. 429-38.
125. Hrobjartsson, A., et al., *Bias due to lack of patient blinding in clinical trials. A systematic review of trials randomizing patients to blind and nonblind sub-studies*. Int J Epidemiol, 2014. **43**(4): p. 1272-83.
126. Nuesch, E., et al., *The importance of allocation concealment and patient blinding in osteoarthritis trials: a meta-epidemiologic study*. Arthritis Rheum, 2009. **61**(12): p. 1633-41.
127. Juni, P., D.G. Altman, and M. Egger, *Systematic reviews in health care: Assessing the quality of controlled clinical trials*. BMJ, 2001. **323**(7303): p. 42-6.
128. Schulz, K.F., *Assessing allocation concealment and blinding in randomised controlled trials: why bother?* Evid Based Nurs, 2001. **4**(1): p. 4-6.
129. Schulz, K.F., I. Chalmers, and D.G. Altman, *The landscape and lexicon of blinding in randomized trials*. Ann Intern Med, 2002. **136**(3): p. 254-9.
130. Hollis, S. and F. Campbell, *What is meant by intention to treat analysis? Survey of published randomised controlled trials*. BMJ, 1999. **319**(7211): p. 670-4.
131. Melander, H., et al., *Evidence b(i)ased medicine--selective reporting from studies sponsored by pharmaceutical industry: review of studies in new drug applications*. BMJ, 2003. **326**(7400): p. 1171-3.
132. Porta, N., C. Bonet, and E. Cobo, *Discordance between reported intention-to-treat and per protocol analyses*. J Clin Epidemiol, 2007. **60**(7): p. 663-9.
133. Tierney, J.F. and L.A. Stewart, *Investigating patient exclusion bias in meta-analysis*. Int J Epidemiol, 2005. **34**(1): p. 79-87.
134. Schulz, K.F., et al., *Blinding and exclusions after allocation in randomised controlled trials: survey of published parallel group trials in obstetrics and gynaecology*. BMJ, 1996. **312**(7033): p. 742-4.
135. Moher, D., K.F. Schulz, and D. Altman, *The CONSORT statement: revised recommendations for improving the quality of reports of parallel-group randomized trials*. JAMA, 2001. **285**(15): p. 1987-91.
136. Chan, A.W., et al., *Outcome reporting bias in randomized trials funded by the Canadian Institutes of Health Research*. CMAJ, 2004. **171**(7): p. 735-40.
137. von Elm, E., et al., *Publication and non-publication of clinical trials: longitudinal study of applications submitted to a research ethics committee*. Swiss Med Wkly, 2008. **138**(13-14): p. 197-203.
138. Furukawa, T.A., et al., *Association between unreported outcomes and effect size estimates in Cochrane meta-analyses*. JAMA, 2007. **297**(5): p. 468-70.
139. Marshall, M., et al., *Unpublished rating scales: a major source of bias in randomised controlled trials of treatments for schizophrenia*. Br J Psychiatry, 2000. **176**: p. 249-52.
140. Liebetrau, A., *Measures of association*. 1983, Newbury Park, CA: Sage Publication.
141. Fleiss, J.L., B. Levin, and M.C. Paik, *Statistical methods for rates and proportions*. 3rd Edition ed. 2003, Hoboken, NJ: Wiley.
142. Byrt, T., *How good is that agreement?* Epidemiology, 1996. **7**(5): p. 561.
143. Hernan, M.A., *The C-Word: Scientific Euphemisms Do Not Improve Causal Inference From Observational Data*. Am J Public Health, 2018. **108**(5): p. 616-619.
144. Glymour, M.M. and R. Hamad, *Causal Thinking as a Critical Tool for Eliminating Social Inequalities in Health*. Am J Public Health, 2018. **108**(5): p. 623.
145. Lederer, D.J., et al., *Control of Confounding and Reporting of Results in Causal Inference Studies. Guidance for Authors from Editors of Respiratory, Sleep, and Critical Care Journals*. Ann Am Thorac Soc, 2019. **16**(1): p. 22-28.
146. Harhay, M.O., et al., *Methodologic Guidance and Expectations for the Development and Reporting of Prediction Models and Causal Inference Studies*. Ann Am Thorac Soc, 2020. **17**(6): p. 679-682.

147. Jorgensen, L., et al., *Evaluation of the Cochrane tool for assessing risk of bias in randomized clinical trials: overview of published comments and analysis of user practice in Cochrane and non-Cochrane reviews*. Syst Rev, 2016. **5**: p. 80.
148. Savovic, J., et al., *Evaluation of the Cochrane Collaboration's tool for assessing the risk of bias in randomized trials: focus groups, online survey, proposed recommendations and their implementation*. Syst Rev, 2014. **3**: p. 37.
149. Higgins, J.P., et al., *A revised tool for assessing risk of bias in randomized trials*. Cochrane Database of Systematic Reviews, 2016(10 (Suppl 1)).
150. Snyder, P.J., et al., *Effects of Testosterone Treatment in Older Men*. N Engl J Med, 2016. **374**(7): p. 611-24.
151. Walther, A., J. Breidenstein, and R. Miller, *Association of Testosterone Treatment With Alleviation of Depressive Symptoms in Men: A Systematic Review and Meta-analysis*. JAMA Psychiatry, 2019. **76**(1): p. 31-40.
152. Savovic, J., et al., *Development of a combined database for meta-epidemiological research*. Res Synth Methods, 2010. **1**(3-4): p. 212-25.
153. Higgins, J.P., et al., *The Cochrane Collaboration's tool for assessing risk of bias in randomised trials*. BMJ, 2011. **343**: p. d5928.
154. Alexander, G.C., et al., *Cardiovascular Risks of Exogenous Testosterone Use Among Men: A Systematic Review and Meta-Analysis*. Am J Med, 2017. **130**(3): p. 293-305.
155. Nian, Y., et al., *Testosterone replacement therapy improves health-related quality of life for patients with late-onset hypogonadism: a meta-analysis of randomized controlled trials*. Andrologia, 2017. **49**(4).
156. Kohn, T.P., et al., *Effects of Testosterone Replacement Therapy on Lower Urinary Tract Symptoms: A Systematic Review and Meta-analysis*. Eur Urol, 2016. **69**(6): p. 1083-90.
157. Corona, G., et al., *THERAPY OF ENDOCRINE DISEASE: Testosterone supplementation and body composition: results from a meta-analysis study*. Eur J Endocrinol, 2016. **174**(3): p. R99-116.
158. Neto, W.K., et al., *Effects of testosterone on lean mass gain in elderly men: systematic review with meta-analysis of controlled and randomized studies*. Age (Dordr), 2015. **37**(1): p. 9742.
159. Kang, D.Y. and H.J. Li, *The effect of testosterone replacement therapy on prostate-specific antigen (PSA) levels in men being treated for hypogonadism: a systematic review and meta-analysis*. Medicine (Baltimore), 2015. **94**(3): p. e410.
160. Grossmann, M., et al., *Effects of testosterone treatment on glucose metabolism and symptoms in men with type 2 diabetes and the metabolic syndrome: a systematic review and meta-analysis of randomized controlled clinical trials*. Clin Endocrinol (Oxf), 2015. **83**(3): p. 344-51.
161. Borst, S.E., et al., *Cardiovascular risks and elevation of serum DHT vary by route of testosterone administration: a systematic review and meta-analysis*. BMC Med, 2014. **12**: p. 211.
162. Corona, G., et al., *Testosterone supplementation and sexual function: a meta-analysis study*. J Sex Med, 2014. **11**(6): p. 1577-92.
163. Amanatkar, H.R., et al., *Impact of exogenous testosterone on mood: a systematic review and meta-analysis of randomized placebo-controlled trials*. Ann Clin Psychiatry, 2014. **26**(1): p. 19-32.
164. Cui, Y., et al., *The effect of testosterone replacement therapy on prostate cancer: a systematic review and meta-analysis*. Prostate Cancer Prostatic Dis, 2014. **17**(2): p. 132-43.
165. Xu, L., et al., *Testosterone therapy and cardiovascular events among men: a systematic review and meta-analysis of placebo-controlled randomized trials*. BMC Med, 2013. **11**: p. 108.
166. Cui, Y. and Y. Zhang, *The effect of androgen-replacement therapy on prostate growth: a systematic review and meta-analysis*. Eur Urol, 2013. **64**(5): p. 811-22.
167. Price, J. and G.C. Leng, *Steroid sex hormones for lower limb atherosclerosis*. Cochrane Database Syst Rev, 2012. **10**: p. CD000188.
168. Toma, M., et al., *Testosterone supplementation in heart failure: a meta-analysis*. Circ Heart Fail, 2012. **5**(3): p. 315-21.

169. Corona, G., et al., *Hypogonadism as a risk factor for cardiovascular mortality in men: a meta-analytic study*. Eur J Endocrinol, 2011. **165**(5): p. 687-701.
170. Corona, G., et al., *Type 2 diabetes mellitus and testosterone: a meta-analysis study*. Int J Androl, 2011. **34**(6 Pt 1): p. 528-40.
171. Fernandez-Balsells, M.M., et al., *Clinical review 1: Adverse effects of testosterone therapy in adult men: a systematic review and meta-analysis*. J Clin Endocrinol Metab, 2010. **95**(6): p. 2560-75.
172. Zarrouf, F.A., et al., *Testosterone and depression: systematic review and meta-analysis*. J Psychiatr Pract, 2009. **15**(4): p. 289-305.
173. Hrobjartsson, A., et al., *Observer bias in randomised clinical trials with binary outcomes: systematic review of trials with both blinded and non-blinded outcome assessors*. BMJ, 2012. **344**: p. e1119.