



UvA-DARE (Digital Academic Repository)

Plant turnover in response to climate change in the Cenozoic: Palynological insights from Myanmar, Southeast Asia and beyond

Huang, H.

Publication date

2021

[Link to publication](#)

Citation for published version (APA):

Huang, H. (2021). *Plant turnover in response to climate change in the Cenozoic: Palynological insights from Myanmar, Southeast Asia and beyond*. [Thesis, fully internal, Universiteit van Amsterdam].

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, P.O. Box 19185, 1000 GD Amsterdam, The Netherlands. You will be contacted as soon as possible.



REFERENCES

REFERENCES

- Abels, H.A., Dupont-Nivet, G., Xiao, G., Bosboom, R.E., Krijgsman, W., 2011. Step-wise change of Asian interior climate preceding the Eocene–Oligocene transition (EOT). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 299, 399–412.
- Adnet, S., Cappetta, H., Beard, K.C., Marivaux, L., Marandat, B., Chaimanee, Y., Jaeger, J.-J., Soe Thura Tun, Aung Naing Soe, 2008. First myliobatiform teeth (Elasmobranchii, Neoselachii) from the Pondaung formation (late middle Eocene) of central Myanmar. *Neues Jahrb. Geol. Palaontol. Abh.* 247, 335–340.
- Aiello-Lammens, M.E., Boria, R.A., Radosavljevic, A., Vilela, B., Anderson, R.P., 2015. spThin: An R package for spatial thinning of species occurrence records for use in ecological niche models. *Ecography* 38, 541–545.
- Aitchison, J.C., Ali, J., Davis, A.M., 2007. When and where did India and Asia collide? *J. Geophys. Res. Solid Earth* 112, B05423.
- Akkiraz, M.S., Akgün, F., Örcen, S., Bruch, A.A., Mosbrugger, V., 2006. Stratigraphic and palaeoenvironmental significance of Bartonian–Priabonian (middle–late Eocene) microfossils from the Başçeşme Formation, Denizli Province, Western Anatolia. *Turk. J. Earth Sci.* 15, 155–180.
- Akkiraz, M.S., Kayseri, M.S., Akgün, F., 2008. Palaeoecology of coal-bearing Eocene sediments in central Anatolia (Turkey) based on quantitative palynological data. *Turk. J. Earth Sci.* 17, 317–360.
- Aleksandrova, G.N., Kodrul, T.M., Jin, J.H., 2015. Palynological and paleobotanical investigations of Paleogene sections in the Maoming Basin, South China. *Stratigr. Geol. Correl.* 23, 300–325.
- Anderberg, A.A., Swenson, U., 2003. Evolutionary lineages in Sapotaceae (Ericales): a cladistic analysis based on ndhF sequence data. *Int. J. Plant Sci.* 164, 763–773.
- Anderson, J.A.R., 1964. The structure and development of the peat swamps of Sarawak and Brunei. *J. Trop. Geogr.* 18, 7–16.
- Anderson, J.A.R., 1976. Observations on the ecology of five peat swamp forests in Sumatra and Kalimantan. *Proceedings of Peat and Podzolic Soils and their Potential for Agriculture in Indonesia*, Soil Research Institute, Bogor, Indonesia, pp. 45–55.
- Anderson, J.A.R., Muller, J., 1975. Palynological study of a Holocene peat and a Miocene coal deposit from NW Borneo. *Rev. Palaeobot. Palynol.* 19, 291–351.

- Antonioli, L., Távora, V.A., Dino, R., 2015. Palynology of carinoliths and limestones from the Baunilha Grande Ecofacies of the Pirabas Formation (Miocene of Pará state, northeastern Brazil). *J. S. Am. Earth Sci.* 62, 134–147.
- Armentrout, J.M., 1991. Paleontologic constraints on depositional modelling: Examples of integration of biostratigraphy and seismic stratigraphy, Gulf of Mexico, in: Weimer, P., Link, M.H. (Eds.), *Seismic Facies and Sedimentary Processes of Submarine Fans and Turbidite Systems*. *Frontiers in Sedimentary Geology Series* Springer, New York, USA, pp. 137–170.
- Armentrout, J.M., 1996. High resolution sequence biostratigraphy: example from the Gulf of Mexico Plio-Pleistocene, in: Howell, J.A., Aitken, J.F. (Eds.), *High Resolution Sequence Stratigraphy: Innovations and Applications*. *Geological Society, London, Special Publications* 104, UK, pp. 65–86.
- Armstrong, K., Stone, G., Nicholls, J., Valderama, E., Anderberg, A., Smedmark, J., Gautier, L., Naciri, Y., Milne, R., Richardson, J., 2014. Patterns of diversification amongst tropical regions compared: a case study in Sapotaceae. *Front. Genet.* 5, 362.
- Asadu, A.N., Ofuyah, W.N., 2017. Miospore biozonation and age characterization of upper Miocene-Pliocene sediments in well X, deep offshore Niger delta. *J. Appl. Geol. Geophys.* 5, 6–13.
- Ashok, K., Guan, Z., Saji, N., Yamagata, T., 2010. Individual and combined influences of ENSO and the Indian Ocean dipole on the Indian summer monsoon. *J. Clim.* 17, 3141–3155.
- Ashton, P., 2014. *On the Forests of Tropical Asia: Lest the Memory Fade*. Kew Publishing, London, UK.
- Ashton, P., Zhu, H., 2020. The tropical-subtropical evergreen forest transition in East Asia: An exploration. *Plant Divers.* 42, 255–280.
- Askin, R.A., Baldoni, A.M., 1998. The Santonian through Paleogene record of Proteaceae in the southern South America-Antarctic Peninsula region. *Aust. Syst. Bot.* 11, 373–390.
- Atta-Peters, D., Salami, M.B., 2004. Late Cretaceous to Early Tertiary pollen grains from offshore Tano Basin, Southwestern Ghana. *Rev. Esp. Micropaleontol.* 36, 451–465.
- Atta-Peters, D., Salami, M.B., 2006. Aptian–Maastrichtian palynomorphs from the offshore Tano Basin, western Ghana. *J. Afr. Earth Sci.* 46, 379–394.

- Aung Naing Soe, Myitta, Soe Thura Tun, Aye Ko Aung, Tin Thein, Marandat, B., Ducrocq, S., Jaeger, J.-J., 2002. Sedimentary facies of the late Middle Eocene Pondaung Formation (central Myanmar) and the palaeoenvironments of its Anthropoid Primates. *C. R. Palevol* 1, 153–160.
- Bacon, C.D., Michonneau, F., Henderson, A.J., Mckenna, M.J., Milroy, A.M., Simmons, M.P., 2013. Geographic and taxonomic disparities in species diversity: dispersal and diversification rates across Wallace's line. *Evolution* 67, 2058–2071.
- Bacon, C.D., Velásquez-Puentes, F.J., Hoorn, C., Antonelli, A., 2018. Iriarteeae palms tracked the uplift of Andean Cordilleras. *J. Biogeogr.* 45, 1653–1663.
- Baker, W.J., 2015. A revised delimitation of the rattan genus *Calamus* (Arecaceae). *Phytotaxa* 197, 139–152.
- Baker, W.J., Coode, M.J.E., Dransfield, J., Dransfield, S., Harley, M.M., Johns, R.J., 1998. Patterns of distribution of Malesian vascular plants, in: Hall, R., Holloway, J.D. (Eds), *Biogeography and Geological Evolution of SE Asia*. Backhuys, Leiden, The Netherlands, pp. 243–258.
- Baker, W.J., Couvreur, T.L.P., 2013a. Global biogeography and diversification of palms sheds light on the evolution of tropical lineages. I. Historical biogeography. *J. Biogeogr.* 40, 274–285.
- Baker, W.J., Couvreur, T.L.P., 2013b. Global biogeography and diversification of palms sheds light on the evolution of tropical lineages. II. Diversification history and origin of regional assemblages. *J. Biogeogr.* 40, 286–298.
- Baker, W.J., Dransfield, J., 2000. Towards a biogeographic explanation of the calamoid palms, in: Wilson, K.L., Morrison, D.A. (Eds.), *Monocots: Systematics and Evolution*. CSIRO, Melbourne, Australia, pp. 545–553.
- Baker, W.J., Dransfield, J., 2016. Beyond Genera Palmarum: Progress and prospects in palm systematics. *Bot. J. Linn. Soc.* 182, 207–233.
- Baker, W.J., Norup, M.V., Clarkson, J.J., Couvreur, T.L.P., Dowe, J.L., Lewis, C.E., Pintaud, J.-C., Savolainen, V., Wilmot, T., Chase, M.W., 2011. Phylogenetic relationships among arecoid palms (Arecaceae: Arecoideae). *Ann. Bot.* 108, 1417–1432.
- Baker, W.J., Savolainen, V., Asmussen-Lange, C.B., Chase, M.W., Dransfield, J., Forest, F., Harley, M.M., Uhl, M.W., Wilkinson, M., 2009. Complete generic-level

REFERENCES

- phylogenetic analyses of palms (Arecaceae) with comparisons of supertree and supermatrix approaches. *Syst. Biol.* 58, 240–256.
- Baksi, S.K., 1962. Palynological investigation of Simsang River Tertiaries, South Shillong Front, Assam. *Bull. Geol. Min. Metall. Soc. India* 26, 1–22.
- Baksi, S.K., 1974. Significant pollen taxa in the stratigraphical analysis of the Tertiary sediments of Assam, in: Surange, K.R. (Ed.), *Aspects and Appraisal of Indian Palaeobotany*. Birbal Sahni Institute of Palaeobotany, Lucknow, India, pp. 502–515.
- Baksi, S.K., Venkatachala, B.S., 1970. *Meyeripollis*, a new genus from the Tertiary sediments of Assam. *J. Geol. Soc. India* 11, 81–83.
- Banin, L., Feldpausch, R., Phillips, O.L., Baker, T.R., Lloyd, J., Affum-Baffoe, K., Arets, E.J.M.M., Berry, N.J., Bradford, M., Brienen, R.J.W., 2012. What controls tropical forest architecture? Testing environmental, structural and floristic drivers. *Glob. Ecol. Biogeogr.* 21, 1179–1190.
- Bao, T., Wang, B., Li, J., Dilcher, D., 2019. Pollination of Cretaceous flowers. *Proc. Nat. Acad. Sci. U.S.A.* 116, 24707–24711.
- Barreda, V.D., Zamaloa, M.C., Gandolfo, M.A., Jaramillo, C., Wilf, P., 2020. Early Eocene spore and pollen assemblages from the Laguna del Hunco fossil lake beds, Patagonia, Argentina. *Int. J. Plant Sci.* 181, 594–615.
- Barrett, C.F., Bacon, C.D., Antonelli, A., Cano, A., Hofmann, T., 2016. An introduction to plant phylogenomics with a focus on palms. *Bot. J. Linn. Soc.* 182, 234–255.
- Bartish, I.V., Antonelli, A., Richardson, J.E., Swenson, U., 2011. Vicariance or long-distance dispersal: historical biogeography of the pantropical subfamily Chrysophylloideae (Sapotaceae). *J. Biogeogr.* 38, 177–190.
- Beck, H.E., Zimmermann, N.E., McVicar, T.R., Vergopolan, N., Berg, A., Wood, E.F., 2018. Present and future Köppen-Geiger climate classification maps at 1-km resolution. *Sci. Data* 5, 180214.
- Behling, H., Berrio, J.C., Hooghiemstra, H., 1999. Late Quaternary pollen records from the middle Caquetá river basin in central Colombian Amazon. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 145, 193–213.
- Beal, E.I., 1998. Stratigraphic and palaeoenvironmental significance of Eocene palynomorphs from the Rusayl Shale Formation, Al Khawd, northern Oman. *Rev. Palaeobot. Palynol.* 102, 249–258.

- Belsky, C.Y., Boltenhagen, E., Potonié, R., 1965. Sporae dispersae der Oberen Kreide von Gabun, Äquatoriales Afrika. *PalZ* 39, 72.
- Bender, F., 1983. *Geology of Burma*. Gebr. Borntraeger, Berlin, Germany.
- Bercovici, A., Hadley, A., Villanueva-Amadoz, U., 2009. Improving depth of field resolution for palynological photomicrography. *Palaeontol. Electron.* 12, 1–12.
- Berger, B.A., Kriebel, R., Spalink, D., Sytsma, K.J., 2016. Divergence times, historical biogeography, and shifts in speciation rates of Myrtales. *Mol. Phylogenet. Evol.* 95, 116–136.
- Berrio, J.C., Hooghiemstra, H., Behling, H., Botero, P., van der Borg, K., 2002. Late Quaternary savanna history of the Colombian Llanos Orientales from Laguna Chenevo and Mozambique: a transect synthesis. *The Holocene* 12, 35–48.
- Berry, E.W., 1929. Tertiary fossil plants from South America. *Proc. U.S. Natl. Mus.* 75, 1–12.
- Bhatia, H., Khan, M.A., Srivastava, G., Hazra, T., Spicer, R.A., Hazra, M., Mehrotra, R.C., Spicer, T.E.V., Bera, S., Roy, K., 2021. Late Cretaceous–Paleogene Indian Monsoon climate vis-à-vis movement of the Indian plate, and the birth of the South Asian Monsoon. *Gondwana Res.* 93, 89–100.
- Bié, G.R., Digbehi, Z.B., Yao, K.R., Tea-Yassi, J., Kangah, K.D., Tahi, I., 2012. Stratigraphie Palynologique du Maastrichtien Supérieur-Eocène Supérieur du Bassin Sédimentaire Offshore de Côte d'Ivoire, Afrique de l'Ouest. *Int. J. Afr. Stud.* 6, 40–57.
- Bignot, G., Blondeau, A., Guernet, C., Perreau, M., Poignant, A., Renard, M., Riveline, J., 1985. Age and characteristics of the Eocene transgression at Gant (Vertes Mountains, Transdanubia, Hungary). *Acta Geol. Hung.* 28, 29–48.
- Birks, H.J.B., Line, J.M., 1992. The use of rarefaction analysis for estimating palynological richness from Quaternary pollen-analytical data. *The Holocene* 2, 1–10.
- Bogotá-Ángel, G., Huang, H., Jardine, P.E., Chazot, N., Salamanca, S., Banks, H., Pardo-Trujillo, A., Plata, A., Dueñas, H., Star, W., Langelaan, R., Eisawi, A., Umeji, O.P., Enuenwemba, L.O., Parmar, S., da Silveira, R.R., Lim, J.Y., Prasad, V., Morley, R.J., Bacon, C.D., Hoorn, C., 2021. Climate and geological change as drivers of Mauritiinae palm biogeography. *J. Biogeogr.* <https://doi.org/10.1111/jbi.14098>
- Bohaty, S.M., Zachos, J.C., 2003. Significant Southern Ocean warming event in the late middle Eocene. *Geology* 31, 1017–1020.

REFERENCES

- Bolaji, T.A., Ndukwe, O.S., Oyebamiji, A.R., Ikegwuonu, O.N., 2020. Palynological age control and paleoenvironments of the Paleogene strata in Eastern Dahomey Basin, Southwestern Nigeria. *Sci. Rep.* 10, 8991.
- Boltenhagen, E., 1967. Spores et pollen du Crétacé supérieur du Gabon. *Pollen et Spores* 9, 335–355.
- Boltenhagen, E., 1976. Senonian microflora of Gabon. *Rev. Micropaléontol.* 18, 191–199.
- Bóna, J., Nagy, E., 1981. Nanoplankton de las secuencias terrígenas del Cretácico Superior de la región oriental de Cuba. *Ciencias de la Tierra y el Espacio* 3, 31–35.
- Bondarenko, O.V., Blokhina, N.I., Bruch, A.A., Henrot, A.-J., Utescher, T., 2017. Quantification of Calabrian vegetation in southern Primorye (Far East of Russia) using multiple proxies. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 467, 253–264.
- Bonnefille, R., 1983. Evidence for a cooler and drier climate in the Ethiopian uplands, 2.5 Ma ago. *Nature* 303, 487–491.
- Boos, W., Kuang, Z., 2010. Dominant control of the South Asian monsoon by orographic insulation versus plateau heating. *Nature* 463, 218–222.
- Bosboom, R.E., Abels, H.A., Hoorn, C., van den Berg, B.C.J., Guo, Z., Dupont-Nivet, G., 2014a. Aridification in continental Asia after the Middle Eocene Climatic Optimum (MECO). *Earth Planet. Sci. Lett.* 389, 34–42.
- Bosboom, R., Dupont-Nivet, G., Grothe, A., Brinkhuis, H., Villa, G., Mandic, O., Stoica, M., Kouwenhoven, T., Huang, W., Yang, W., Guo, Z., 2014b. Timing, cause and impact of the late Eocene stepwise sea retreat from the Tarim Basin (west China). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 403, 101–118.
- Bosboom, R.E., Dupont-Nivet, G., Houben, A.J.P., Brinkhuis, H., Villa, G., Mandic, O., Stoica, M., Zachariasse, W.-J., Guo, Z., Li, C., 2011. Late Eocene sea retreat from the Tarim Basin (West China) and concomitant Asian paleoenvironmental change. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 299, 385–398.
- Braman, D.R., Sweet, A.R., 1999. Terrestrial palynomorph biostratigraphy of the Cypress Hills, Wood Formation, and Turtle Mountain areas (Upper Cretaceous–Paleocene) of western Canada. *Can. J. Earth Sci.* 36, 725–741.
- Brummitt, R.K., Pando, F., Hollis, S., Brummitt, N.A., 2001. World Geographical Scheme for Recording Plant Distributions. International Working Group on Taxonomic Databases for Plant Sciences (TDWG). Royal Botanic Gardens, Kew, Richmond, UK.

- Brunig, E.F., 1974. Ecological Studies in Kerangas Forests of Sarawak and Brunei. Borneo Literature Bureau for Sarawak Forest Department, Sarawak, Malaysia.
- Caballero, R., Huber, M., 2013. State-dependent climate sensitivity in past warm climates and its implications for future climate projections. *Proc. Natl. Acad. Sci. U.S.A.* 110, 14162–14167.
- Cailliez, F., 1983. The analytical solution of the additive constant problem. *Psychometrika* 48, 305–308.
- Carpenter, R.J., Macphail, M.K., Jordan, G.J., Hill, R.S., 2015. Fossil evidence for open, Proteaceae-dominated heathlands and fire in the Late Cretaceous of Australia. *Am. J. Bot.* 12, 2092–2107.
- Cássia-Silva, C., Freitas, C.G., Alves, D.M., Bacon, C.D., Collevatti, R.G., 2019. Niche conservatism drives a global discrepancy in palm species richness between seasonally dry and moist habitats. *Glob. Ecol. Biogeogr.* 28, 814–825.
- Chakraborty, M., 2004. Palynology of Lakadong sandstone exposed around Bhalukurung, North Cachher hills, Assam. *Palaeobotanist* 53, 113–121.
- Chandra, A., Kumar, M., 1997. Palynology of the Late Tertiary sediments (DSDP Site 218) in the Bengal Fan, Indian Ocean. *Palaeobotanist* 46, 51–69.
- Chao, A., Jost, L., 2012. Coverage-based rarefaction and extrapolation: standardizing samples by completeness rather than size. *Ecology* 93, 2533–2547.
- Chatterjee, S., Scotese, C.R., 1999. The breakup of Gondwana and the evolution and biogeography of the Indian plate. *Proc. Indian Natl. Sci. Acad.* 65, 397–425.
- Chavasseau, O., Chaimanee, Y., Coster, P., Emonet, E.-G., Soe Aung Naing, Aung Aung Kyaw, Aye Maung, Rugbumrung, M., Hla Shwe, Jaeger, J.-J., 2010. First record of a chalicothere from the Miocene of Myanmar. *Acta Palaeontol. Pol.* 55, 13–22.
- Cheng, X, Fan, L., Gu, W., 2019. *Comprehensive Practice of Exploration and Evaluation Techniques in Complex Reservoirs*. Petroleum Industry Press and Springer Nature Singapore Pte Ltd., Springer, Singapore.
- Cheng, Z., Weng, C., Foong, S., Dai, L., 2020. A study on modern pollen rain and pollen morphology in the tropical western Malay Peninsula and its implications for paleoenvironmental reconstructions in the Sunda region. *Rev. Palaeobot. Palynol.* 279, 104236.

REFERENCES

- Chiadikobi, K.C., Chiaghanam, O.I., Onyemesili, O.C., Omoboriwo. A.O., 2018. Palynological study of the Campano-Maastrichtian Nkporo Group of Anambra Basin, southeastern, Nigeria. *World News Nat. Sci.* 20, 31–53.
- Chmura, C.A., 1973. Upper Cretaceous (Campanian–Maastrichtian) angiosperm pollen from the western San Joaquin Valley, California, USA. *Palaeontogr. Abt. B* 141, 89–171.
- Christenhusz, M.J.M., Byng, J.W., 2016. The number of known plants species in the world and its annual increase. *Phytotaxa* 261, 201–217.
- Coetzee, J.A., 1978. Climate and biological changes in southwestern Africa during the Late Cainozoic. *Palaeocol. Afr.* 10, 13–29.
- Cohen, K.M., Harper, D.A.T., Gibbard, P.L., 2020. ICS International Chronostratigraphic Chart 2020/03. International Commission on Stratigraphy, IUGS. www.stratigraphy.org. (accessed 23 December 2020)
- Collinson, M.E., 1993. Taphonomy and fruiting ecology of recent and fossil *Nypa*. *Spec. Pap. Palaeontol.* 49, 165–180.
- Colmenares, O.A., Teran, L., 1993. A biostratigraphic study of Paleogene sequences in southwestern Venezuela. *Palynology* 17, 67–89.
- Colwell, R.K., Chao, A., Gotelli, N.J., Lin, S.Y., Mao, C.X., Chazdon, R.L., Longino, J.T., 2012. Models and estimators linking individual-based and sample-based rarefaction, extrapolation, and comparison of assemblages. *J. Plant Ecol.* 5, 3–21.
- Condamine, F.L., Rolland, J., Morlon, H., 2013. Macroevolutionary perspectives to environmental change. *Ecol. Lett.* 16, 72–85.
- Cookson, I.C., 1950. Fossil pollen grains of Proteaceous type from Tertiary deposits in Australia. *Aust. J. Biol. Sci.* 3, 166–177.
- Cookson, I.C., Pike, K.M., 1954. Some dicotyledonous pollen types from Cainozoic deposits in the Australian region. *Aust. J. Bot.* 2, 197–219.
- Correa, E., Jaramillo, C., Manchester, S., Gutierrez, M., 2010. A fruit and leaves of Rhamnaceous affinities from the late Cretaceous (Maastrichtian) of Colombia. *Am. J. Bot.* 97, 71–79.
- Couvreur, T.L.P., Baker, W.J., 2013. Tropical rain forest evolution: palms as a model group. *BMC Biol.* 11, 48.
- Couvreur, T.L.P., Dauby, G., Blach-Overgaard, A., Deblauwe, V., Dessein, S., Droissart, V., Hardy, O.J., Harris, D.J., Janssens, S.B., Ley, A.C., Mackinder, B.A., Sonké, B.,

- Sosef, M.S.M., Stévant, T., Svenning, J.-C., Wieringa, J.J., Faye, A., Missoup, A.D., Tolley, K.A., Nicolas, V., Ntie, S., Fluteau, F., Robin, C., Guillocheau, F., Barboni, D., Sepulchre, P., 2021. Tectonics, climate and the diversification of the tropical African flora and fauna. *Biol. Rev.* 96, 16–51.
- Couvreur, T.L.P., Forest, F., Baker, W.J., 2011. Origin and global diversification patterns of tropical rain forests: inferences from a complete genus-level phylogeny of palms. *BMC Biol.* 9, 44.
- Couvreur, T.L.P., Kissling, W.D., Condamine, F.L., Svenning, J.-C., Rowe, N.P., Baker, W.J., 2015. Global diversification of a tropical plant growth form: environmental correlates and historical contingencies in climbing palms. *Front. Genet.* 5, 452.
- Coxall, H.K., Pearson, P.N., 2007. The Eocene–Oligocene transition, in: Williams, M., Haywood, A.M., Gregory, F.J., Schmidt, D.N. (Eds.), *Deep-time Perspectives on Climate Change: Marrying the Signal from Computer Models and Biological Proxies*. The Geological Society, London, UK, pp. 351–387.
- Crane, P.R., Lidgard, S., 1989. Angiosperm diversification and paleolatitudinal gradients in Cretaceous floristic diversity. *Science* 246, 675–678.
- Crié, L., 1892. Recherches sur les palmiers silicifiés des terrains Crétacés de l'Anjou. *Bull. Soc. Études Sci. Angers* 21, 97–103.
- Crisp, M.D., Cook, L.G., 2011. Cenozoic extinctions account for low diversity of extant gymnosperms compared with angiosperms. *New Phytol.* 192, 997–1009.
- Cruikshank, R.D., Ko, K., 2003. Geology of an amber locality in the Hukawng Valley, Northern Myanmar. *J. Asian Earth Sci.* 21, 441–455.
- D'Apolito, C., Absy, M.L., Latrubesse, E.M., 2013. The Hill of Six Lakes revisited: new data and re-evaluation of a key Pleistocene Amazon site. *Quat. Sci. Rev.* 76, 140–155.
- D'Apolito, C., da Silva-Caminha, S.A.F., Jaramillo, C., Dino, R., Soares, E.A.A., 2019. The Pliocene–Pleistocene palynology of the Negro River, Brazil. *Palynology* 43, 223–243.
- Davies, E.H., 2001. Palynological analysis and age assignments of two Burmese amber sample sets. Unpublished report by Branta Biostratigraphy Ltd. for Leeward Capital Corp.
- De Bonis, L., Solé, F., Chaimanee, Y., Aung Naing Soe, Chit Sein, Lazzari, V., Chavasseau, O., Jaeger, J.-J., 2018. New hyaenodonts (Mammalia) from the middle Eocene of Myanmar. *C. R. Palevol* 17, 357–365.

- de la Parra, F., 2009. Palynological changes across the Cretaceous-Tertiary boundary in Colombia, South America. MSc Thesis, University of Florida, Gainesville, USA.
- de Terra, H., 1944. Component geographic factors of the natural regions of Burma. *Ann. Am. Assoc. Geogr.* 34, 67–96.
- Deconto, R.M., Pollard, D., 2003. Rapid Cenozoic glaciation of Antarctica induced by declining atmospheric CO₂. *Nature* 421, 245–249.
- Delcourt, H.R., Delcourt, P.A., Webb III, T., 1982. Dynamic plant ecology: The spectrum of vegetational change in space and time. *Quat. Sci. Rev.* 1, 153–175.
- Descombes, P., Gaboriau, T., Albouy, C., Heine, C., Leprieur, F., Pellissier, L., 2018. Linking species diversification to palaeo-environmental changes: A process-based modelling approach. *Glob. Ecol. Biogeogr.* 27, 233–244.
- Dettmann, M.E., Jarzen, D.M., 1998. The early history of the Proteaceae in Australia: the pollen record. *Aust. Syst. Bot.* 11, 401–438.
- Digbehi, Z.B., Guédé, K.E., Yao, N., Affian, K., Toé, B.K.K., Yao, K.R., Tahi, I., 2011. Palynostratigraphy and depositional palaeoenvironment of Cretaceous–Palaeogene (K–Pg) Boundary deposits of Abidjan margin (Cote d'Ivoire). *J. Geogr. Reg. Plann.* 4, 644–655.
- Digbehi, Z.B., Yao, K.R., Tea, Y.J., Boblai, G., 1996. Contribution à l'étude palynologique et paléoenvironnementale du Campanien et du Maastrichtien du Bassin offshore de Côte d'Ivoire. *Géol. Méditerran.* 23, 155–171.
- Dilcher, D.L., Manchester, S.R., 1986. Investigations of angiosperms from the Eocene of North America: Leaves of the Engelhardiaceae (Juglandaceae). *Bot. Gaz.* 147, 189–199.
- Ding, L., Spicer, R., Yang, J., Xu, Q., Cai, F., Li, S., Lai, Q., Wang, H., Spicer, T.E.V., Yue, Y., Shukla, A., Srivastava, G., Khan, M.A., Bera, S., Mehrotra, R., 2017. Quantifying the rise of the Himalaya orogen and implications for the South Asian monsoon. *Geology* 45, 215–218.
- Donoghue, M.J., Edwards, E.J., 2014. Biome shifts and niche evolution in plants. *Annu. Rev. Ecol. Evol. Syst.* 45, 547–572.
- Doubinger, J., 1973. Pollen and spores from the Paleocene coal basin of Cerrejón (Guajira Province, Colombia). *Proceedings de la 1996 Congres de National des Sociétés Savantes, Toulouse, 1971*, 5, 253–262.
- Dransfield, J., 1974. *A Short Guide to Rattans*. Regional Center for Tropical Biology (BIOTROP), Bogor, Indonesia.

- Dransfield, J., Uhl, N.W., Asmussen, C.B., Baker, W.J., Harley, M.M., Lewis, C.E., 2008. *Genera Palmarum: The Evolution and Classification of Palms*. Royal Botanic Gardens, Kew, Richmond, UK.
- Dueñas, H., 1980. Palynology of Oligocene–Miocene strata of borehole Q-E-22, Planeta Rica, Northern Colombia. *Rev. Palaeobot. Palynol.* 30, 313–328.
- Duke, N.O., 1991. *Nypa* in the mangroves of Central America: introduced or relict? *Principes* 35, 127–132.
- Dupont-Nivet, G., Hoorn, C., Konert, M., 2008. Tibetan uplift prior to the Eocene–Oligocene climate transition: Evidence from pollen analysis of the Xining Basin. *Geology* 36, 987–990.
- Dupont-Nivet, G., Krijgsman, W., Langereis, C.G., Abels, H.A., Dai, S., Fang, X., 2007. Tibetan plateau aridification linked to global cooling at the Eocene–Oligocene transition. *Nature* 445, 635–638.
- Dutta, S., Mathews, R.P., Singh, B.D., Tripathi, S.M., Singh, A., Saraswati, P.K., Banerjee, S., Mann, U., 2011. Petrology, palynology and organic geochemistry of Eocene lignite of Matanomadh, Kutch Basin, western India: Implications to depositional environment and hydrocarbon source potential. *Int. J. Coal Geol.* 85, 91–102.
- Ediger, V.Ş., Bati, Z., Alişan, C., 1990. Paleopalynology and paleoecology of *Calamus*-like disulcate pollen grains. *Rev. Palaeobot. Palynol.* 62, 97–105.
- Eisawi, A., Schrank, E., 2008. Upper Cretaceous to Neogene palynology of the Melut Basin, Southeast Sudan. *Palynology* 32, 101–129.
- Eisawi, A., Schrank, E., 2009. Terrestrial palynology and age assessment of the Gedaref Formation (eastern Sudan). *J. Afr. Earth Sci.* 54, 22–30.
- Eiserhardt, W.L., Couvreur, T.L.P., Baker, W.J., 2017. Plant phylogeny as a window on the evolution of hyperdiversity in the tropical rainforest biome. *New Phytol.* 214, 1408–1422.
- Eiserhardt, W.L., Svenning, J.-C., Kissling, W.D., Balslev, H., 2011. Geographical ecology of the palms (Arecaceae): Determinants of diversity and distributions across spatial scales. *Ann. Bot.* 108, 1391–1416.
- Eldrett, J.S., Greenwood, D.R., Harding, I.C., Huber, M., 2009. Increased seasonality through the Eocene to Oligocene transition in northern high latitudes. *Nature* 459, 969–973.

REFERENCES

- Elith, J., Phillips, S.J., Hastie, T., Dudík, M., Chee, Y.E., Yates, C.J., 2011. A statistical explanation of MaxEnt for ecologists. *Divers. Distrib.* 17, 43–57.
- Ellison, A.M., Farnsworth, E.J., Merkt, R.E., 1999. Origins of mangrove ecosystems and the mangrove biodiversity anomaly. *Glob. Ecol. Biogeogr.* 8, 95–115.
- Engelhardt, D.W., Wrenn, J.H., 1994. Palynology and paleoenvironments of Eocene sediments, Saw area, Salin Basin, Myanmar (Burma). *Palynology* 18, 243–244.
- Environment Systems Research Institute (ESRI), 2019. Guidebooks of ArcMap 10.7. Website. <http://desktop.arcgis.com/en/arcmap/latest/map/projections/mollweide.htm>
- Erdtman, G., 1952. *Pollen Morphology and Plant Taxonomy: Angiosperms*. Almqvist & Wiksell, Stockholm, Sweden.
- Escobar, L.E., 1982. Estudio palinológico de la Formación Amaga. *Boletín de Ciencias de la Tierra (Universidad Nacional de Colombia, Bogotá)* 7–8, 117–129.
- Falcon-Lang, H., Mages, V., Collinson, M., 2016. The oldest *Pinus* and its preservation by fire. *Geology* 44, 303–306.
- Faye, A., Pintaud, J.C., Baker, W.J., Vigouroux, Y., Sonke, B., Couvreur, T.L.P., 2016. Phylogenetics and diversification history of African rattans (Calamoideae, Ancistrophyllinae). *Bot. J. Linn. Soc.* 182, 256–271.
- Feng, X., Tang, B., Kodrul, T.M., Jin, J., 2013. Winged fruits and associated leaves of *Shorea* (Dipterocarpaceae) from the Late Eocene of South China and their phytogeographic and paleoclimatic implications. *Am. J. Bot.* 100, 574–581.
- Ferguson, D.K., Zetter, R., Paudyal, K.N., 2007. The need for the SEM in palaeopalynology. *C. R. Palevol* 6, 423–430.
- Ferguson, I.K., Harley, M.M., 1993. The significance of new and recent work on pollen morphology of the Palmae. *Kew Bull.* 48, 205–243.
- Fick, S.E., Hijmans, R.J., 2017. WorldClim 2: New 1-km spatial resolution climate surfaces for global land areas. *Int. J. Climatol.* 37, 4302–4315.
- Filatoff, J., Hughes, W., 1996. Late Cretaceous to Recent palaeoenvironments of the Saudi Arabian Red Sea. *J. Afr. Earth Sci.* 22, 535–548.
- Fine, P.V.A., Ree, R.H., 2006. Evidence for a time-integrated species-area effect on the latitudinal gradient in tree diversity. *Am. Nat.* 168, 796–804.
- Fo, A., Fa, L., 2018. Miospore Biozonation of Cenomanian–Santonian Succession in FAMO-1 well, Gongola Sub Basin, Upper Benue Trough, Nigeria. *J. Appl. Sci. Environ. Manag.* 22, 1171–1176.

- Foster, C.B., 1982. Illustrations of Early Tertiary (Eocene) plant microfossils from the Yaamba Basin, Queensland. Queensland Department of Mines, Brisbane, Australia.
- Foster, G.L., Royer, D.L., Lunt, D.J., 2017. Future climate forcing potentially without precedent in the last 420 million years. *Nat. Commun.* 8, 14845.
- François, L., Bruch, A.A., Utescher, T., Spicer, R.A., Spicer, T., 2017. Reconstructing Cenozoic vegetation from proxy data and models – a NECLIME synthesis (Editorial). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 467, 1–4.
- Frederiksen, N.O., 1980. Sporomorphs from the Jackson Group (Upper Eocene) and adjacent strata of Mississippi and western Alabama. U.S. Geol. Surv. Prof. Paper 1084, 1–75.
- Frederiksen, N.O., 1984. Stratigraphic, paleoclimatic, and biogeographic significance of Tertiary sporomorphs from Massachusetts. U.S. Geol. Surv. Prof. Paper 1308, 25 pp.
- Frederiksen, N.O., 1994. Middle and late Palaeocene angiosperm pollen from Pakistan. *Palynology* 18, 91–137.
- Friis, E.M., Pedersen, K.R., Crane, P.R., 2004. Araceae from the Early Cretaceous of Portugal: evidence on the emergence of monocotyledons. *Proc. Natl. Acad. Sci. U.S.A.* 101, 16565–16570.
- Galeano, G., Bernal, R., 2010. *Palmas de Colombia: Guía de campo*. Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Bogotá D.C., Colombia.
- Garzon, S., Warny, S., Bart, P.J., 2012. A palynological and sequence-stratigraphic study of Santonian–Maastrichtian strata from the Upper Magdalena Valley basin in central Colombia. *Palynology* 36, 112–133.
- GBIF.org. GBIF Occurrence Download (<https://doi.org/10.15468/dl.ag40lq>). (accessed on 5 August 2019)
- GBIF.org. GBIF Occurrence Download (<https://doi.org/10.15468/dl.bzjzjl>). (accessed on 5 August 2019)
- GBIF.org. GBIF Occurrence Download (<https://doi.org/10.15468/dl.rliz8n>). (accessed on 5 August 2019)
- GBIF.org. GBIF Occurrence Download (<https://doi.org/10.15468/dl.wvfy3m>). (accessed on 5 August 2019)
- GBIF.org. GBIF Occurrence Download (<https://doi.org/10.15468/dl.prgnvt>). (accessed on 16 June 2020)

REFERENCES

- Gee, C.T., 1990. On the fossil occurrences of the mangrove palm *Nypa*, in: Knobloch, E., Kvaček, Z. (Eds.), *Paleofloristic and Paleoclimatic Changes in the Cretaceous and Tertiary*. Prague, Czech Republic, pp. 315–319.
- Gee, C.T., 2001. The mangrove palm *Nypa* in the geologic past of the New World. *Wetl. Ecol. Manag.* 9, 181–194.
- Germeraad, J.H., Hopping, C.A., Muller, J., 1968. Palynology of Tertiary sediments from tropical areas. *Rev. Palaeobot. Palynol.* 6, 189–348.
- González-Guzmán, A.E., 1967. A palynological study on the upper Los Cuervos and Mirador formations (Lower and Middle Eocene; Tibú area, Colombia). Ph.D. Thesis, University of Amsterdam, Amsterdam, The Netherlands.
- Gotelli, N., Colwell, R.K., 2001. Quantifying biodiversity: procedures and pitfalls in the measurement and comparison of species richness. *Ecol. Lett.* 4, 379–391.
- Gottwald, H.P.J., 1994. Tertiäre Kieselhölzer aus dem Chindwinn-Bassin im nordwestlichen Myanmar (Birma). *Doc. Naturae (Munich)* 86, 1–90.
- Gough, A., Hall, R., 2017. Oligocene Fluvio-Deltaic Depositional Environments Salin Sub-Basin, Central Myanmar, in: *American Geophysical Union Fall Meeting 2017, New Orleans*, abstract #EP21F-1904.
- Gough, A., Hall, R., BouDagher-Fadel, M.K., 2020. Mid-Cenozoic fluvio-deltaic to marine environments of the Salin Sub-basin, Central Myanmar. *J. Asian Earth Sci.* 190, 104143.
- Govaerts, R., Dransfield, J., Zona, S., Hodel, D.R., Henderson, A., 2015. World Checklist of Arecaceae, <http://wmsp.science.kew.org/>, facilitated by the Royal Botanic Gardens, Kew. (Retrieved on June 2015)
- Govindan, A., Mallikarjuna, U.B., 2019. An Integrated Stratigraphic and Paleoenvironmental Studies from Narmada-Tapti Block, South Cambay Basin, Western India. *J. Geol. Soc. India* 94, 171–178.
- Gower, J.C., 1971. A general coefficient of similarity and some of its properties. *Biometrics* 27, 857–874.
- Greenwood, D.R., Moss, P.T., Rowett, A.I., Vadala, A.J., Keefe, R.L., 2003. Plant communities and climate change in southeastern Australia during the early Paleogene. *Spec. Pap. Geol. Soc. Am.* 369, 365–380.
- Greenwood, D.R., Archibald, S.B., Mathewes, R.W., Moss, P.T., 2005. Fossil biotas from the Okanagan Highlands, southern British Columbia and northeastern Washington

- State: climates and ecosystems across an Eocene landscape. *Can. J. Earth Sci.* 42, 167–185.
- Grimaldi, D.A., Peñalver, E., Barrón, E., Herhold, H.W., Engel, M.S., 2019. Direct evidence for eudicot pollen-feeding in a Cretaceous stinging wasp (Angiospermae; Hymenoptera, Aculeata) preserved in Burmese amber. *Commun. Biol.* 2, 1–10.
- Grimm, E.C., 1987. CONISS: a FORTRAN 77 program for stratigraphically constrained cluster analyses by the method of incremental sum of squares. *Comput. Geosci.* 13, 13–35.
- Grimm, E.C., 1991. *Tilia and tiliagraph*. Illinois State Museum, Springfield, USA.
- Gruas-Cavagnetto, C. 1978. Etude palynologique de l'Eocene du Bassin Anglo-Parisien. *Mém. Soc. Geol. Fr. N. Ser.* 131, 1–164.
- Guédé, K.É., Slimani, H., Jean-Paul Yao, N., Chekar, M., Jean-Claude Koffi, N., M'Hamdi, A., Mouah, R., Digbehi, B.Z., 2019. Late Cretaceous to Early Eocene dinoflagellate cysts from the “12 frères” borehole, Fresco, southwestern Côte d'Ivoire: Biostratigraphy and paleobiogeographic implication. *J. Afr. Earth Sci.* 150, 744–756.
- Guimarães, J.T.F., Nogueira, A.C.R., da Silva, J.B.C., Soares, J.L., Alves, R., Kern, A.K., 2015. Palynology of the Middle Miocene–Pliocene Novo Remanso Formation, Central Amazonia, Brazil. *Ameghiniana* 52, 107–134.
- Guler, M.V., Berbach, L., Archangelsky, A., Archangelsky, S., 2015. Quistes de dinoflagelados y polen asociado del Cretácico Inferior (Formación Springhill) de la Cuenca Austral, Plataforma Continental Argentina. *Rev. Bras. Paleontol.* 18, 307–324.
- Guo, Z.T., Ruddiman, W.F., Hao, Q.Z., Wu, H.B., Qiao, Y.S., Zhu, R.X., Peng, S.Z., Wei, J.J., Yuan, B.Y., Liu, T.S., 2002. Onset of Asian desertification by 22 Myr ago inferred from loess deposits in China. *Nature* 416, 159–163.
- Haberle, S., 1997. Upper Quaternary Vegetation and Climate History of the Amazon Basin: Correlating Marine and Terrestrial Pollen Records, in: Flood, R.D., Piper, D.J.W., Klaus, A., Peterson, L.C. (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results* (vol. 155). Ocean Drilling Program, Texas A&M University, Texas, USA, pp. 381–396.
- Halbritter, H., Ulrich, S., Grímsson, F., Weber, M., Zetter, R., Hesse, M., Buchner, R., Svojtka, M., Frosch-Radivo, A., 2018. *Illustrated Pollen Terminology*. Second edition. Springer, Vienna, Austria.

REFERENCES

- Han, F., Rydin, C., Bolinder, K., Dupont-Nivet, G., Abels, H.A., Koutsodendris, A., Zhang, K., Hoorn, C., 2016. Steppe development on the Northern Tibetan Plateau inferred from Paleogene ephedroid pollen. *Grana* 55, 71–100.
- Handique, G.K., 1993. Stratigraphy, depositional environment and hydrocarbon potential of upper Assam Basin, India. Symposium on Biostratigraphy of Mainland Southeast Asia: Facies and Palaeontology, Chiang Mai, Thailand, vol. 1, pp. 151–169.
- Handique, G.K., Dutta, S.K., Neog, A., 1992. Stratigraphy, depositional environment and hydrocarbon prospects in Moran Oilfield, Upper Assam. *Geophytology* 22, 217–228.
- Hardenbol, J., Thierry, J., Farley, M.B., De Graciansky, P.-C., Vail, P.R., 1998. Mesozoic and Cenozoic chronostratigraphic framework of European basins. *SEPM Special Publ.* 60, 3–13.
- Harley, M.M., 1986. Distinguishing pollen characters for the Sapotaceae. *Can. J. Bot.* 64, 3091–3100.
- Harley, M.M., 1991. The Pollen Morphology of the Sapotaceae. *Kew Bull.* 46, 379–491.
- Harley, M.M., 2006. A summary of fossil records for Arecaceae. *Bot. J. Linn. Soc.* 151, 39–67.
- Harley, M.M., Baker, W.J., 2001. Pollen aperture morphology in Arecaceae: application within phylogenetic analyses, and a summary of the fossil record of palm-like pollen. *Grana* 40, 45–77.
- Harley, M.M., Morley, R.J., 1995. Ultrastructural studies of some fossil and extant palm pollen, and the reconstruction of the biogeographical history of subtribes Iguanurinae and Calaminae. *Rev. Palaeobot. Palynol.* 85, 153–182.
- Harrington, G.J., 2008. Comparisons between Palaeocene–Eocene paratropical swamp and marginal marine pollen floras from Alabama and Mississippi, USA. *Palaeontology* 51, 611–622.
- Harris, W.K., 1965. Basal Tertiary microfloras from the Princetown area, Victoria. *Palaeontogr. Abt. B* 115, 75–106.
- Harrison, T.M., Copeland, P., Kidd, W.S.F., Yin, A., 1992. Raising Tibet. *Science* 255, 1663–1670.
- Hartwich, S.J., Conran, J.G., Bannister, J.M., Lindqvist, J.K., 2010. Calamoid fossil palm leaves and fruits (Arecaceae: Calamoideae) from late Eocene Southland, New Zealand. *Aust. Syst. Bot.* 23, 131–140.

- Hay, W.W., Floegel, S., 2012. New thoughts about the Cretaceous climate and oceans. *Earth-Sci. Rev.* 115, 262–271.
- Hekel, H., 1972. Pollen and spore assemblages from Queensland Tertiary sediments. Publication 355, Geological Survey of Queensland.
- Hel-hama, 2013. Wikipedia Commons. (https://commons.wikimedia.org/wiki/File:Burma_administrative_divisions.svg) with a CC BY-SA 3.0 license (<https://creativecommons.org/licenses/by-sa/3.0/>). (accessed 8 September 2020)
- Helmstetter, A.J., Amoussou, B.E.N., Bethune, K., Kandem, N.G., Kakai, R.G., Sonké, B., Couvreur, T.L.P., 2020. Phylogenomic approaches reveal how climate shapes patterns of genetic diversity in an African rain forest tree species. *Mol. Ecol.* 29, 1–14.
- Henderson, A., 2020. A revision of *Calamus* (Arecaceae, Calamoideae, Calamiaceae, Calaminae). *Phytotaxa* 445, 1–656.
- Herman, A.B., Spicer, R.A., Aleksandrova, G.N., Yang, J., Kodrul, T.M., Maslova, N.P., Spicer, T.E.V., Chen, G., Jin, J.H., 2017. Eocene–early Oligocene climate and vegetation change in southern China: evidence from the Maoming Basin. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 479, 126–137.
- Herngreen, G.F.W., 1975. Palynology of Middle and Upper Cretaceous strata in Brazil. *Meded. Rijks Geol. Dienst N. Ser.* 26, 39–91.
- Herngreen, G.F.W., 1998. Cretaceous sporomorph provinces and events in the equatorial region. *Zentralblatt für Geol. Paläontol.* 1996, 1313–1323.
- Herngreen, G.F.W., Chlonova, A.F., 1981. Cretaceous Microfloral Provinces. *Pollen et Spores* 23, 441–555.
- Herngreen, G.F.W., Kedves, M., Rovnina, L.V., Smirnova, S.B., 1996. Cretaceous palynofloral provinces: a review, in: Jansonius, J., McGregor, D.C. (Eds.), *Palynology: Principles and Applications 3*. American Association of Stratigraphic Palynologists Foundation, Dallas, USA, pp. 1157–1188.
- Herrera, F., Manchester, S.R., Jaramillo, C., MacFadden, B., da Silva-Caminha, S.A., 2010. Phytogeographic history and phylogeny of the Humiriaceae. *Int. J. Plant Sci.* 171, 392–408.
- Hesse, M., Zetter, R., 2007. The fossil pollen record of Araceae. *Plant Syst. Evol.* 263, 93–115.
- Ho, K.F., 1978. Stratigraphic framework for oil exploration in Sarawak. *Bull. Geol. Soc. Malays.* 10, 1–13.

- Hofmann, C.-C., 2018. Light and scanning electron microscopic investigations of pollen of Ericales (Ericaceae, Sapotaceae, Ebenaceae, Styracaceae and Theaceae) from five lower and mid-Eocene localities. *Bot. J. Linn. Soc.* 187, 550–578.
- Hofmann, C.-C., Pancost, R., Ottner, F., Egger, H., Taylor, K., Mohamed, O., Zetter, R., 2012. Palynology, biomarker assemblages and clay mineralogy of the Early Eocene Climate Optimum (EECO) in the transgressive Krappfeld succession (Eastern Alps, Austria). *Aust. J. Earth Sci.* 105, 224–239.
- Holtum, R.E., 1953. A Revised Flora of Malaya: an Illustrated Systematic Account of the Malayan Flora, including Commonly Cultivated Plants. Government Printing Office, Singapore.
- Hooghiemstra, H., 1984. Vegetational and climatic history of the high plain of Bogotá, Colombia. *Dissertationes Botanicae*, vol. 79. J Cramer, Vaduz, 368 pp.
- Horn, C., 1993. Marine incursions and the influence of Andean tectonics on the Miocene depositional history of northwestern Amazonia: Results of a palynostratigraphic study. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 105, 267–309.
- Horn, C., 1994a. An environmental reconstruction of the palaeo-Amazon river system (Middle–Late Miocene, NW Amazonia). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 112, 187–238.
- Horn, C., 1994b. Fluvial palaeoenvironments in the Amazonas Basin (Early Miocene to early Middle Miocene, Colombia). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 109, 1–54.
- Horn, C., 2001. Pleistocene deposits of the Amazon Fan: a palynological analysis of holes 940a, 944a, and 946a (ODP leg 155), in: Goodman, D.K., Clarke, R.T. (Eds.), *Proceedings of the IX International Palynological Congress*. American Association of Stratigraphic Palynologist Foundation, Houston, Texas, USA, pp. 329–338.
- Horn, C., 2006. Mangrove forests and marine incursions in Neogene Amazonia (Lower Apaporis River, Colombia). *Palaios* 21, 197–209.
- Horn, C., Bogotá-A, G.R., Romero-Baez, M., Lammertsma, E.I., Flantua, S.G.A., Dantas, E.L., Dino, R., do Carmo, D.A., Chemale Jr, F., 2017. The Amazon at sea: Onset and stages of the Amazon River from a marine record, with special reference to Neogene plant turnover in the drainage basin. *Glob. Planet. Change* 153, 51–65.

- Hoorn, C., Straathof, J., Abels, H.A., Xu, Y.D., Utescher, T., Dupont-Nivet, G., 2012. A late Eocene palynological record of climate change and Tibetan Plateau uplift (XiningBasin, China). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 344, 16–38.
- Hoorn, C., Wesselingh, F.P., ter Steege, H., Bermudez, M.A., Mora, A., Sevink, J., Sanmartín, I., Sanchez-Meseguer, A., Anderson, C.L., Figueiredo, J.P., Jaramillo, C., Riff, D., Negri, F.R., Hooghiemstra, H., Lundberg, J., Stadler, T., Säkkinen, T., Antonelli, A., 2010. Amazonia through time: Andean uplift, climate change, landscape evolution, and biodiversity. *Science* 330, 927–931.
- Hsieh, T.C., Ma, K.H., Chao, A., 2014. iNEXT: An R package for interpolation and extrapolation in measuring species diversity.
- Htway, O., Matsumoto, J., 2011. Climatological onset dates of summer monsoon over Myanmar. *Int. J. Climatol.* 31, 382–393.
- Huang, H., Licht, A., Morley, R., Dupont-Nivet, G., Zaw Win, Westerweel, J., Littell, V., Hnin Hnin Swe, Myat Kaythi, Day Wa Aung, Roperch, P., Poblete, F., Kyaing Sein, Jardine, P., Philip, A., Hoorn, C., 2018. Palynology of the central Myanmar basin corroborates Eocene–Oligocene monsoonal conditions in south-east Asia, in: Abstract volume of the European Palaeobotany and Palynology Conference, 12–17 August 2018, Dublin, Ireland.
- Huang, H., Morley, R., Licht, A., Dupont-Nivet, G., Grímsson, F., Zetter, R., Westerweel, J., Zaw Win, Day Wa Aung, Hoorn, C., 2020. Eocene palms from central Myanmar in a South-East Asian and global perspective: evidence from the palynological record. *Bot. J. Lin. Soc.* 194, 177–206.
- Huang, H., Pérez-Pinedo, D., Morley, R.J., Dupont-Nivet, G., Philip, A., Zaw Win, Day Wa Aung, Licht, A., Jardine, P.E., Hoorn, C. At a crossroads: the late Eocene flora of central Myanmar owes its composition to colliding plate collision and tropical climate. *Rev. Palaeobot. Palynol.*, in revision.
- Huber, M., Goldner, A., 2012. Eocene monsoons. *J. Asian Earth Sci.* 44, 3–23.
- Hutchinson, D.K., Coxall, H.K., Lunt, D.J., Steinthorsdottir, M., de Boer, A.M., Baatsen, M., von der Heydt, A., Huber, M., Kennedy-Asser, A.T., Kunzmann, L., Ladant, J.-B., Lear, C.H., Moraweck, K., Pearson, P.N., Piga, E., Pound, M.J., Salzmann, U., Scher, H.D., Sijp, W.P., Śliwińska, K.K., Wilson, P.A., Zhang, Z., 2020. The Eocene–Oligocene transition: A review of marine and terrestrial proxy data, models and model-data comparisons. *Clim. Past* 17, 269–315.

- Ikegwuonu, O.N., Umeji, O.P., Chiaghanam, O.I., Nwozor, K.K., Ndukwe, O.S., Chiadikobi, K.C., 2020. Palynomorph assemblage biozonation of Paleogene strata in Bende-Umuahia Area, Niger Delta Basin, southeastern Nigeria. *J. Palaeogeogr.* 9, 13.
- Inglis, G.N., Bragg, F., Burls, N., Evans, D., Foster, G.L., Huber, M., Lunt, D.J., Siler, N., Steinig, S., Tierney, J.E., Wilkinson, R., Anagnostou, E., de Boer, A.M., Jones, T.D., Edgar, K.M., Hollis, C.J., Hutchinson, D.K., Pancost, R.D., 2020. Global mean surface temperature and climate sensitivity of the EECO, PETM and latest Paleocene. *Clim. Past* 16, 1953–1968.
- Intergovernmental Panel on Climate Change (IPCC), 2014. *Climate Change 2014: Impacts, Adaptation and Vulnerability. Contribution of working group II to the fifth assessment report of the intergovernmental panel on climate change.* Cambridge University Press, Cambridge, UK.
- International Commission on Stratigraphy (ICS), 2020. TimeScale Creator. <https://timescalecreator.org/index/index.php> (accessed 12 March 2021).
- Isnard, S., Rowe, N.P., 2008. The climbing habit in palms: Biomechanics of the cirrus and flagellum. *Am. J. Bot.* 95, 1538–1547.
- Ivanov, D.A., Ashraf, A.R., Mosbrugger, V., 2007. Late Oligocene and Miocene climate and vegetation in the Eastern Paratethys area (northeast Bulgaria), based on pollen data. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 255, 342–360.
- Jacobs, B.F., Pan, A.D., Scotese, C.R., 2010. Chapter 5: a review of the Cenozoic vegetation history of Africa, in: Werdelin, L., Sanders, W.J. (Eds.), *Cenozoic Mammals of Africa.* University of California Press, Berkeley, California, USA, pp. 57–72.
- Jacques, F.M.B., Guo, S.-X., Su, T., Xing, Y.-W., Huang, Y.-J., Liu, Y.-S., Ferguson, D.K., Zhou, Z.-K., 2011a. Quantitative reconstruction of the Late Miocene monsoon climates of southwest China: a case study of the Lincang flora from Yunnan Province. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 304, 318–327.
- Jacques, F.M.B., Su, T., Spicer, R.A., Xing, Y., Huang, Y., Wang, W., Zhou, Z., 2011b. Leaf physiognomy and climate: are monsoon systems different? *Glob. Planet. Change* 76, 56–62.
- Jaeger, J.-J., Tin Thein, Benammi, M., Chaimanee, Y., Soe Aung Naing, Lwin Thit, Than Tun, San Wai, Ducrocq, S., 1999. A new primate from the middle Eocene of Myanmar and the Asian early origin of anthropoids. *Science* 286, 528–530.

- Jais, J.H.M., 1997. Oligocene to Pliocene quantitative stratigraphic palynology of the southern Malay Basin, offshore Malaysia. Ph.D. Thesis, Sheffield University, Sheffield, UK.
- Jan du Chêne, R.E., Adegoke, O.S., Adediran, S.A., Petters, S.W., 1978a. Palynology and foraminifera of the Lokoja Sandstone (Maastrichtian), Bida Basin, Nigeria. *Rev. Esp. Micropaleontol.* 10, 379–9393.
- Jan du Chêne, R.E., Onyike, M.S., Sowunmi, M.A., 1978b. Some new Eocene pollen of the Ogwashi-Asaba Formation, southeastern Nigeria. *Rev. Esp. Micropaleontol.* 10, 285–322.
- Jaramillo, C.A., Bayona, G., Pardo-Trujillo, A., Rueda, M., Torres, V., Harrington, G.J., Mora, G., 2007. The palynology of the Cerrejón Formation (Upper Paleocene) of northern Colombia. *Palynology* 31, 153–189.
- Jaramillo, C.A., Dilcher, D.L., 2001. Middle Paleogene palynology of Central Colombia, South America: A study of pollen and spores from tropical latitudes. *Palaeontogr. Abt. B* 258, 87–259.
- Jaramillo, C., Ochoa, D., Contreras, L., Pagani, M., Carvajal-Ortiz, H., Pratt, L.M., Krishnan, S., Cardona, A., Romero, M., Quiroz, L., Rodriguez, G., Rueda, M.J., de la parra, F., Morón, S., Green, W., Bayona, G., Montes, C., Quintero, O., Ramirez, R., Mora, G., Schouten, S., Bermudez, H., Navarrete, R., Parra, F., Alvarán, M., Osorno, J., Crowley, J.L., Valencia, V., Vervoort, J., 2010. Effects of rapid global warming at the Paleocene-Eocene boundary on Neotropical vegetation. *Science* 330, 957–961.
- Jaramillo, C., Romero, I., D'Apolito, C., Bayona, G., Duarte, E., Louwye, S., Escobar, J., Luque, J., Carrillo-Briceño, J.D., Zapata, V., Mora, A., Schouten, S., Zavada, M., Harrington, G., Ortiz, J., Wesselingh, F.P., 2017. Miocene flooding events of western Amazonia. *Science* 3, e1601693.
- Jaramillo, C., Rueda, M.J., Mora, G., 2006. Cenozoic plant diversity in the Neotropics. *Science* 311, 1893–1896.
- Jaramillo, C., Rueda, M., Torres, V., 2011. A Palynological zonation for the Cenozoic of the Llanos and Llanos foothills of Colombia. *Palynology* 35, 46–84.
- Jaramillo, C., Sepulchre, P., Cardenas, D., Correa-Metrio, A., Moreno, J.E., Trejos, R., Vallejos, D., Hoyos, N., Martínez, C., Carvalho, D., Escobar, J., Oboh-Ikuenobe, F., Prámparo, M.B., Pinzón, D., 2020. Drastic vegetation change in the Guajira Peninsula (Colombia) during the Neogene. *Paleoceanogr. Paleoclimatol.* 35, e2020PA003933.

- Jardine, P.E., 2011. Spatial and temporal diversity trends in an extra-tropical megathermal vegetation type: The Early Paleogene pollen and spore record from the U.S. Gulf Coast. Ph.D. Thesis, University of Birmingham, Birmingham, UK.
- Jardine, P.E., Harrington, G.J., Sessa, J.A., Dašková, J., 2018. Drivers and constraints on floral latitudinal diversification gradients. *J. Biogeogr.* 45, 1408–1419.
- Jardiné, S., Magloire, L., 1965. Palynologie et stratigraphie du Crétacé des Bassins du Sénégal et de Côte d'Ivoire. *Mém. Bur. Rec. Géol. Min.* 32, 187–245.
- Jetz, W., Fine, P.V., 2012. Global gradients in vertebrate diversity predicted by historical area-productivity dynamics and contemporary environment. *PLoS Biol.* 10, e1001292.
- Jiménez-Moreno, G., 2006. Progressive substitution of a subtropical forest for a temperate one during the middle Miocene climate cooling in Central Europe according to palynological data from cores Tengelic-2 and Hidas-53 (Pannonian Basin, Hungary). *Rev. Palaeobot. Palynol.* 142, 1–14.
- Johnson, N.M., Stix, J., Tauxe, L., Cerveny, P.F., Tahirkheli, R.A.K., 1985. Paleomagnetic chronology, fluvial processes, and tectonic implications of the Siwalik deposits near Chinji village. Pakistan. *J. Geol.* 93, 27–40.
- Jones, E.L., 1961. Plant microfossils of the laminated sediments of the Lower Eocene Wilcox Group in south-central Arkansas. Ph.D. Thesis, University of Oklahoma, Norman, USA.
- Kar, R. K., Bhattacharya, M., 1992. Palynology of Rajpardi Lignite, Cambay Basin and Gujradam and Akri Lignite, Kutch Basin. *Palaeobotanist* 39, 250–263.
- Kaska, H.V., 1989. A spore and pollen zonation of Early Cretaceous to Tertiary nonmarine sediments of central Sudan. *Palynology* 13, 79–90.
- Kayseri-Özer, M.S., 2013. Spatial distribution of climatic conditions from the Middle Eocene to Late Miocene based on palynoflora in Central, Eastern and Western Anatolia. *Geodin. Acta* 26, 122–157.
- Kedves, M., 1992. Études palynologiques des couches du Tertiaire Inférieur de la Région Parisienne. VII. *Acta Bil. Szeged.* 38, 33–46.
- Kershaw, A.P., 1970. Pollen morphological variation in the Casuarinaceae. *Pollen et Spores* 12, 145–161.
- Kershaw A.P., 1983. A Holocene pollen diagram from Lynch's crater, north-eastern Queensland, Australia. *New Phytol.* 94, 669–682.

- Kershaw, A., 1997. A bioclimatic analysis of early to middle Miocene brown coal floras, Latrobe Valley, south-eastern Australia. *Aust. J. Bot.* 45, 373–387.
- Kershaw, A.P., Nix, H.A., 1988. Quantitative palaeoclimatic estimates from pollen data using bioclimatic profiles of extant taxa. *J. Biogeogr.* 15, 589–602.
- Khan, A.M., 1976. Palynology of Tertiary sediments from Papua New Guinea. I. New form genera and species from upper Tertiary sediments. *Aust. J. Bot.* 24, 753–781.
- Khan, M.A., Spicer, R.A., Spicer, T.E.V., Bera, S., 2016. Occurrence of *Shorea* Roxburgh ex C. F. Gaertner (Dipterocarpaceae) in the Neogene Siwalik forests of eastern Himalaya and its biogeography during the Cenozoic of Southeast Asia. *Rev. Palaeobot. Palynol.* 233, 236–254.
- Khanolkar, S., Sharma, J., 2019. Record of early to middle Eocene paleoenvironmental changes from lignite mines, western India. *J. Micropalaeontol.* 38, 1–24.
- Khorsand Rosa, R., Koptur, S., 2013. New findings on the pollination biology of *Mauritia flexuosa* (Arecaceae) in Roraima, Brazil: Linking dioecy, wind, and habitat. *Am. J. Bot.* 100, 613–621.
- Kingsley, O.K., Umeji, O.P., 2018. Oil shale prospects of Imo Formation Niger Delta Basin, southeastern Nigeria: Palynofacies, organic thermal maturation and source rock perspective. *J. Geol. Soc. India* 92, 498–506.
- Kisel, Y., McInnes, L., Toomey, N.H., Orme, C.D.L., 2011. How diversification rates and diversity limits combine to create large-scale species–area relationships. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* 366, 2514–2525.
- Kissling, W.D., Balslev, H., Baker, W.J., Dransfield, J., Gödel, B., Lim, J.Y., Onstein, R.E., Svenning, J.-C., 2019. PalmTraits 1.0, a species-level functional trait database of palms worldwide. *Sci. Data* 6, 178.
- Kissling, W.D., Eiserhardt, W.L., Baker, W.J., Borchsenius, F., Couvreur, T.L.P., Balslev, H., Svenning, J.-C., 2012. Cenozoic imprints on the phylogenetic structure of palm species assemblages worldwide. *Proc. Natl. Acad. Sci. U.S.A.* 109, 7379–7384.
- Klages, J.P., Salzmann, U., Bickert, T., Hillenbrand, C.-D., Gohl, K., Kuhn, G., Bohaty, S.M., Titschack, J., Müller, J., Frederichs, T., Bauersachs, T., Ehrmann, W., van de Flierdt, T., Pereira, P.S., Larter, R.D., Lohmann, G., Niezgodzki, I., Uenzelmann-Neben, G., Zundel, M., Spiegel, C., Mark, C., Chew, D., Francis, J.E., Nehrke, G., Schwarz, F., Smith, J.A., Freudenthal, T., Esper, O., Pälike, H., Ronge, T.A., Dziadek,

REFERENCES

- R., the Science Team of Expedition PS104, 2020. Temperate rainforests near the South Pole during peak Cretaceous warmth. *Nature* 580, 81–86.
- Klaus, S., Morley, R.J., Plath, M., Zhang, Y.P., Li, J.T., 2016. Biotic interchange between the Indian subcontinent and mainland Asia through time. *Nat. Commun.* 7, Article 12132.
- Knox, E.M., 1949. The spores of *Lycopodium*, *Phylloglossum*, *Selaginella* and *Isoetes* and their value in the study of microfossils of Paleozoic age. *Trans. Proc. Bot. Soc. Edinb.* 35, 211–357.
- Kochummen, K.M., 1989. Anacardiaceae, in: Ng, F.S.P. (Ed.), *Tree Flora of Malaya*, vol. 4. Longman Malaysia Sdn. Bhd., Petaling Jaya, pp. 9–57.
- Kocsis, A.T., Reddin, C.J., Alroy, J., Kiessling, W., 2019. The R package divDyn for quantifying diversity dynamics using fossil sampling data. *Methods Ecol. Evol.* 10, 735–743.
- Kominz, M.A., Browning, J.V., Miller, K.G., Sugarman, P.J., Mizintseva, S., Scotese, C.R., 2008. Late Cretaceous to Miocene sea-level estimates from the New Jersey and Delaware coastal plain coreholes: an error analysis. *Basin Res.* 20, 211–226.
- Kovar-Eder, J., Jechorek, H., Kvacek, Z., Parashiv, V., 2008. The integrated plant record: the ultimate tool to reconstruct Neogene zonal vegetation in Europe. *Palaios* 23, 97–111.
- Kovar-Eder, J., Kvaček, Z., 2003. Towards vegetation mapping based on the fossil plant record. *Acta Univ. Carol. Geol.* 46, 7–13.
- Kraak, M.J., Ormeling, F.J., 2003. *Cartography, Visualization of Geospatial Data* (2nd ed.). Prentice Hall, London, UK.
- Kress, W.J., DeFilipps, R.A., Farr, E., Daw Yin Yin Kyi, 2003. A checklist of the trees, shrubs, herbs, and climbers of Myanmar (revised from the original works by Lacey, J.H., Rodger, R. Hundley H.G., U Chit Ko Ko on the “List of trees, shrubs, herbs and principal climbers etc. recorded from Burma”). *Contrib. U. S. Natl. Herb.* 45, pp. 1–590.
- Kristyarin, D.A., Rahardjo, A.T., Bambang, P., 2016. Paleocology and Paleoclimate of Tanjung Formation Deposition, Based on Palynological Data From Siung Malopot, Central Borneo. *IOP Conf. Ser. Earth Environ. Sci.* 29, 012022.
- Krutzsch, W., 1959. Mikropaläontologische (sporenpaläontologische) Untersuchungen in der Braunkohle des Geiseltales. *Geologie Beih.* 21/22, 1–425.

- Krutzsch, W., 1961. Über Funde von “ephedroidem” Pollen im deutschen Tertiär. *Geologie* 10, 15–53.
- Krutzsch, W., 1970. Atlas der mittel- und jungtertiären dispersen Sporen- und Pollen- sowie der Mikroplanktonformen des nördlichen Mitteleuropas, Lieferung VII. Deutscher Verlag der Wissenschaften, Berlin, Germany.
- Krutzsch, W. 1989. Paleogeography and historical phytogeography (paleochorology) in the Neophyticum. *Plant Syst. Evol.* 162, 5–61.
- Krutzsch, W., Lenk, G., 1969. Über 5 neue Pollenarten aus dem Maastricht der Bohrung Colbitz 10 (Calvoeder Scholle, DDR).–*Monatsber. Dtsch. Akad. Wiss. Berlin* 11, 938–945.
- Kuhnhäuser, B.G., Bellot, S., Couvreur, T.L.P., Dransfield, J., Henderson, A., Schley, R., Chomicki, G., Eiserhardt, W.L., Hiscock, S.J., Baker, W.J., 2021. A robust phylogenomic framework for the calamoid palms. *Mol. Phylogenet. Evol.* 157, 107067.
- Kumar, A., Takahashi, K., 1991. Palynology of the Tertiary sediments of Southern Assam, India. *Bull. Fac. Lib. Arts, Nagasaki Univ. Nat. Sci.* 31, 515–659.
- Kumar, M., 1994. Palynological and palaeoecological studies of Palaeocene coal seams in Jarain and Laitrymbai areas, Jaintia Hills, Meghalaya, India. *Geophytology* 23, 203–207.
- Kumar, M., 1996. Palynostratigraphy and palaeoecology of Early Eocene palynoflora of Rajpardi lignite, Bharuch District, Gujarat. *Palaeobotanist* 43, 110–121.
- Kvaček, J., Herman, A.B., 2004. Monocotyledons from the Early Campanian (Cretaceous) of Grünbach, Lower Austria. *Rev. Palaeobot. Palynol.* 128, 323–353.
- Kwetche, P., Ntamak-Nida, M.J., Nitcheu, A.L.D., Etame, J., Owono, F.M., Mbese, C.O., Kissaaka, J.B.I., Ngou, G.N., Bourquin, S., Bilong, P., 2018. Facies analysis and sequence stratigraphy of Missole Outcrops: N’Kapa Formation of the south-eastern edge of Douala Sub-Basin (Cameroon). *Earth Sci. Res. Can. Cent. Sci. Edu.* 7, 35–54.
- Lai Lai Aung, Ei Ei Zin, Pwint Theingi, Naw Elvera, Phyu Phyu Aung, Thu Thu Han, Yamin Oo, Skaland, R.G., 2017. Myanmar climate report. *MET rep.* 9, 1–105.
- Laliberté, E., Legendre, P., Shipley, B., 2014. Measuring functional diversity (FD) from multiple traits, and other tools for functional ecology. *R package v. 1.0-12.*

REFERENCES

- Lang, J., Kogbe, C., Alidu, S., Alzouma, K.A., Bellion, G., Dubois, D., Durand, A., Guiraud, R., Houessou, A., de Klasz, I., Romann, E., Salard-Cheboldaeff, M., Trichet, J., 1990. The Continental terminal in West Africa. *J. Afr. Earth Sci.* 10, 79–99.
- Lasso, C.A., Rial, A., González-B, V., 2013. Morichales y Cananguchales de la Orinoquia y la Amazonia: Colombia y Venezuela. Parte I, Edition: Serie Recursos Hidrobiológicos y Pesqueros Continentales de Colombia. Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, Bogotá D.C., Colombia.
- Latham, R.E., Ricklefs, R.E., 1993. Continental comparisons of temperate-zone tree species diversity, in: Ricklefs, R.E., Schluter, D. (Eds.), *Species Diversity: Historical and Geographical Perspectives*. University of Chicago Press, Chicago, Illinois, USA, pp. 294–314.
- Leandro, L.M., Vieira, C.E.L., Santos, A., Fauth, G., 2019. Palynostratigraphy of two Neogene boreholes from the northwestern portion of the Solimões Basin, Brazil. *J. S. Am. Earth Sci.* 89, 211–218.
- Lee, J., Wang, B., Seo, K., Ha, K., Kitoh, A., Liu, J., 2015. Effects of mountain uplift on global monsoon precipitation. *Asia-Pac. J. Atmos. Sci.* 51, 275–290.
- Leffingwell, H.A., 1971. Palynology of the Lance (Late Cretaceous) and Fort Union (Paleocene) Formations in the type Lance area, Wyoming. *Geol. Soc. Am. Spec. Pap.* 127, 64 pp.
- Legoux, O., 1978. Quelques especes de pollen caracteristiques du Neogene Nigeria. *Bull. Cent. Rech. Explor. Prod. Elf-Aquitaine* 2, 265–317.
- Leidelmeyer, P., 1966. The Paleocene and Lower Eocene pollen flora of Guyana. *Leidse Geol. Meded.* 38, 49–70.
- Leite, F.P.R., 2004. Palinologia, in: Rossetti, D.F., Goes, A.M. (Eds.), *O Néogeno da Amazônia Oriental*. Museu Paraense Emilio Goeldi, Coleção Friedrich Katzer, Belém, Para, Brazil.
- Leite, F.P.R., Silva-Caminha, S.A.F.D., D'Apolito, C., 2020. New Neogene index pollen and spore taxa from the Solimões Basin (western Amazonia), Brazil. *Palynology*, 1–27.
- Lelono, E.B., 2000. Palynological study of the Eocene Nanggulan Formation, Central Java, Indonesia. Ph.D. Thesis, Royal Holloway University of London, Egham, UK.
- Lelono, E.B., 2007a. Gondwanan Palynomorphs from the Paleogene Sediment of East Java: ?The Evidence of Earlier Arrival. *LEMIGAS Sci. Contrib.* 30, 1–12.

- Lelono, E.B., 2007b. Palinomorf Eosen dari Selat Makasar. Lembaran Publikasi LEMIGAS 41, 1–10.
- Lelono, E.B., Morley, R.J., 2011. Oligocene palynological succession from the East Java Sea, in: Hall, R., Wilson, M. (Eds.), Southeast Asian Gateway Evolution. Geological Society, London, Special Publications, pp. 333–345.
- Lenz, O.K., Diekmeyer, S., Wilde, V., 2020. *Nypa* pollen diversity in middle Eocene mangroves at the edge of the southern Proto-North Sea (Helmstedt, northern Germany) – a statistical approach. *Grana* 60, 97–118.
- Li, J.F., Ferguson, D.K., Yang, J., Feng, G.P., Ablav, A.G., Wang, Y.F., Li, C.S., 2009. Early Miocene vegetation and climate in Weichang District, North China. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 280, 47–63.
- Li, J.X., Yue, L.P., Roberts, A.P., Hirt, A.M., Pan, F., Guo, L., Xu, Y., Xi, R.G., Guo, L., Qiang, X.K., Gai, C.C., Jiang, Z.X., Sun, Z.M., Liu, Q.S., 2018a. Global cooling and enhanced Eocene Asian mid-latitude interior aridity. *Nat. Commun.* 9, Article 3026.
- Li, M.-Y., 1989. Sporo-pollen from Shanghu Formation of Early Paleocene in Nanxiong Basin, Guangdong. *Acta Palaeontol. Sin.* 28, 741–749.
- Li, S., Mao, L., Spicer, R.A., Lebreton-Anberree, J., Su, T., Sun, M., Zhou, Z., 2015. Late Miocene vegetation dynamics under monsoonal climate in southwestern China. *Paleogeogr. Paleoclimatol. Paleoecol.* 425, 14–40.
- Li, S., Xing, Y., Valdes, P.G., Huang, Y., Su, T., Farnsworth, A., Lunt, D.J., Tang, H., Kennedy, A., Zhou, Z., 2018b. Oligocene climate signals and forcings in Eurasia revealed by plant macrofossils and modelling results. *Gondwana Res.* 61, 115–127.
- Licht, A., Boura, A., De Franceschi, D., Ducrocq, S., Soe Aung Naing, Jaeger, J.-J., 2014a. Fossil woods from the late middle Eocene Pondaung Formation, Myanmar. *Rev. Palaeobot. Palynol.* 202, 29–46.
- Licht, A., Boura, A., De Franceschi, D., Utescher, T., Chit Sein, Jaeger, J.-J., 2015. Late middle Eocene fossil wood of Myanmar: implications for the landscape and the climate of the Eocene Bengal Bay. *Rev. Palaeobot. Palynol.* 216, 44–54.
- Licht, A., Dupont-Nivet, G., Zaw Win, Hnin Hnin Swe, Myat Kaythi, Roperch, P., Ugrai, T., Littell, V., Park, D., Westerweel, J., Jones, D., Poblete, F., Day Wa Aung, Huang, H., Hoorn, C., Kyaing Sein. 2019. Paleogene evolution of the Burmese forearc basin and implications for the history of India-Asia convergence. *Geol. Soc. Am. Bull.* 131, 730–748.

REFERENCES

- Licht, A., France-Lanord, C., Reisberg, L., Fontaine, C., Soe Aung Naing, Jaeger, J.-J., 2013. A palaeo Tibet-Myanmar connection? Reconstructing the Late Eocene drainage system of central Myanmar using a multi-proxy approach. *J. Geol. Soc.* 170, 929–939.
- Licht, A., Reisberg, L., France-Lanord, C., Soe Aung Naing, Jaeger, J.-J., 2016. Cenozoic evolution of the central Myanmar drainage system: insights from sediment provenance in the Minbu Sub-Basin. *Basin Res.* 28, 237–251.
- Licht, A., van Cappelle, M., Abels, H.A., Ladant, J.-B., Trabucho-Alexandre, J., France-Lanord, C., Donnadiou, Y., Vandenberghe, J., Rigaudier, T., Lécuyer, C., Terry Jr, D., Adriaens, R., Boura, A., Guo, Z., Aung Naing Soe, Quade, J., Dupont-Nivet, G., Jaeger, J.-J., 2014b. Asian monsoons in a late Eocene greenhouse world. *Nature* 513, 501–506.
- Licht, A., Zaw Win, Westerweel, J., Cogné, N., Morley, C., Chantraprasert, S., Poblete, F., Ugrai, T., Nelson, B., Day Wa Aung, Dupont-Nivet, G., 2020. Magmatic history of central Myanmar and implications for the evolution of the Burma Terrane. *Gondwana Res.* 87, 303–319.
- Lim, J.Y., Marshall, C.R., 2017. The true tempo of evolutionary radiation and decline revealed on the Hawaiian archipelago. *Nature* 543, 710–713.
- Lima, N.E., Lima-Ribeiro, M.S., Tinoco, C.F., Terribile, L.C., Collevatti, R.G., 2014. Phylogeography and ecological niche modelling, coupled with the fossil pollen record, unravel the demographic history of a Neotropical swamp palm through the Quaternary. *J. Biogeogr.* 41, 673–686.
- Lindeman, J.C., 1953. The vegetation of the Coastal Region of Suriname. *Landbouwproefstation Agricultural Experiment Station Bulletin* 63, 135.
- Liu, Y.C., Tihelka, E., Yamamoto, S., Yin, Z.W., Huang, D.Y., Tian, L., Cai, C.Y., 2020. The first fossil record of the rove beetle subfamily Protopselaphinae (Coleoptera: Staphylinidae) from mid-Cretaceous Burmese amber. *Cretac. Res.* 110, 104416.
- Liu, Z., Pagani, M., Zinniker, D., DeConto, R., Huber, M., Brinkhuis, H., Shah, S.R., Leckie, R.M., Pearson, A., 2009. Global cooling during the Eocene–Oligocene climate transition. *Science* 323, 1187–1190.
- Loizeau, P.-A., Barrera, G., Manen, J.-F., Broenniman, O., 2005. Towards an understanding of the distribution of *Ilex* L. (Aquifoliaceae) on a world-wide scale, in: Friis, I., Balslev, H. (Eds.), *Plant Diversity and Complexity Patterns. Local, Regional and Global Dimensions*, vol. 55. *Biologiske Skrifter*, pp. 501–520.

- Lorente, M.A., 1986. Palynology and palynofacies of the upper Tertiary in Venezuela. Ph.D. Thesis, University of Amsterdam, Amsterdam, The Netherlands.
- Louca, S., Pennell, M.W., 2020. Extant timetrees are consistent with a myriad of diversification histories. *Nature* 580, 502–505.
- Lyle, M., Olivarez-Lyle, A., Backman, J., Tripathi, A., 2005. Biogenic sedimentation in the Eocene equatorial Pacific – the stuttering greenhouse and Eocene carbonate compensation depth, in: Wilson, P.A., Lyle, M., Firth, J.V. (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results*, pp. 1–35.
- Ma, L., Wang, Y., Fan, W., Geng, H., Cai, Y., Zhong, H., Liu, H., Xing, X., 2014. Petrogenesis of the early Eocene I-type granites in west Yingjiang (SW Yunnan) and its implication for the eastern extension of the Gangdese batholiths. *Gondwana Res.* 25, 401–419.
- Mackinder, B.A., Pennington, R.T., 2011. Monograph of *Berlinia* (Leguminosae). *Syst. Bot. Monogr.* 91, 1–117.
- Macphail, M.K., 1999. Palynostratigraphy of the Murray Basin, inland southeastern Australia. *Palynology* 23, 199–242.
- Macphail, M.K., Colhoun, E.A., Fitzsimons, S.J., 1995. Key periods in the evolution of the Cenozoic vegetation and flora in western Tasmania: the Late Pliocene. *Aust. J. Bot.* 43, 505–526.
- Macphail, M.K., Hill, R.S., 2019. What was the vegetation in northwest Australia during the Paleogene, 66–23 million years ago? *Aust. J. Bot.* 66, 556–574.
- Macphail, M.K., Hill, R.S., Forsyth, S.M., Wells, P.M., 1991. A Late Oligocene–Early Miocene cool climate flora in Tasmania. *Alcheringa* 15, 87–106.
- Macphail, M.K., Jordan, G.J., 2015. Tropical palms and arums at near-polar latitudes: fossil pollen evidence from the Tamar and Macquarie Grabens, Northern Tasmania. *Pap. Pro. R. Soc. Tasmania* 149, 23–28.
- Macphail, M.K., Stone, M.S., 2004. Age and palaeoenvironmental constraints on the genesis of the Yandi channel iron deposits, Marillana Formation, Pilbara, northwestern Australia. *Aust. J. Earth Sci.* 51, 497–520.
- Mahmoud, M.S., Schrank, E., 2007. Late Cretaceous spores, pollen and dinoflagellates from two boreholes (Nuqra-1 and 3) in the Aswan area, southeast Egypt. *Rev. Paléobiol.* 26, 593–613.

- Mairal, M., Sanmartín, I., Pellissier, L., 2017. Lineage-specific climatic niche drives the tempo of vicariance in the Rand Flora. *J. Biogeogr.* 44, 911–923.
- Malécot, V., Lobreau-Callen, D., 2005. A survey of species assigned to the fossil pollen genus *Anacolosidites*. *Grana* 44, 314–336.
- Manchester, S.R., 1989. Early history of the Juglandaceae. *Plant Syst. Evol.* 162, 231–250.
- Mandal, J., 1997. Palynofossils from the Tertiary (Barail Group) of Nagaland: Palaeoecological interpretation and age. *Palaeobotanist* 45, 98–108.
- Mandal, J., Guleria, J.S., 2006. Palynology of Vastan lignite (Surat district), Gujarat: its age, palaeoecology and depositional environment. *Paleobotanist* 55, 51–56.
- Mandal, J., Kumar, M., 2000. Stratigraphic significance of some angiosperm pollen from the Tinali oilfield, Upper Assam, India. *Palaeobotanist* 49, 197–207.
- Mandaokar, B.D., 1999. Occurrence of palynofossils from the Tirap River section (Disang Group), Tinsukia District. Assam. *Palaeobotanist* 48, 239–243.
- Mandaokar, B.D., 2003. Palynology and palaeoecological consideration of Middle Bhuvan Formation (Miocene), Lawngtlai, Mizoram, India. *Gondwana Geol. Mag. Sp.* 6, 185–193.
- Mandaokar, B.D., 2004. Age and depositional environment of the Upper Bhuvan Formation of Champhai area (Eastern Mizo Hills), India—A palynological approach. *Palaeobotanist* 53, 143–153.
- Mao, L., Foong, S.Y., 2013. Tracing ancestral biogeography of *Sonneratia* based on fossil pollen and their probable modern analogues. *Palaeoworld* 22, 133–143.
- Marivaux, L., Ducrocq, S., Jaeger, J., Marandat, B., Sudre, J., Chaimanee, Y., Soa Thura Tun, Wanna Htoon, Aung Naing Soe, 2005. New remains of *Pondaungimys anomaluropsis* (Rodentia, Anomaluroidea) from the latest middle Eocene Pondaung Formation of central Myanmar. *J. Vertebr. Paleontol.* 25, 214–227.
- Martin, H.A., 1977. The history of *Ilex* (Aquifoliaceae) with special reference to Australia: Evidence from pollen. *Aust. J. Bot.* 25, 655–673.
- Martin, H.A., 1994. Australian Tertiary phytogeography: evidence from Palynology, in: Hill, R.S. (Ed.), *History of the Australian Vegetation: Cretaceous to Recent*. University of Cambridge Press, Cambridge, UK, pp. 104–142.
- Martínez, L., Archangelsky, S., Prámparo, M., Archangelsky, A., 2016. Early Cretaceous palm pollen tetrads from Patagonia, Argentina. *Cretac. Res.* 59, 129–139.

- Mathews, R.P., Tripathi, S.M., Banerjee, S., Dutta, S., 2013. Palynology, palaeoecology and palaeodepositional environment of Eocene lignites and associated sediments from Matanomadh mine, Kutch Basin, western India. *J. Geol. Soc. India* 82, 236–248.
- Mathur, Y.K., Jain, A.K., 1980. Palynology and age of the Dras Volcanics near Shergol, Ladakh, Jammu and Kashmir, India. *Geosci. J.* 1, 55–74.
- Matthews, K.J., Maloney, K.T., Zahirovic, S., Williams, S.E., Seton, M., Müller, R.D., 2016. Global plate boundary evolution and kinematics since the late Paleozoic. *Glob. Planet. Change* 146, 226–250.
- Matzke, N.J., 2014. Model selection in historical biogeography reveals that founder-event speciation is a crucial process in island clades. *Syst. Biol.* 63, 951–970.
- Maurin, T., Rangin, C., 2009. Structure and kinematics of the Indo-Burmese Wedge: Recent and fast growth of the outer wedge. *Tectonics* 28, TC2010.
- Mbesse, C.O., 2013. La limite Paléocène–Éocène dans le Bassin de Douala Biostratigraphie et essai de reconstitution des paléoenvironnements. Ph.D. Thesis, Université of Liège, Liège, Belgium.
- McGowran, B., Archer, M., Bock, P., Darragh, T.A., Godthelp, H., Hageman, S., Hand, S.J., Hill, R., Li, Q., Maxwell, P.A., McNamara, K.J., MacPhail, M., Mildenhall, D., Partridge, A.D., Richardson, J., Shafik, S., Truswell, E.M., Warne, M., 2000. Australasian palaeobiogeography: the Paleogene and Neogene record, in: Wright, A.J., Young, G.C., Talent, J.A., Laurie, J.R. (Eds.), *Palaeobiogeography of Australasian Faunas and Floras*. Memoir 23 of the association of the Australasian Palaeontologists, Canberra, Australia, pp. 405–470.
- McLoughlin, S., 2001. The breakup history of Gondwana and its impact on pre-Cenozoic floristic provincialism. *Aust. J. Bot.* 49, 271–300.
- Mehrotra, R.C., Tiwari, R.P., Mazumder, B.I., 2003. *Nypa* megafossils from the Tertiary sediments of northeast India. *Geobios* 36, 83–92.
- Meijer, N., Dupont-Nivet, G., Abels, H.A., Kaya, M.Y., Licht, A., Xiao, M., Zhang, Y., Roperch, P., Poujol, M., Lai, Z., Guo, Z., 2019. Central Asian moisture modulated by proto-Paratethys Sea incursions since the early Eocene. *Earth Planet. Sci. Lett.* 510, 73–84.
- Mejia, K.M., Kahn, F., 1996. Biología, ecología y utilización del Irapay (*Lepidocaryum gracile* Martius). *Folia Amazónica* 8, 19–28.

- Melo, W.A., Freitas, C.G., Bacon, C.D., Collevatti, R.G., 2018. The road to evolutionary success: insights from the demographic history of an Amazonian palm. *Heredity* 121, 183–195.
- Meseguer, A.S., Lobo, J.M., Cornuault, J., Beerling, D., Ruhfel, B.R., Davis, C.C., Jousselin, E., Sanmartín, I., 2018. Reconstructing deep-time paleoclimate legacies in the clusioid Malpighiales unveils their role in the evolution and extinction of the boreotropical flora. *Glob. Ecol. Biogeogr.* 27, 616–628.
- Meseguer, A.S., Lobo, J.M., Ree, R., Beerling, D.J., Sanmartin, I., 2015. Integrating fossils, phylogenies, and niche models into biogeography to reveal ancient evolutionary history: the case of *Hypericum* (Hypericaceae). *Syst. Biol.* 64, 215–232.
- Metcalf, I., 2013. Gondwana dispersion and Asian accretion: tectonic and palaeogeographic evolution of eastern Tethys. *J. Asian Earth Sci.* 66, 1–33.
- Miao, Y., Wu, F., Herrmann, M., Yan, X., Meng, Q., 2013. Late early Oligocene East Asian summer monsoon in the NE Tibetan Plateau: Evidence from a palynological record from the Lanzhou Basin, China. *J. Asian Earth Sci.* 75, 46–57.
- Miki, A., 1977. Late Cretaceous pollen and spore floras of northern Japan: composition and interpretation. *J. Fac. Sci., Hokkaido Univ., Ser. 4. Geol. Mineral.* 17, 399–436.
- Miller, K.G., Kominz, M.A., Browning, J.V., Wright, J.D., Mountain, G.S., Katz, M.E., Sugarman, P.J., Cramer, B.S., Christie-Blick, N., Pekar, S.F., 2005. The Phanerozoic record of global sea-level change. *Science* 310, 1293–1298.
- Mitchell, A.H.G., 1993. Cretaceous–Cenozoic tectonic events in the western Myanmar (Burma)-Assam region. *J. Geol. Soc. London* 150, 1089–1102.
- Moe Zat, Day Wa Aung, 2018. Sedimentology and Sequence Stratigraphy of Letkat Formation in Kalewa-Mawleik Area, Sagaing Region. The Second Myanmar National Conference on Earth Sciences (MNCES, 2018), 29–30 November 2018, Hinthada University, Hinthada, Myanmar.
- Mohammed, Z., Awad, Z., Eisawi, A., 2017. Upper Cretaceous to Neogene palynology of the Rawat Basin, White Nile State, Sudan. *J. Earth Sci. Clim. Change* 8, 397.
- Molnar, P., Boos, W.R., Battisti, D.S., 2010. Orographic Controls on Climate and Paleoclimate of Asia: Thermal and Mechanical Roles for the Tibetan Plateau. *Annu. Rev. Earth Planet. Sci.* 38, 77–102.

- Moltzer, J.G., Binda, P.L., 1981. Micropaleontology and palynology of the middle and upper members of the Shumaysi Formation, Saudi Arabia. Bull. Fac. Earth Sci. King Abdulaziz Univ. A 4, 57–76.
- Moltzer, J.G., Binda, P.L., 1984. Age and depositional environment of the Middle and upper members of the Shumaysi formation, Saudi Arabia. Geol. Surv. Egypt Ann. 14, 269–278.
- Moreno-Domínguez, R., Cascales-Miñana, B., Ferrer, J., Diez, J.B., 2016. *Acrostitium*, a pioneering fern of floodplain areas from the Late Oligocene Sariñena Formation of the Iberian Peninsula. PLoS One 11, e0162334.
- Morley, C.K., Tin Tin Naing, Searle, M., Robinson, S.A., 2020. Structural and tectonic development of the Indo-Burma ranges. Earth-Sci. Rev. 200, 102992.
- Morley, R.J., 1976. Vegetation change in West Malaysia during the late Quaternary period: a palynological study of selected lowland and lower montane sites. Ph.D. Thesis, the University of Hull, Hull, UK.
- Morley, R.J., 1978. Palynology of Tertiary and Quaternary sediments in Southeast Asia, in: Proceeding of Indonesian Petroleum Association 6th Annual Convention, pp. 255–276.
- Morley, R.J., 1982. Fossil pollen attributable to *Alangium* Lamarck (Alangiaceae) from the Tertiary of Malesia. Rev. Palaeobot. Palynol. 36, 65–94.
- Morley, R.J., 1991. Tertiary stratigraphic palynology in Southeast Asia: current status and new directions. Bull. Geol. Soc. Malays. 28, 1–36.
- Morley, R.J., 1996. Biostratigraphic characterisation of systems tracts in Tertiary sedimentary basins, in: Proceedings of the International Symposium on Sequence Stratigraphy in SE Asia, IPA Jakarta, pp. 49–71.
- Morley, R.J., 1998. Palynological evidence for Tertiary plant dispersal in the SE Asian region in relation to plate tectonics and climate, in: Hall, R., Halloway, J.D. (Eds.), Biogeography and Geological Evolution of SE Asia. Backhuys, Leiden, The Netherlands, pp. 211–234.
- Morley, R.J., 2000. Origin and Evolution of Tropical Rain Forests. John Wiley and Sons, Ltd, Chichester, UK.
- Morley, R.J., 2003. Interplate dispersal paths for megathermal angiosperms. Perspect. Plant Ecol. Evol. Syst. 6, 5–20.

- Morley, R.J., 2007. Cretaceous and Tertiary climate change and the past distribution of megathermal rainforests, in: Bush, M.B., Flenley, J.R. (Eds.), *Tropical Rainforest Responses to Climatic Changes*. Praxis Publishing, Chichester, UK, pp. 1–31.
- Morley, R.J., 2012. A review of the Cenozoic palaeoclimate history of Southeast Asia, in: Gower, D.J., Johnson, K.G., Richardson, J.E., Rosen, B.R., Rüber, L., Williams, S.T. (Eds.), *Biotic Evolution and Environmental Change in Southeast Asia*. Cambridge University Press, Cambridge, UK, pp. 79–114.
- Morley, R.J., 2013. Cenozoic ecological history of South East Asian peat mires based on the comparison of coals with present day and Late Quaternary peats. *J. Limnol.* 72, 36–59.
- Morley, R.J., 2014. Rifting and mountain building across Sundaland, a palynological and sequence biostratigraphic perspective, in: *Proceedings of the 38th Annual Convention and Exhibition of Indonesian Petroleum Association*. Indonesian Petroleum Association, Jakarta, Indonesia.
- Morley, R.J., 2018a. Assembly and division of the South and South-East Asian flora in relation to tectonics and climate change. *J. Trop. Ecol.* 34, 209–234.
- Morley, R.J., 2018b. The complex history of mountain building and the establishment of mountain biota in Southeast Asia and Eastern Indonesia, in: Hoorn, C., Perrigo, A., Antonelli, A. (Eds.). *Mountains, Climate and Diversity*. Wiley-Blackwell, Oxford, UK, pp. 475–493.
- Morley, R.J., 2020. Palynology in Indonesia: an overview of Quaternary studies, Cenozoic stratigraphic palynology and Sequence biostratigraphy. FOSI IAGI Webinar, 4 July 2020, online.
- Morley, R.J., Bui, V.D., Nguyen, T.T., Kullman, A.J., Bird, R.T., Nguyen, V.K., Nguyen, H.C., 2019. High resolution Palaeogene sequence stratigraphic framework for the Cuu Long Basin, offshore Vietnam, driven by climate change and tectonics, established from sequence biostratigraphy. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 530, 113–135.
- Morley, R.J., Hasan, S.S., Morley, H.P., Jais, J.H.M., Mansor, A., Aripin, M.R., Nordin, M.H., Rohaizar, M.H., 2021. Sequence biostratigraphic framework for the Oligocene to Pliocene of Malaysia: High-frequency depositional cycles driven by polar glaciation. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 561, 110058.

- Morley, R.J., Lelono, E.B., Nugrahaningsih, L., Hasjim, N., 2000. LEMIGAS Tertiary Palynology Project: aims, progress and preliminary results from the Middle Eocene to Late Pliocene of Sumatra and Java, in: *Paleontology Series, Part 1: Environmental Photochemistry*, vol. 10. Geological Research and Development Centre, pp. 27–47.
- Morley, R.J., Morley, H.P., 2010. Neogene climate history of the Makassar Straits, with emphasis on the Attaka region, East Kalimantan, Indonesia. 34th Annual Convention and Exhibition Indonesia Petroleum Association. Indonesian Petroleum Association, Jakarta, Indonesia.
- Morley, R.J., Morley, H.P., 2013. Mid Cenozoic freshwater wetlands of the Sunda region. *J. Limnol.* 72, 18–35.
- Morley, R.J., Morley, H.P., 2018. Montane pollen indicates character of Mid Cenozoic uplands across Sunda Shelf, in: *PESGB SEAPEX Asia Pacific E&P Conference*, 27–28 June 2018, Olympia Exhibition Centre, London, UK.
- Morley, R.J., Morley, H.P., 2021. The prelude to the Holocene: tropical Asia during the Pleistocene, in: Navnith, K., Kumaran, P., Padmalal, D. (Eds.), *Holocene Climate Change and Environment*. Elsevier, in press.
- Morley, R.J., Morley, H.P., Restrepo-Pace, P., 2003. Unravelling the tectonically controlled stratigraphy of the West Natuna Basin by means of palaeo-derived mid-Tertiary climate changes, in: *Proceeding of the Indonesian Petroleum Association 29th Annual Convention and Exhibition*. Indonesian Petroleum Association, Jakarta, Indonesia, pp. 264–288.
- Mosbrugger, V., Utescher, T., 1997. The coexistence approach—a method for quantitative reconstructions of Tertiary terrestrial palaeoclimate data using plant fossils. *Paleogeogr. Palaeoclimatol. Palaeoecol.* 134, 61–86.
- Muller, J., 1966. Montane pollen from the Tertiary of NW. Borneo. *Blumea* 14, 231–235.
- Muller, J., 1968. Palynology of the Pedawan and Plateau Sandstone Formations (Cretaceous–Eocene) in Sarawak, Malaysia. *Micropaleontology* 14, 1–37.
- Muller, J., 1970. Palynological evidence on early differentiation of angiosperms. *Biol. Rev. Camb. Philos. Soc.* 45, 417–50.
- Muller, J., 1972. Palynological evidence for change in geomorphology, climate and vegetation in the Mio-Pliocene of Malesia, in: Ashton, P.S., Ashton, M. (Eds.), *The Quaternary Era in Malesia*. Geography Department, University of Hull, Miscellaneous Series 13, pp. 6–34.

- Muller, J., 1981. Fossil pollen of extant angiosperms. *Bot. Rev.* 47, 1–142.
- Muller, J., di Giacomo, E., van Erve, A.W., 1987. A palynological zonation for the Cretaceous, Tertiary, and Quaternary of northern South America. *Am. Assoc. Strat. Palynol. Contrib. Ser.* 19, 7–76.
- Muscarella, R., Emilio, T., Phillips, O.L., Lewis, S.L., Slik, F., Baker, W.J., Couvreur, T.L.P., Eiserhardt, W.L., Svenning, J.C., Affum-Baffoe, K., Aiba, S.I., de Almeida, E.C., de Almeida, S.S., de Oliveira, E.A., Álvarez-Dávila, E., Alves, L.F., Alvez-Valles, C.M., Carvalho, F.A., Guarín, F.A., Andrade, A., Aragão, L.E.O.C., Murakam, A.A., Arroyo, L., Ashton, P.S., Corredor, G.A.A., Baker, T.R., de Camargo, P.B., Barlow, J., Bastin, J.F., Bengone, N.N., Berenguer, E., Berry, N., Blanc, L., Böhning-Gaese, K., Bonal, D., Bongers, F., Bradford, M., Brambach, F., Brearley, F.Q., Brewer, S.W., Camargo, J.L.C., Campbell, D.G., Castilho, C.V., Castro, W., Catchpole, D., Martínez, C.E.C., Chen, S., Chang, P., Cho, P., Chutipong, W., Clark, C., Collins, M., Comiskey, J.A., Medina, M.N.C., Costa, F.R.C., Culmsee, H., David-Higuita, H., Davidar, P., del Aguila-Pasquel, J., Derroire, G., Fiore, A.D., Do, T.V., Doucet, J.L., Dourdain, A., Drake, D.R., Ensslin, A., Erwin, T., Ewango, C.E.N., Ewers, R.M., Fauset, S., Feldpausch, T.R., Ferreira, J., Ferreira, L.V., Fischer, M., Franklin, J., Fredriksson, G.M., Gillespie, T.W., Gilpin, M., Gonmadje, C., Gunatilleke, A.U.N., Hakeem, K.R., Hall, J.S., Hamer, K.C., Harris, D.J., Harrison, R.D., Hector, A., Hemp, A., Herault, B., Pizango, C.G.H., Coronado, E.N.H., Hubau, W., Hussain, M.S., Ibrahim, F.H., Imai, N., Joly, C.A., Joseph, S., K, A., Kartawinata, K., Kassi, J., Killeen, T.J., Kitayama, K., Klitgård, B.B., Kooyman, R., Labrière, N., Larney, E., Laumonier, Y., Laurance, S.G., Laurance, W.F., Lawes, M.J., Levesley, A., Lisingo, J., Lovejoy, T., Lovett, J.C., Lu, X., Lykke, A.M., Magnusson, W.E., Mahayani, N.P.D., Malhi, Y., Mansor, A., Peña, J.L.M., Marimon-Junior, B.H., Marshall, A.R., Melgaco, K., Bautista, C.M., Mihindou, V., Millet, J., Milliken, W., Mohandass, D., Mendoza, A.L.M., Mugerwa, B., Nagamasu, H., Nagy, L., Seuaturien, N., Nascimento, M.T., Neill, D.A., Neto, L.M., Nilus, R., Vargas, M.P.N., Nurtjahya, E., de Araújo, R.N.O., Onrizal, O., Palacios, W.A., Palacios-Ramos, S., Parren, M., Paudel, E., Morandi, P.S., Pennington, R.T., Pickavance, G., Pipoly III, J.J., Pitman, N.C.A., Poedjirahajoe, E., Poorter, L., Poulsen, J.R., Prasad, P.R.C., Prieto, A., Puyravaud, J.P., Qie, L., Quesada, C.A., Ramírez-Angulo, H., Razafimahaimodison, J.C., Reitsma, J.M., Requena-Rojas, E.J., Correa, Z.R., Rodriguez, C.R., Roopsind, A., Rovero, F.,

- Rozak, A., Lleras, A.R., Rutishauser, E., Rutten, R., Punchi-Manage, R., Salomão, R.P., Sam, H.V., Sarker, S.K., Satdichanh, M., Schietti, J., Schmitt, C.B., Marimon, B.S., Senbeta, F., Sharma, L.N., Sheil, D., Sierra, R., Silva-Espejo, J.E., Silveira, M., Sonké, B., Steininger, M.K., Steinmetz, R., Stévant, T., Sukumar, R., Sultana, A., Sunderland, T.C.H., Suresh, H.S., Tang, T., Tanner, E., Steege, H.T., Terborgh, J.W., Theilade, I., Timberlake, T., Torres-Lezama, A., Umunay, P., Uriarte, M., Gamarra, L.V., van de Bult, M., van der Hout, P., Martinez, R.V., Vieira, I.C.G., Vieira, S.A., Vilanova, E., Cayo, J.V., Wang, O., Webb, C.O., Webb, E.L., White, L., Whitfeld, T.J.S., Wich, S., Willcock, W., Wiser, S.K., Young, K.R., Zakaria, Z., Zang, R., Zartman, C.E., Zo-Bi, I.C., Balslev, H., 2020. The global abundance of tree palms. *Glob. Ecol. Biogeogr.* 29, 1495–1514.
- Muscarella, R., Galante, P.J., Soley-Guardia, M., Boria, R.A., Kass, J.M., Uriarte, M., Anderson, R.P., 2014. ENMeval: An R package for conducting spatially independent evaluations and estimating optimal model complexity for Maxent ecological niche models. *Methods Ecol. Evol.* 5, 1198–1205.
- Myers, N., Mittermeier, R., Mittermeier, G.C., Dafonseca, G.A.B., Kent, J., 2000. Biodiversity hotspots for conservation priorities. *Nature* 403, 853–858.
- Nagappa, Y., 1959, Foraminiferal biostratigraphy of the Cretaceous–Eocene succession in the India-Pakistan-Burma region. *Micropaleontology* 5, 145–177.
- Navarro, J.A., Galeano, G., Bernal, R., 2011. Impact of leaf harvest on populations of *Lepidocaryum tenue*, an Amazonian understory palm used for thatching. *Trop. Conserv. Sci.* 4, 25–38.
- Ng, F.S.P., 1972. Juglandaceae, in: Whitmore, T.C. (Ed.), *Tree flora of Malaya*, vol. 1. Longman, Kuala Lumpur and London, pp. 233–236.
- Ngon Ngon, G.F., Etame, J., Ntamak-Nida, M.J., Mbesse, C.O., Mbaï, J.S., Bayiga, E. C., Gerard, M., 2016. Geochemical and palaeoenvironmental characteristics of Missole I iron duricrusts of the Douala sub-basin (Western Cameroon). *C. R. Geosci.* 348, 127–137.
- Nguyen, H.C., Morley, R.J., Bui, V.D., Cao, D.H.Y., Nguyen, V.S., 2011. Sequence biostratigraphy of the late Oligocene to Miocene succession in the northern Song Hong Basin, offshore Vietnam. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 569, 110322.

REFERENCES

- Ni, J., Yu, G., Harrison, S.P., Prentice, C., 2010. Palaeovegetation in China during the late Quaternary: Biome reconstructions based on a global scheme of plant functional types. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 289, 44–61.
- Nogueira, A.C.R., Silveira, R., Guimarães, J.T.F., 2013. Neogene–Quaternary sedimentary and paleovegetation history of the eastern Solimões Basin, central Amazon region. *J. S. Am. Earth Sci.* 46, 89–99.
- Nürk, N.M., Linder, H.P., Onstein, R.E., Larcombe, M.J., Hughes, C.E., Piñeiro Fernández, L., Schlüter, P.M., Valente, L., Beierkuhnlein, C., Cutts, V., Donoghue, M.J., Edwards, E.J., Field, R., Flantua, S.G.A., Higgins, S.I., Jentsch, A., Liede-Schumann, S., Pirie, M.D., 2020. Diversification in evolutionary arenas—Assessment and synthesis. *Ecol. Evol.* 10, 6163–6182.
- Ochoa, D., Hoorn, C., Jaramillo, C., Bayona, G., Parra, M., de la Parra, F., 2012. The final phase of tropical lowland conditions in the axial zone of the Eastern Cordillera of Colombia: Evidence from three palynological records. *J. S. Am. Earth Sci.* 39, 157–169.
- Okeke, K.K., Umeji, O.P., 2016. Palynostratigraphy, palynofacies and palaeoenvironment of deposition of Selandian to Aquitanian sediments, southeastern Nigeria. *J. S. Am. Earth Sci.* 120, 102–124.
- Oloto, I.N., 1990. Palynological assemblage from the Danian of South-West Nigeria. *Acta Palaeobot.* 30, 23–39.
- Oloto, I.N., Promise, W., 2014. Biostratigraphic study and paleoenvironmental reconstruction of cores from offshore (South Western) Niger Delta, Nigeria. *Int. J. Sci. Technol. Res.* 3, 219–286.
- Onstein, R.E., Baker, W.J., Couvreur, T.L.P., Faurby, S., Svenning, J.-C., Kissling, W. D., 2017. Frugivory-related traits promote speciation of tropical palms. *Nat. Ecol. Evol.* 1, 1903–1911.
- Onuigbo, E.N., Okoro, A.U., Etu-Efeotor, J.O., Akpunonu, E.O., Okeke, H.C., 2015. Paleocology of Enugu and Mamu Formations in Anambra Basin, Southeastern Nigeria. *Adv. Appl. Sci. Res.* 6, 23–39.
- Osorio-Granada, E., Pardo-Trujillo, A., Restrepo-Moreno, S.A., Gallego, F., Muñoz, J., Plata, A., Trejos-Tamayo, R., Vallejo, F., Barbosa-Espitia A., Cardona-Sánchez, F.J., Foster, D.A., Kamenov, G., 2020. Provenance of Eocene–Oligocene sediments in the

- San Jacinto Fold Belt: Paleogeographic and geodynamic implications for the Northern Andes and the southern Caribbean. *Geosphere* 16, 210–228.
- Palamarczuk, S., Barreda, V.D., 2000. Palinología del Paleogeno tardío-Neogeno temprano, pozo Aries x-1, plataforma continental Argentina, Tierra del Fuego. *Ameghiniana* 37, 221–234.
- Palazzesi, L., Barreda, V., Cuitiño, J., Guler, M., Tellería, M., Santos, R.V., 2014. Fossil pollen records indicate that Patagonian desertification was not solely a consequence of Andean uplift. *Nat. Commun.* 5, 3558.
- Pälike, H., Lyle, M.W., Nishi, H., Raffi, I., Ridgwell, A., Kusali, G., Klaus, A., Acton, G., Anderson, L., Backman, J., Baldauf, J., Beltran, C., Bohaty, S.M., Busch, P.B.W., Channell, J.E.T., Chun, C.O.J., Delaney, M., Dewangan, P., Jones, T.D., Edgar, K.M., Evans, H., Fitch, P., Foster, G.L., Gussone, N., Hasegawa, H., Hathorne, E.C., Hayashi, H., Herrle, J.O., Holbourn, A., Hovan, S., Hyeong, K., Iijima, K., Ito, T., Kamikuri, S.-i., Kimoto, K., Kuroda, J., Leon-Rodriguez, L., Malinverno, A., Moore Jr., T.C., Murphy, B.H., Murphy, D.P., Nakamura, H., Ogane, K., Ohneiser, C., Richter, C., Robinson, R., Rohling, E.J., Romero, O., Sawada, K., Scher, H., Schneider, L., Sluijs, A., Takata, H., Tian, J., Tsujimoto, A., Wade, B.S., Westerhold, T., Wilkens, R., Williams, T., Wilson, P.A., Yamamoto, Y., Yamamoto, S., Yamazaki, T., Zeebe, R.E., 2012. A Cenozoic record of the equatorial Pacific carbonate compensation depth. *Nature* 488, 609–614.
- Pälike, H., Norris, R.D., Herrle, J.O., Wilson, P.A., Coxall, H.K., Lear, C.H., Shackleton, N.J., Tripathi, A., Wade, B.S., 2006. The heartbeat of the Oligocene climate system. *Science* 314, 1894–1898.
- Palynodata Inc., White, J.M., 2008. Palynodata Datafile: 2006 version, with introduction by White JM. Geological Survey of Canada Open File 5793, 1 CD-ROM.
- Pan, A.D., Jacobs, B.F., Dransfield, J., Baker, W.J., 2006. The fossil history of palms in Africa and new records from the Late Oligocene (~28–27 Myr) of northwestern Ethiopia. *Bot. J. Linn. Soc.* 151, 69–81.
- Paradis, E., Claude, J., Strimmer, K., 2004. APE: Analyses of phylogenetics and evolution in R language. *Bioinformatics* 20, 289–290.
- Pardo-Trujillo, A., Jaramillo, C.A., 2014. Palynology and Paleoenvironments of Eastern Cordillera Paleogene deposits of Colombia: 35 million years of Neotropical vegetation history, in: Rangel, J.O. (Ed.), *Colombia Diversidad Biológica XIV: La región de la*

REFERENCES

- Orinoquia de Colombia. Universidad Nacional de Colombia Instituto de Ciencias Naturales, Bogotá D.C., Colombia, pp. 1–30.
- Pardo-Trujillo, A., Jaramillo, C.A., Oboh-Ikuenobe, F.E., 2003. Paleogene palynostratigraphy of the eastern middle Magdalena Valley, Colombia. *Palynology* 27, 155–178.
- Pardo-Trujillo, A., Roche, E., 2009. Paleocene–Eocene palynology and palynofacies from northeastern Colombia and western Venezuela. *Cuadernos de Investigación* 41, Universidad de Caldas, Caldas, Colombia.
- Pearson, P.N., Foster, G.L., Wade, B.S., 2009. Atmospheric carbon dioxide through the Eocene-Oligocene climate transition. *Nature* 461, 1110–1113.
- Pennington, T.D., 1991. *The Genera of the Sapotaceae*. Kew Publishing, London, UK.
- Penny, D., 2001. A 40,000 year palynological record from north-east Thailand: implications for biogeography and palaeo-environmental reconstruction. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 171, 97–128.
- Phadtare, N.R., Kulkarni, A.R., 1984. Palynological assemblage of lignite exposure of Ratnagiri District, in: Badvae, R.M., Maharashtra Association for the Cultivation of Science (Eds.), *Proceedings of the X Indian Colloquium on the Indian Micropalaeontology and Stratigraphy*. Maharashtra Association of Cultivation of Science, Pune, India, pp. 515–531.
- Phadtare, N.R., Thakur, B., 1990. Fossil pollen of *Alangium* from the Eocene lignite of Gujarat, India, with comments on its stratigraphic antiquity. *Rev. Palaeobot. Palynol.* 63, 281–297.
- Phillips, S., Anderson, R.P., Dudik, M., Schapire, R., Blair, M., 2017. Opening the black box: An open-source release of Maxent. *Ecography* 40, 887–893.
- Pieters, P.E., Trail, D.S., Supriatna, S., 1987. Correlation of Early Tertiary rocks across Kalimantan, in: *Proceedings Indonesian Petroleum Association 16th Annual Convention*. IPA87-11/11, pp. 1–16.
- Pivnik, D.A., Nahm, J., Tucker, R.S., Smith, G.O., Nyein, K., Nyunt, M., Maung, P.H., 1998. Polyphase deformation in a fore-arc/back-arc basin, Salin sub-basin, Myanmar (Burma). *Am. Assoc. Pet. Geol. Bull.* 82, 1837–1856.
- Playford, G., 1982. Neogene palynomorphs from the Huon Peninsula, Papua New Guinea. *Palynology* 6, 29–54.

- Pocknall, D.T., 1982. Palynology of Late Oligocene Pomahaka estuarine bed sediments, Waikoikoi, Southland, New Zealand. *N. Z. J. Bot.* 20, 263–287.
- Pocknall, D.T., Erlich, R.N., Stein, J.A., Bergen, J.A., Lorente, M.A., 2001. The palynoflora succession across the Cretaceous to Paleocene transition zone, Mérida Andes, western Venezuela, in: Goodman, D.K., Clarke, R.T. (Eds.), *Proceedings of the IX International Palynological Congress*, Houston, Texas, USA. American Association of Stratigraphic Palynologists Foundation, USA, pp. 171–179.
- Pocknall, D.T., Jarzen, D.M., 2012. *Grimsdalea magnaclavata* Germeraad, Hopping & Muller: an enigmatic pollen type from the Neogene of northern South America. *Palynology* 36, 134–143.
- Poinar, G., 2018. Burmese amber: evidence of Gondwanan origin and Cretaceous dispersion. *Hist. Biol.* 31, 1304–1309.
- Pole, M.S., 1998. The Proteaceae record in New Zealand. *Aust. Syst. Bot.* 11, 343–372.
- Pole, M.S., Macphail, M.K., 1996. Eocene *Nypa* from Regatta Point, Tasmania. *Rev. Palaeobot. Palynol.* 92, 55–67.
- Potonié, R., 1960. Sporologie der eozänen Kohle von Kalewa in Burma. *Senckenb. Leth.* 41, 451–481.
- Poux, C., Chevret, P., Huchon, D., de Jong, W.W., Douzery, E.J.P., 2006. Arrival and diversification of Caviomorph rodents and Platyrrhine primates in South America. *Syst. Biol.* 55, 228–244.
- Prakash, U., 1965. Fossil wood of *Lagerstroemia* from the Tertiary of Burma. *Curr. Sci.* 16, 484–485.
- Prakash, U., 1971. Fossil woods from the Tertiary of Burma. *Palaeobotanist* 20, 48–70.
- Prakash, U., Bande, M.B., 1980. Some more fossil woods from the Tertiary of Burma. *Palaeobotanist* 26, 261–278.
- Prasad, V., Farooqui, A., Murthy, S., Sarate, O.S., Bajpai, S., 2018a. Palynological assemblage from the Deccan Volcanic Province, central India: insights into early history of angiosperms and the terminal Cretaceous paleogeography of peninsular India. *Cretac. Res.* 86, 186–198.
- Prasad, V., Farooqui, A., Tripathi, S.K.M., Garg, R., Thakur, B., 2009. Evidence of Late Palaeocene–Early Eocene equatorial rain forest refugia in southern Western Ghats, India. *J. Biosci.* 34, 777–797.

REFERENCES

- Prasad, V., Singh, I.B., Bajpai, S., Garg, R., Thakur, B., Singh, A., Saravanan, N., Kapur, V.V., 2013. Palynofacies and sedimentology-based high-resolution sequence stratigraphy of the lignite-bearing muddy coastal deposits (early Eocene) in the Vastan Lignite Mine, Gulf of Cambay, India. *Facies* 59, 737–761.
- Prasad, V., Utescher, T., Sharma, A., Singh, I.B., Garg, R., Gogoi, B., Srivastava, J., Uddandam, P.R., Joachimski, M.M., 2018b. Low-latitude vegetation and climate dynamics at the Paleocene Eocene transition—A study based on multiple proxies from the Jathang section in northeastern India. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 497, 139–156.
- Prell, W.L., Kutzbach, J.E., 1992. Sensitivity of the Indian monsoon to forcing parameters and implications for its evolution. *Nature* 360, 647–652.
- Privé-Gill, C., Vozenin-Serra, C., Ducrocq, S., Soe Aung Naing, Jaeger, J.-J., 2004. Bois de la Formation Pondaung, Eocène moyen du Myanmar. Implications paléoenvironnementales. *Palaeontogr. Abt. B* 267, 57–65.
- Punt, W., Hoen, P.P., Blackmore, S., Nilsson, S., Le Thomas, A., 2007. Glossary of pollen and spore terminology. *Rev. Palaeobot. Palynol.* 143, 1–81.
- Quental, T.B., Marshall, C.R., 2010. Diversity dynamics: molecular phylogenies need the fossil record. *Trends Ecol. Evol.* 25, 434–441.
- Quental, T.B., Marshall, C.R., 2011. The molecular phylogenetic signature of clades in decline. *PLoS One* 6, e25780.
- R Core Team, 2018. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.r-project.org>.
- R Core Team, 2019. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.r-project.org>.
- Raatz, G.V., 1938. Mikrobotanisch-stratigraphische Untersuchungen der Braunkohle des Muskauer Bogens. *Abh. Preuss. Geol. Landesanst. (Berlin)* 183, 1–48.
- Rabosky, D.L., 2014. Automatic detection of key innovations, rate shifts, and diversity-dependence on phylogenetic trees. *PLoS One* 9, e89543.
- Rachman, R.S., Winantris, Muljana, B., 2021. Age and depositional Environment of Walat Formation Based on Palynological Analysis in Sukabumi Regency, West Java, Indonesia. *Pak. J. Geol.* 5, 1–7.
- Ramanujam, C.G.K., 1987. Palynology of the Neogene Warkalli Beds of Kerala State in South India. *J. Palaeontol. Soc. India* 32, 26–46.

- Ramanujam, C.G.K., Reddy, P.R., Rao, M.G., 1991. Palynoassemblages of the subsurface Tertiary strata at Pattanakad, Alleppey District, Kerala State. *J. Palaeontol. Soc. India* 36, 51–58.
- Ramírez-Arriaga, E., Pramparo, M.B., Martínez-Hernández, E., Valiente-Baunet, A., 2006. Palynology of the Paleogene Cuayuca Formation (stratotype sections), southern Mexico: chronostratigraphical and palaeoecological implications. *Rev. Palaeobot. Palynol.* 141, 259–275.
- Ramli, N., 1988. Stratigraphy and palaeofacies development of Carigali's operating areas in the Malay Basin, South China Sea. *Geol. Soc. Malaysia Bull.* 22, 153–187.
- Rao, K.P., Ramanujam, C.G.K., 1982. Palynology of the Neogene Quilon beds of Kerala State in south India II. Pollen of dicotyledons and discussion. *Palaeobotanist* 30, 68–100.
- Rao, M.R., Sahni, A., Rana, R.S., Verma, P., 2013. Palynostratigraphy and depositional environment of Vastan Lignite Mine (Early Eocene), Gujarat, western India. *J. Earth Syst. Sci.* 122, 289–307.
- Ratnam, J., Tomlinson, K.W., Rasquinha, D.N., Sankaran, M., 2016. Savannahs of Asia: antiquity, biogeography, and an uncertain future. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* 371, 20150305.
- Rawat, M.S., Mukherjee, J.S., Venkatachala, B.S., 1977. Palynology of the Kadi Formation, Cambay Basin, India, in: Venkatachala, B.S., Sastri, V.V. (Eds.), *Proceedings of the 4th Colloquium on Indian Micropalaeontology and Stratigraphy, Dehradun, 1974–1975*. Institute of Petroleum Exploration, Oil and Natural Gas Commission, India, pp. 179–192.
- Raymer, J.D., 2010. Cretaceous/Paleogene boundary biostratigraphy and palynofacies of the Alo-1 well, Southeastern Nigeria. MSc Thesis, Missouri University of Science and Technology, Rolla, USA.
- Ree, R.H., Smith, S.A. 2008. Maximum likelihood inference of geographic range evolution by dispersal, local extinction, and cladogenesis. *Syst. Biol.* 57, 4–14.
- Regali, M.S.P., Uesugui, N., Santos, A.S., 1974. Palinología dos sedimentos Mesozoicos do Brasil (II). *Bol. Téc. Petrobrás* 17, 263–301.
- Reichgelt, T., Kennedy, E.M., Mildenhall, D.C., Conran, J.G., Greenwood, D.R., Lee, D.E., 2013. Quantitative palaeoclimate estimates for Early Miocene southern New Zealand: evidence from Foulden Maar. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 378, 36–44.

REFERENCES

- Reichgelt, T., West, C.K., Greenwood, D.R., 2018. The relation between global palm distribution and climate. *Sci. Rep.* 8, 4721.
- Reimann, K.U., Aye Thaug, 1981. Results of palynostratigraphical investigations of the Tertiary sequence in the Chindwin Basin/Northwestern Burma, in: Bharadwaj, D.C. (Ed.), *Proceedings of IV International Palynological Conference*. Birbal Sahni Institute of Palaeobotany, Lucknow, India, pp. 380–395.
- Renema, W., Bellwood, D.R., Braga, J.C., Bromfield, K., Hall, R., Johnson, K.G., Lunt, P., Meyer, C.P., McMonagle, L.B., Morley, R.J., O’Dea, A., Todd, J.A., Wesselingh, F.P., Wilson, M.E.J., Pandolfi, J.M., 2008. Hopping Hotspots: global Shifts in Marine Biodiversity. *Science* 321, 654–657.
- Renner, S., 2004. Plant dispersal across the Tropical Atlantic by wind and sea currents. *Int. J. Plant Sci.* 165, S23–S33.
- Revell, L.J., 2012. phytools: An R package for phylogenetic comparative biology (and other things). *Methods Ecol. Evol.* 3, 217–223.
- Richardson, J., Bakar, A., Tosh, J., Armstrong, K., Smedmark, J., Anderberg, A., Slik, F., Wilkie, P., 2014. The influence of tectonics, sea-level changes and dispersal on migration and diversification of *Isonandreae* (Sapotaceae). *Bot. J. Linn. Soc.* 174, 130–140.
- Riegel, W., Lenz, O.K., Wilde, V., 2015. From open estuary to meandering river in a greenhouse world: an ecological case study from the middle Eocene of Helmstedt, Northern Germany. *Palaios* 30, 304–326.
- Riegel, W., Wilde, V., Lenz, O.K., 2012. The early Eocene of Schöningen (N-Germany)—an interim report. *Aust. J. Earth Sci.* 105, 88–109.
- Rodríguez-Forero, G., Oboh-Ikuenobe, F.E., Jaramillo-Muñoz, C., Rueda-Serrano, M. J., Cadena-Rueda, E., 2012. Palynology of the Eocene Esmeraldas Formation, Middle Magdalena Valley Basin, Colombia. *Palynology* 36, 96–111.
- Ronquist, F., Sanmartín, I., 2011. Phylogenetic methods in biogeography. *Annu. Rev. Ecol. Evol. Syst.* 42, 441–464.
- Ruddle, K., 1979. The distribution of sago-producing palms. *Bull. Natl. Mus. Ethnol.* 3, 572–594.
- Rull, V., 1998. Biogeographical and evolutionary considerations of *Mauritia* (Arecaceae), based on palynological evidence. *Rev. Palaeobot. Palynol.* 100, 109–122.

- Rull, V., 2001. A morphometric study of Early Miocene *Mauritiidites* from northern South America: palaeoecological and evolutionary implications. *Grana* 40, 163–167.
- Rull, V., 2003. Contribution of quantitative ecological methods to the interpretation of stratigraphically homogeneous pre-Quaternary sediments: A palynological example from the Oligocene of Venezuela. *Palynology* 27, 75–98.
- Sah, S.C.D., 1974. Palaeogene biostratigraphy of Shillong Plateau, in: Surange, K.R., Lakhanpal, R.N., Bharadwaj, D.C. (Eds.), *Aspects and Appraisal of Indian Palaeobotany*. Birbal Sahni Institute of Palaeobotany, Lucknow, India, pp. 525–533.
- Sah, S.C.D., Datta, S.K., 1966. Palyno-stratigraphy of the sedimentary formations of Assam. Stratigraphical position of the Cherra formation. *Palaeobotanist* 15, 72–86.
- Salamanca, S., van Soelen, E.E., Teunissen van Manen, M.L., Flantua, S.G.A., Santos, R.V., Roddaz, M., Dantaz, E.L., van Loon, E., Damsté, J.S.S., Kim, J.-H., Hoorn, C., 2016. Amazon forest dynamics under changing abiotic conditions in the early Miocene (Colombian Amazonia). *J. Biogeogr.* 43, 2424–2437.
- Salard-Cheboldaeff, M., 1978. Sur la palynoflore Maestrichtienne et Tertiaire du Bassin sédimentaire Littoral du Cameroun. *Pollen et Spores* 20, 215–260.
- Salard-Cheboldaeff, M., 1979. Palynologie maestrichtienne et tertiaire du cameroun. Etude qualitative et repartition verticale des principales especes. *Rev. Palaeobot. Palynol.* 28, 365–388.
- Salard-Cheboldaeff, M., 1990. Intertropical African palynostratigraphy from Cretaceous to late quaternary times. *J. Afr. Earth Sci. (Middle East)* 11, 1–24.
- Salgado-Labouriau, M.L., 1997. Late Quaternary paleoclimate in the savannas of South America. *J. Quat. Sci.*, 12, 371–379.
- Samant, B., 2000. Palynostratigraphy and age of the Bhavnagar lignite, Gujarat, India. *Palaeobotanist* 49, 101–118.
- Samant, B., Mohabey, D.M., 2009. Palynoflora from Deccan volcano-sedimentary sequence (Cretaceous–Palaeogene transition) of central India: implications for spatio-temporal correlation. *J. Biosci.* 34, 811–823.
- Samant, B., Phadtre, N.R., 1997. Stratigraphic palynoflora of the early Eocene Rajpardi lignite, Gujarat and the lower age limit of the Tarkeshwar Formation of South Cambay Basin, India. *Palaeontogr. Abt. B* 245, 1–108.
- Samant, B., Tapaswi, P.M., 2001. Palynology of the Early Eocene Surat lignite deposits of Gujarat, India. *J. Palaeontol. Soc. India* 46, 121–132.

- Sander, N.L., Pérez-Zaval, F., Da Silva, C.J., Arruda, J.C., Pulido, M.T., Barelli, M.A.A., Rosi, A.B., Viana, A.P., Boechat, M.S.B., Bacon, C.D., Cibrián-Jaramillo, A., 2018. Rivers shape population genetic structure in *Mauritia flexuosa* (Arecaceae). *Ecol. Evol.* 8, 6589–6598.
- Sarmiento, G., 1991. Palinología de la Formación Guaduas—estratigrafía y sistemática. *Bol. Geol. INGEOMINAS* 32, 45–126.
- Saxena, R.K., Khare, S., 2004. Palynological investigation of the Jayamkondacholapuram Well 12, Tiruchirapalli District, Tamil Nadu, India. *Geophytology* 34, 73–93.
- Saxena, R.K., Trivedi, G.K., 2006. A catalogue of Tertiary spores and pollen from India. Diamond Jubilee Special Publication, Birbal Sahni Institute of Palaeobotany, Lucknow, India.
- Saxena, R.K., Trivedi, G.K., 2009. Palynological investigation of the Kopili Formation (Late Eocene) in North Cachar Hills, Assam, India. *Acta Palaeobot.* 49, 253–277.
- Schneider, H., Schmidt, A.R., Heinrichs, J., 2016. Burmese amber fossils bridge the gap in the Cretaceous record of polypod ferns. *Perspect. Plant Ecol. Evol. Syst.* 18, 70–80.
- Schrank, E., 1987. Biostratigraphic importance of microfloras from the Late Cretaceous Clastic Series of Northwestern Sudan. *Cretac. Res.* 8, 29–42.
- Schrank, E., 1994. Palynology of the Yesomma Formation in the northern Somalia: A study of pollen, spores and associated phytoplankton from the Late Cretaceous Palmae Province. *Palaeontogr. Abt. B* 231, 63–112.
- Selling, O.H., 1944. Studies in the Recent and fossil species of *Schizaea*, with particular reference to their spore characters. *Meddel. Goteborgs Bot. Tradg.* 16, 1–112.
- Sevastjanova, I., Hall, R., Rittner, M., Saw Mu Tha Lay Paw, Tin Tin Naing, Alderton, D.H., Comfort, G., 2016. Myanmar and Asia united, Australia left behind long ago. *Gondwana Res.* 32, 24–40.
- Shaffer, B.L., 1987. The potential of calcareous nannofossils for recognizing Plio–Pleistocene climatic cycles and sequence boundaries on the shelf, in: *Proceedings, Gulf Coast Section SEPM 8th Annual Research Conference*, pp. 142–145.
- Shi, G., Jacques, F.M., Li, H., 2014. Winged fruits of *Shorea* (Dipterocarpaceae) from the Miocene of Southeast China: Evidence for the northward extension of dipterocarps during the mid-Miocene climatic optimum. *Rev. Palaeobot. Palynol.* 200, 97–107.

- Shiono, T., Kusumoto, B., Yasuhara, M., Kubota, Y., 2018. Roles of climate niche conservation and range dynamics in woody plant diversity patterns through the Cenozoic. *Glob. Ecol. Biogeogr.* 27, 865–874.
- Shukla, A., Mehrotra, R.C., Spicer, R.A., Spicer, T.E.V., Kumar, M., 2014. Cool equatorial terrestrial temperatures and the South Asian monsoon in the Early Eocene: evidence from the Gurha Mine, Rajasthan, India. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 412, 187–198.
- Silveira, R.R., Souza, P.A.D., 2015. Palinologia (grãos de pólen de angiospermas) das formações Solimões e Iça (bacia do Solimões), nas regiões de Coari e Alto Solimões, Amazonas. *Rev. Bras. Paleontol.* 18, 455–474.
- Singh, H.P., Singh, R.Y., Sah, S.C.D., 1979. Further observations on the genus *Dandotiaspora*, Sah, Kar & Singh, 1971. *Palaeobotanist* 26, 185–189.
- Singh, R.S., 1990. Palynology of Langrin Coalfield, South Shillong Plateau, Meghalaya. *Palaeobotanist* 38, 217–228.
- Smedmark, J.E., Anderberg, A.A., 2007. Boreotropical migration explains hybridization between geographically distant lineages in the pantropical clade Sideroxyleae (Sapotaceae). *Am. J. Bot.* 94, 1491–1505.
- Smith, B., Wilson, J.B., 1996. A consumer's guide to evenness indices. *Oikos* 76, 70–82.
- Soares, E.A.A., D'Apolito, C., Jaramillo, C., Harrington, G., Caputo, M.V., Barbosa, R. O., dos Santos, E.B., Dino, R., Gonçalves, A.D., 2017. Sedimentology and palynostratigraphy of a Pliocene-Pleistocene (Piacenzian to Gelasian) deposit in the lower Negro River: Implications for the establishment of large rivers in Central Amazonia. *J. S. Am. Earth Sci.* 79, 215–229.
- Soares, E.A.A., Dino, R., Soares, D.P., Antonioli, L., da Silva, M.A.L., 2015. New sedimentological and palynological data from surface Miocene strata in the central Amazonas Basin area. *Braz. J. Geol.* 45, 337–357.
- Soe Moe Lwin, Si Si Thu, Myo Min Aung, Kyaw Kyaw Khine, Hayman Aung, Moe Thidar, Suang Khan Pau, 2017. Paleo-environment of the Miocene sediments, central belt, Myanmar, in: *Papers of the First Myanmar National Conference on Earth Sciences (MNCES, 2017)*. University of Monywa, Monywa, Myanmar, pp. 348–359.
- Song, Y.G., Fragnière, Y., Meng, H.H., Li, Y., Bétrisey, S., Corrales, A., Manchester, S., Deng, M., Jasińska, A.K., Vãn Sâm, H., Kozłowski, G., 2020. Global biogeographic

- synthesis and priority conservation regions of the relict tree family Juglandaceae. *J. Biogeogr.* 47, 643–657.
- Song, Z., Wang, W., Huang, F., 2004. Fossil pollen records of extant angiosperms in China. *Bot. Rev.* 70, 425–458.
- Songtham, W., Ratanasthien, B., Mildenhall, D.C., Singharajwarapana, S., Kandharosa W., 2003. Oligocene–Miocene Climatic Changes in Northern Thailand Resulting from Extrusion Tectonics of Southeast Asian Land mass. *Sci. Asia* 29, 221–233.
- Sorrel, P., Eymard, I., Leloup, P.H., Maheo, G., Olivier, N., Sterb, M., Gourbet, L., Wang, G.C., Jing, W., Lu, H.J., Li, H.B., Xu, Y.D., Zhang, K.X., Cao, K., Chevalier, M.L., Replumaz, A., 2017. Wet tropical climate in SE Tibet during the Late Eocene. *Sci. Rep.* 7, 7809.
- Sowunmi, M.A., 1981. Aspects of late Quaternary vegetational changes in West Africa. *J. Biogeogr.* 8, 457–474.
- Spicer, R.A., 2017. Tibet, the Himalaya, Asian monsoons and biodiversity – In what ways are they related? *Plant Divers.* 39, 233–244.
- Spicer, R.A., Yang, J., Herman, A.B., Kodrul, T.M., Aleksandrova, G.N., Maslova, N.P., Spicer, T.E.V., Ding, L., Xu, Q., Shukla, A., Srivastava, G., Mehrotra, R., Liu, X.Y., Jin, J.H., 2017. Paleogene monsoons across India and South China: Drivers of biotic change. *Gondwana Res.* 49, 350–363.
- Spicer, R.A., Yang, J., Herman, A.B., Kodrul, T., Maslova, N., Spicer, T.E.V., Aleksandrova, G., Jin J., 2016. Asian Eocene monsoons as revealed by leaf architectural signatures. *Earth Planet. Sci. Lett.* 449, 61–68.
- Srivastava, J., Prasad, V., 2015. Effect of global warming on diversity pattern in *Nypa* mangroves across Paleocene–Eocene transition in the paleo-equatorial region of the Indian sub-continent. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 429, 1–12.
- Srivastava, S.K., 1972. Some spores and pollen from the Paleocene Oak Hill Member of the Naheola Formation, Alabama (U.S.A.). *Rev. Palaeobot. Palynol.* 14, 217–285.
- Srivastava, S.K., Binda, P.L., 1991. Depositional history of the Early Eocene Shumaysi Formation, Saudi Arabia. *Palynology* 15, 47–61.
- Srivastava, S.K., Braman, D.R., 2013. The palynostratigraphy of the Edmonton Group (Upper Cretaceous) of Alberta, Canada. *Palynology* 37, 127.
- Stamp, D.L., 1925. *The Vegetation of Burma from an Ecological Standpoint*. University of Rangoon, Rangoon, Myanmar.

- Stein, A., Gerstner, K., Kreft, H., Arita, H., 2014. Environmental heterogeneity as a universal driver of species richness across taxa, biomes and spatial scales. *Ecol. Lett.* 17, 866–880.
- Stockey, R.A., Hoffman, G.L., Rothwell, G.W., 1997. The fossil monocot *Limnobiophyllum scutatatum*: Resolving the phylogeny of Lemnaceae. *Am. J. Bot.* 84, 355–368.
- Storch, D., Bohdalková, E., Okie, J., 2018. The more-individuals hypothesis revisited: the role of community abundance in species richness regulation and the productivity-diversity relationship. *Ecol. Lett.* 21, 920–937.
- Stover, L.E., Evans, P.R., 1973. Upper Cretaceous–Eocene spore-pollen zonation, offshore Gippsland Basin, Australia, in: Glover, J.E., Playford, G. (Eds.), *Mesozoic and Cainozoic Palynology: Essays in Honour of Isabel Cookson*. Geological Society of Australia Special Publication 4, pp. 55–72.
- Stover, L.E., Partridge, A.D., 1973. Tertiary and Late Cretaceous spores and pollen from the Gippsland Basin, southeastern Australia. *Proc. R. Soc. Victoria* 85, 237–286.
- Su, T., Spicer, R.A., Wu, F., Farnsworth, A., Huang, J., Del Rio, C., Deng, T., Ding, L., Huang, Y., Hughes, A., Jia, L., Jin, J., Li, S., Liang, S., Liu, J., Liu, X., Sherlock, S., Spicer, T., Srivastava, G., Tang, H., Valdes, P., Wang, T., Widdowson, M., Wu, M., Xing, Y., Xu, C., Yang, J., Zhang, C., Zhang, S., Zhang, X., Zhao, F., Zhou, Z., 2020. A Middle Eocene lowland humid subtropical “Shangri-La” ecosystem in central Tibet. *Proc. Natl. Acad. Sci. U.S.A.* 117, 32989–32995.
- Sun, J., Windley, B.F., 2015. Onset of aridification by 34 Ma across the Eocene-Oligocene transition in Central Asia. *Geology* 43, 1015–1018.
- Sun, X., Li, M., Zhang, Y., Lei, Z., Kong, Z., Li, P., Ou, Q., Liu, Q., 1981. Pollen and Spores, in: Hou, Y. (Ed.), *Tertiary Paleontology of North Continental Shelf of South China Sea*. Guangdong Science and Technology Press, Guangzhou, China.
- Sun, X., Wang, P., 2005. How old is the Asian monsoon system? – Palaeobotanical records from China. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 222, 181–222.
- Svenning, J.-C., Normand, S., Kageyama, M., 2008. Glacial refugia of temperate trees in Europe: insights from species distribution modelling. *J. Ecol.* 96, 1117–1127.
- Swenson, U., Anderberg, A.A., 2005. Phylogeny, character evolution, and classification of Sapotaceae (Ericales). *Cladistics* 21, 101–130.

- Takahashi, K., 1982. Miospores from the Eocene Nanggulan Formation in the Yogyakarta Region, Central Java. *Trans. Proc. Palaeontol. Soc. Jpn. N. Ser.* 126, 303–326.
- Takai, M., Nishioka, Y., Thaug-Htike, Maung Maung, Kyaw Khaing, Zin-Maung-Maung-Thein, Tsubamoto, T., Egi, N., 2016. Late Pliocene *Semnopithecus* fossils from central Myanmar: rethinking of the evolutionary history of cercopithecoid monkeys in Southeast Asia. *Hist. Biol.* 28, 171–187.
- Tang, H., Li, S., Su, T., Spicer, R.A., Zhang, S., Li, S., Liu, J., Lauretano, V., Witkowski, C., Spicer, T.E.V., Deng, W., Wu, M., Ding, W., Zhou, Z., 2020. Early Oligocene vegetation and climate of southwestern China inferred from palynology. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 560, 109988.
- Tardif, D., Fluteau, F., Donnadieu, Y., Le Hir, G., Ladant, J.-B., Sepulchre, P., Licht, A., Poblete, F., Dupont-Nivet, G., 2020. The onset of Asian monsoons: A modelling perspective. *Clim. Past* 16, 847–865.
- Teodoridis, V., Kovar-Eder, J., Marek, P., Kvacek, Z., Mazouch, P., 2011. The integrated plant record vegetation analysis: Internet platform and online application. *Acta Mus. Natl. Pragae* 67, 159–165.
- ter Steege, H., Pitman, N.C.A., Sabatier, D., Baraloto, C., Salomao, R.P., Guevara, J.E., Phillips, O.L., Castilho, C.V., Magnusson, W.E., Molino, J.-F.; Monteagudo, A., Nunez Vargas, P., Carlos Montero, J., Feldpausch, T.R., Coronado, E.N.H., Killeen, T.J., Mostacedo, B., Vasquez, R., Assis, R.L., Terborgh, J., Wittmann, F., Andrade, A., Laurance, W.F., Laurance, S.G.W., Marimon, B.S., Marimon, B.-H., Guimaraes Vieira, I.C., Amaral, I.L., Brienen, R., Castellanos, H., Cardenas Lopez, D., Duivenvoorden, J.F., Mogollon, H.F., de Almeida Matos, F.D., Davila, N., Garcia-Villacorta, R., Stevenson Diaz, P.R., Costa, F., Emilio, T., Levis, C., Schietti, J., Souza, P., Alonso, A., Dallmeier, F., Duque Montoya, A.J., Fernandez Piedade, M.T., Araujo-Murakami, A., Arroyo, L., Gribel, R., Fine, P.V.A., Peres, C.A., Toledo, M., Gerardo, A.A.C., Baker, T.R., Ceron, C., Engel, J., Henkel, T.W., Maas, P., Petronelli, P., Stropp, J., Eugene Zartman, C., Daly, D., Neill, D., Silveira, M., Rios Paredes, M., Chave, J., de Andrade Lima, D., Jorgensen, P.M., Fuentes, A., Schoengart, J., Cornejo Valverde, F., Di Fiore, A., Jimenez, E.M., Penuela Mora, M.C., Fernando Phillips, J., Rivas, G., van Andel, T.R., von Hildebrand, P., Hoffman, B., Zent, E.L., Malhi, Y., Prieto, A., Rudas, A., Ruschell, A.R., Silva, N., Vos, V., Zent, S., Oliveira, A.A., Cano Schutz, A., Gonzales, T., Nascimento, M.T., Ramirez-Angulo, H., Sierra, R., Tirado,

- M., Umana Medina, M.N., van der Heijden, G., Vela, C.I.A., Vilanova Torre, E., Vriesendorp, C., Wang, O., Young, K.R., Baider, C., Balslev, H., Ferreira, C., Mesones, I., Torres-Lezama, A., Urrego Giraldo, L.E., Zagt, R., Alexiades, M.N., Hernandez, L., Huamantupa-Chuquimaco, I., Milliken, W., Palacios Cuenca, W., Pauletto, D., Valderrama Sandoval, E., Valenzuela Gamarra, L., Dexter, K.G., Feeley, K., 2013. Hyperdominance in the Amazonian tree flora. *Science* 342, 1243092.
- Thanikaimoni, G., Caratini, C., Venkatachala, B.S., Ramanujam, C.G.K., Kar, R.K., 1984. Selected Tertiary angiosperm pollens from India and their relationship with African Tertiary pollens. *Travaux Sect. Sci. Technol. IFP* 19, 1–93.
- Thompson, R., Anderson, K., Pellier, R., Strickland, L., Bartlein, P., Shafer, S., 2012. Quantitative estimation of climatic parameters from vegetation data in North America by the mutual climatic range technique. *Quat. Sci. Rev.* 51, 18–39.
- Thornthwaite, C.W., 1948. An approach toward a rational classification of climate, *Geogr. Rev.* 38, 55–94.
- Tomlinson, P.B., 1986. *The Botany of Mangroves*. Cambridge University Press, Cambridge, UK.
- Tong, G., Zheng, M., Wang, W., Yuan, H., Liu, J., Shen, Z., 2001. Eocene palynological assemblages and environment in the Baise Basin of Guangxi. *J. Stratigr.* 25, 273–278.
- Tripathi, S.K.M., Kumar, M., Srivastava, D., 2009. Palynology of Lower Palaeogene (Thanetian-Ypresian) coastal deposits from the Barmer Basin (Akli Formation, Western Rajasthan, India): Palaeoenvironmental and palaeoclimatic implications. *Geol. Acta* 7, 147–160.
- Tripathi, S.K.M., Saxena, R.K., Prasad, V., 2000. Palynological investigation of the Tura Formation (Early Eocene) exposed along Tura-Dalu Road, West Garo Hills, Meghalaya, India. *Palaeobotanist* 49, 239–251.
- Tripathi, S.K.M., Srivastava, D., 2010. Palynological investigation, facies analysis and palaeoenvironmental interpretations from late Palaeocene to early Eocene lignites and associated sediments of Barmer, western India. *Palaeobotanist* 59, 1–32.
- Tripathi, S.K.M., Srivastava, D., 2012. Palynology and palynofacies of the early Palaeogene lignite bearing succession of Vastan, Cambay Basin, Western India. *Acta Palaeobot.* 52, 157–175.

- Trivedi, G.K., Ranhotra, P.S., 2015. Palynofloral evidence for palaeoecology and depositional environment of the Kopili Formation (Late Eocene), Jaintia Hills, Meghalaya. *J. Geol. Soc. India* 86, 33–40.
- Trivedi, G.K., Saxena, R.K., 2000. Palynofloral investigation of the Kopili Formation (Late Eocene) exposed near Umrongso in North Cachar Hills District, Assam, India. *Palaeobotanist* 49, 269–280.
- Tschudy, R.H., 1973. Stratigraphic distribution of significant Eocene palynomorphs of the Mississippi Embayment. *U.S. Geol. Surv. Prof. Pap.* 743-B, p. 24.
- Umeji, O.P., 2003. Palynological data from the road section at the Ogbunike Tollgate, Onitsha, southeastern Nigeria. *J. Min. Geol.* 39, 95–102.
- Urrego, G., Galeano, L. E., Sánchez, S., Peñuela, C., 2013. Paleoecología, ecología y etnobotánica de los cananguchales de la Amazonia Colombiana, in: C. A. Lasso, A. Rial, V. González-B (Eds.), *Morichales y Cananguchales de la Orinoquia y la Amazonia: Colombia y Venezuela* (vol. 1). Instituto de Investigación de Recursos Biológicos Alexander von Humboldt (IAvH), Bogotá D.C., Colombia, pp. 217–246.
- Vajda, V., Bercovici, A., 2014. The global vegetation pattern across the Cretaceous–Paleogene mass extinction interval: A template for other extinction events. *Glob. Planet. Change* 122, 29–49.
- Vajda-Santivaner, V., 1999. Miospores from upper Cretaceous–Paleocene strata in northwestern Bolivia. *Palynology* 23, 181–196.
- Valdes, P.J., Armstrong, E., Badger, M.P.S., Bradshaw, C.D., Bragg, F., Crucifix, M., Davies-Barnard, T., Day, J.J., Farnsworth, A., Gordon, C., Hopcroft, P.O., Kennedy, A.T., Lord, N.S., Lunt, D.J., Marzocchi, A., Parry, L.M., Pope, V., Roberts, W.H.G., Stone, E.J., Tourte, G.J.L., Williams, J.H.T., 2017. The BRIDGE HadCM3 family of climate models: HadCM3@bristol v1.0. *Geosci. Model Dev.* 10, 3715–3743.
- Valdiya, K.S., 2002. Emergence and evolution of Himalaya: reconstructing history in the light of recent studies. *Prog. Phys. Geogr.* 26, 360–399.
- van Dam, J.A., 2006. Geographic and temporal patterns in the late Neogene (12–3 Ma) aridification of Europe: the use of small mammals as paleoprecipitation proxies. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 238, 90–218.
- van der Hammen, T., 1956. Description of some genera and species of fossil pollen and spores. *Bol. Geol. (Bogotá)* 4, 111–117.

- van der Hammen, T., 1957. Climatic periodicity and evolution of South American Maestrichtian and Tertiary floras. *Bol. Geol. (Bogotá)* 5, 49–91.
- van der Hammen, T., Absy, M.L., 1994. Amazonia during the last glacial. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 109, 247–261.
- van der Hammen, T., Garcia de Mutis, C., 1966. The Paleocene pollen flora of Colombia. *Leidse Geol. Meded.* 35, 105–114.
- van Gorsel, J.T., Lunt, P., Morley, R.J., 2014. Introduction to Cenozoic biostratigraphy of Indonesia-SE Asia. *Berita Sedimentol.* 29, 6–40.
- van Hoeken-Klinkenberg, P.M.J., 1964. A palynological investigation of some Upper Cretaceous sediments in Nigeria. *Pollen et Spores* 6, 209–231.
- Van Sickel, W.A., Kominz, M.A., Miller, K.G., Browning, J.V., 2004. Late Cretaceous and Cenozoic sea-level estimates: backstripping analysis of borehole data, onshore New Jersey. *Basin Res.* 16, 451–465,
- van Steenis, C.G.G.J., 1962. The land-bridge theory in botany. *Blumea* 11, 235–372.
- van Steenis, C.G.G.J., 1963. Transpacific floristic affinities particularly in the tropical zone, in: Gressitt, J.K. (Ed.), *Pacific Basin Biogeography*. Bishop Museum Press, Hawaii, USA, pp. 219–31.
- van Steenis, C.G.G.J., Schippers-Lammertse, A.F., 1965. Concise plant-geography of Java, in: Backer, C.A., Backhuizen van den Brink, R.C. (Eds.), *Flora of Java*, vol. 2. N.V.P. Noordhoff, Groningen, The Netherlands, pp. 1–72.
- Venkatachala, B.S., 1974. Palynological zonation of the Mesozoic and Tertiary subsurface sediments in the Cauvery basin, in: Surange, K.R., Lakhanpal, R.N., Bharadwaj, D.C. (Eds.), *Aspects and Appraisal of Indian Palaeobotany*. Birbal Sahni Institute of Paleobotany, Lucknow, India, pp. 476–494.
- Venkatachala, B.S., Kar, R.K., Rajanikanth, A., Ly, A., 1998. Palynology of the Cretaceous Tertiary transition in an Atlantic bore core, off Senegal Coast, West Africa. *Palaeobotanist* 47, 60–67.
- Venkatachala, B.S., Rawat, M.S., 1972. Palynology of the Tertiary sediments in the Cauvery Basin. I. Palaeocene–Eocene palynoflora from the subsurface, in: Ghosh, A.K. (Ed.), *Seminar in Paleopalynology and Indian Stratigraphy*, 1971. Botany Department, Calcutta University, Calcutta, India, pp. 292–335.

REFERENCES

- Venkatachala, B.S., Rawat, M.S., 1973. Palynology of the Tertiary sediments in the Cauvery Basin 2. Oligocene–Miocene palynoflora from the surface. *Palaeobotanist* 20, 238–263.
- Venkatachala, B.S., Sharma, K.D., 1984. Palynological zonation in subsurface sediments in Narsapur Well No. 1, Godavari-Krishna Basin, India, in: Badve, R.M. (Ed.), *Proceedings of the 10th Indian Colloquium on the Indian Micropalaeontology and Stratigraphy, 1982*. Maharashtra Association for the Cultivation of Science, Pune, India, pp. 445–466.
- Vergara, S.L.E., Rodriguez, C.H.G., 1997. The Upper Cretaceous and lower Paleocene of the Eastern Bogota Plateau and Llanos Thrustbelt, Colombia: Alternative appraisal to the nomenclature and sequence stratigraphy. *Geol. Colombiana* 22, 51–79.
- Verma, P., Garg, R., Rao, M.R., Bajpai, S., 2020. Palynofloral diversity and palaeoenvironments of early Eocene Akri lignite succession, Kutch Basin, western India. *Paleobiodivers. Paleoenviron.* 100, 605–627.
- Vidal, G., 1988. A palynological preparation method. *Palynology* 12, 215–220.
- Vimal, K.P., 1952. Spores and pollen from Tertiary lignites from Dandot, West Punjab (Pakistan). *India Acad. Sci. Proc.* 36, 135–147.
- Vinken, R., 1988. The Northwest European Tertiary Basin. *Geol. Jahrb. Reihe A* 100, 1–508.
- Walker, J.D., Geissman, J.W., Bowring, S.A., Babcock, L.E., compilers., 2018. *Geologic Time Scale v. 5.0*. Geological Society of America.
- Wang, B., Ding, Q., 2008. Global monsoon: Dominant mode of annual variation in the tropics. *Dyn. Atmospheres Oceans* 44, 165–183.
- Wang, B., Ho, L., 2002. Rainy seasons of the Asian-Pacific monsoon. *J. Clim.* 15, 386–398.
- Wang, J.-G., Wu, F.-Y., Tan, X.-C., Liu, C.-Z., 2014. Magmatic evolution of the Western Myanmar Arc documented by U–Pb and Hf isotopes in detrital zircon. *Tectonophysics* 612–613, 97–105.
- Wang, X., Kraatz, B., Meng, J., Carrapa, B., DeCelles, P., Clementz, M., Abdulov, S., Chen, F., 2016. Central Asian aridification during the late Eocene to early Miocene inferred from preliminary study of shallow marine-eolian sedimentary rocks from northeastern Tajik Basin. *Sci. China Earth Sci.* 59, 1242–1257.

- Warny, S., Jarzen, D.M., Evans, A., Hesp, P., Bart, P., 2012. Environmental significance of abundant and diverse hornwort spores in a potential submerged Paleindian site in the Gulf of Mexico. *Palynology* 36, 234253.
- Warny, S., Jarzen, D.M., Haynes, S.J., MacLeod, K.G., Huber, B.T., 2018. Late Cretaceous (Turonian) angiosperm pollen from Tanzania: a glimpse of past vegetation from a warmer climate. *Palynology* 43, 608–620.
- Wasiljeff, J., Kaakinen, A., Salminen, J.M., Zhang, Z.Q., 2020. Magnetostratigraphic constraints on the fossiliferous Ulanatal sequence in Inner Mongolia, China: Implications for Asian aridification and faunal turnover before the Eocene-Oligocene boundary. *Earth Planet. Sci. Lett.* 535, 1–15.
- Watanasak, M., 1988. Mid-Tertiary palynology of onshore and offshore Thailand. Ph.D. Thesis, University of Adelaide, Adelaide, Australia.
- Watanasak, M., 1990. Mid tertiary palynostratigraphy of Thailand. *J. Southeast Asian Earth Sci.* 4, 203–218.
- Wei, L.J., Liu, X.H., Yan, F.H., Mai, X.S., Li, G.W., Liu, X.B., Zhou, X.J., 2011. Palynological evidence sheds new light on the age of the Liuqu Conglomerates in Tibet and its geological significance. *Sci. China Earth Sci.* 54, 901–911.
- Werner, W.L., 1997. Pines and other conifers in Thailand—a Quaternary relic? *J. Quatern. Res.* 12, 451–454.
- Westerhold, T., Marwan, N., Drury, A.J., Liebrand, D., Angini, C., Anagnostou, E., Barnet, J.S., Bohaty, S.M., Vleeschouwer, D.D., Florindo, F., Frederichs, T., Hodell, D.A., Holbourn, A.E., Kroon, D., Lauretano, V., Littler, K., Lourens, L.J., Lyle, M., Pälike, H., Röhl, U., Tian, J., Wilkens, R.H., Wilson, P.A., Zachos, J. C., 2020. An astronomically dated record of Earth's climate and its predictability over the last 66 million years. *Science* 369, 1383–1387.
- Westerweel, J., Licht, A., Dupont-Nivet, G., Roperch, P., Myat Kay Thi, Hnin Hnin Swe, Huang, H., Zaw Win, Day Wa Aung, 2020. Burma Terrane collision and northward indentation in the Eastern Himalayas recorded in the Eocene–Miocene Chindwin Basin (Myanmar). *Tectonics* 39, e2020TC006413.
- Westerweel, J., Roperch, P., Licht, A., Dupont-Nivet, G., Zaw Win, Poblete, F., Ruffet, G., Hin Hin Swe, Myat Kai Thi, Day Wa Aung, 2019. Burma Terrane part of the Trans-Tethyan arc during collision with India according to palaeomagnetic data. *Nat. Geosci.* 12, 863–868.

- White, N.M., Pringle, M., Garzanti, E., Bickle, M., Najman, Y., Chapman, H., Friend, P., 2002. Constraints on the exhumation and erosion of the High Himalayan Slab, NW India, from foreland basin deposits. *Earth Planet. Sci. Lett.* 195, 29–44.
- Whitmore, T.C., 1973. *Palms of Malaysia*. Oxford University Press, London, UK.
- Wijmstra, T.A., 1969. Palynology of the Alliance Well. *Neth. J. Geosci.* 48, 125–133.
- Willis, K.J., McElwain, J., 2014. *The Evolution of Plants*, 2nd Edition. Oxford University Press, Oxford, UK.
- Winantris, W., Hamdani, H., Harlia, E., 2017. Paleoenvironment of Tanjung Formation Barito Basin Central Kalimantan Based on palynological data. *J. Geosci. Eng. Environ. Technol.* 2, 110–116.
- Wing, S., Greenwood, D.R., 1993. Fossils and fossil climate: the case for equable continental interiors in the Eocene. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* 341, 243–252.
- Witts, D., Hall, R., Nichols, G., Morley, R.J., 2012. A new depositional and provenance model for the Tanjung Formation, Barito Basin, SE Kalimantan, Indonesia. *J. Asian Earth Sci.* 56, 77–104.
- Wu, J., Zhang, K., Xu, Y., Wang, G., Garziane, C.N., Leloup, P.H., Sorrel, P., Mahéo, G., 2018. Paleoelevations in the Jianchuan Basin of the southeastern Tibetan Plateau based on stable isotope and pollen grain analyses. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 53, 93–108.
- Wysocka, A., Pha, P.D., Durska, E., Czarniecka, U., Thang, D.V., Filipek, A., Cuong, N.Q., Tuan, D.M., Huyen, N.X., Tha, H.V., Staniszewski, R., 2020. The Na Duong Basin (North Vietnam): A key for understanding Paleogene basin evolution in relation to the left-lateral Cao Bang-Tien Yen Fault. *J. Asian Earth Sci.* 195, 104350.
- Xia, K., Su, T., Liu, Y.S., Xing, Y.W., Jacques, F.M.B., Zhou, Z.K., 2009. Quantitative climate reconstructions of the late Miocene Xiaolongtan megaflora from Yunnan, southwest China. *Palaeogeogr. Palaeoclimat. Palaeoecol.* 276, 80–86.
- Xiao, G.Q., Abels, H.A., Yao, Z.Q., Dupont-Nivet, G., Hilgen, F.J., 2010. Asian aridification linked to the first step of the Eocene-Oligocene climate transition (EOT) in obliquity-dominated terrestrial records (Xining Basin, China). *Clim. Past* 6, 501–513.
- Xie, S.P., Xu, H., Saji, N.H., Wang, Y., Liu, W.T., 2006. Role of narrow mountains in large-scale organization of Asian monsoon convection. *J. Clim.* 19, 3420–3429.

- Xie, Y., Wu, F., Fang, X., 2019. Middle Eocene East Asian monsoon prevalence over southern China: Evidence from palynological records. *Glob. Planet. Change* 175, 13–26.
- Xie, Y., Wu, F., Fang, X., Zhang, D., Zhang, W., 2020. Early Eocene southern China dominated by desert: Evidence from a palynological record of the Hengyang Basin, Hunan Province. *Glob. Planet. Change* 195, 103320.
- Xing, L., Gu, L., 2020. The possible earliest epizoochorous fruit preserved in mid-Cretaceous Burmese amber. *Cretac. Res.* 114, 104498.
- Xing, Y., Gandolfo, M.A., Onstein, R.E., Cantrill, D.J., Jacobs, B.F., Jordan, G.J., Lee, D.E., Popova, S., Srivastava, R., Su, T., Vikulin, S.V., Yabe, A., Linder, H.P., 2016. Testing the biases in the rich Cenozoic angiosperm macrofossil record. *Int. J. Plant Sci.* 177, 371–388.
- Xing, Y.W., Utescher, T., Jacques, F.M.B., Su, T., Liu, Y.S., Huang, Y.J., Zhou, Z.K., 2012. Paleoclimatic estimation reveals a weak winter monsoon in southwestern China during the late Miocene: evidence from plant macrofossils. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 358–360, 19–26.
- Yakzan, A.M., Harun, A., Nasib, B.M., Morley, R.J., 1996. Integrated biostratigraphic zonation for the Malay Basin. *Bull. Geol. Soc. Malays.* 39, 157–184.
- Yang, B., Deng, M., Zhang, M., Aung Zaw Moe, Ding, H., Mya Bhone Maw, Pyae Pyae Win, Corlett, R.T., Tan, Y., 2020. Contributions to the flora of Myanmar from 2000 to 2019. *Plant Divers.* 42, 292–301.
- Yao, W., Ding, L., Cai, F., Wang, H., Xu, Q., Than Zaw, 2017. Origin and tectonic evolution of Upper Triassic turbidites in the Indo-Burman ranges, West Myanmar. *Tectonophysics* 721, 90–105.
- Yao, Y., Bera, S., Ferguson, D.K., Mosbrugger, V., Paudyal, K.N., Jin, J., Li, C., 2009. Reconstruction of paleovegetation and paleoclimate in the Early and Middle Eocene, Hainan Island, China. *Clim. Change* 92, 169–189.
- Yap, S.K., 1989. Sapindaceae, in: Ng, F.S.P. (Ed.), *Tree Flora of Malaya*, vol. 4. Longman, Kuala Lumpur, Malaysia, pp. 434–461.
- Yuan, Q., Barbolini, N., Rydin, C., Gao, D., Wei, H., Fan, Q., Qin, Z., Du, Y., Shan, J., Shan, F., Vajda, V., 2020. Aridification signatures from fossil pollen indicate a drying climate in east-central Tibet during the late Eocene. *Clim. Past* 16, 2255–2273.

REFERENCES

- Zachos, J., Pagani, M., Sloan, L., Thomas, E., Billups, K., 2001. Trends, rhythms, and aberrations in global climate 65 Ma to present. *Science* 292, 686–693.
- Zachos, J.C., Dickens, G.R., Zeebe, R.E., 2008. An early Cenozoic perspective on greenhouse warming and carbon-cycle dynamics. *Nature* 451, 279–283.
- Zachos, J.C., Wara, M.W., Bohaty, S., Delaney, M.L., Petrizzo, M.R., Brill, A., Bralower, T.J., Premoli-Silva, I., 2003. A Transient Rise in Tropical Sea Surface Temperature During the Paleocene-Eocene Thermal Maximum. *Science* 302, 1551–1554.
- Zamaloa, M.C., 2000. Palinoflora y ambiente en el Terciario del nordeste de Tierra del Fuego, Argentina. *Rev. Mus. Argent. Cienc. Nat. N. Ser. 2*, 43–51.
- Zamaloa, M.C., Gandolfo, M.A., González, C.C., Romero, E.J., Cúneo, N.R., Wilf, P., 2006. Casuarinaceae from the Eocene of Patagonia, Argentina. *Int. J. Plant Sci.* 167, 1279–1289.
- Zaw Zaw, Fan, Z.-X., Bräuning, A., Xu, C., Liu, W.-J., Gaire, N.P., Panthi, S., Kay Zin Than, 2020. Drought reconstruction over the past two centuries in southern Myanmar using teak tree-rings: linkages to the Pacific and Indian Oceans. *Geophys. Res. Lett.* 47, e2020GL087627.
- Zetter, R., 1989. Methodik und Bedeutung einer routinemäßig kombinierten lichtmikroskopischen und rasterelektronischen Untersuchung fossiler Mikroflora. *Cour. Forschunsinst. Senckenb.* 109, 41–50.
- Zetter, R., Farabee, M.J., Pigg, K.B., Manchester, S.T., DeVore, M., Nowak, M.D., 2011. Palynology of the late Paleocene silicified shale at Almond, North Dakota, USA. *Palynology* 35, 179–211.
- Zetter, R., Ferguson, D.K., 2001. Trapaceae pollen in the Cenozoic. *Acta Palaeobot.* 41, 321–339.
- Zetter, R., Hesse, M., Frosch-Radivo, A., 2001. Early Eocene zona-aperturate pollen grains of the *Proxapertites* type with affinity to Araceae. *Rev. Palaeobot. Palynol.* 117, 267–279.
- Zetter, R., Hofmann, C.-C., 2001. New aspects of the palynoflora of the lowermost Eocene (Krappfeld Area, Carinthia), in: Piller, W.E., Rasser, M.W. (Eds.), *Paleogene of the Eastern Alps*. Verlag der Österreichischen Akademie der Wissenschaften, Wien, Österreich, pp. 473–507.

- Zhang, W., Li, H., Shih, C., Zhang, A., Ren, D., 2018. Phylogenetic analyses with four new Cretaceous bristletails reveal interrelationships of Archaeognatha and Gondwana origin of Meinertellidae. *Cladistics* 34, 384–406.
- Zhao, L., Lu, H., Wang, H., Meadows, M., Ma, C., Tang, L., Lei, F., and Zhang, H., 2020a. Vegetation dynamics in response to evolution of the Asian Monsoon in a warm world: Pollen evidence from the Weihe Basin, central China. *Glob. Planet. Change* 193, 103269.
- Zhao, Y., Tzedakis, P.C., Li, Q., Qin, F., Cui, Q., Liang, C., Zhao, H., Birks, H.J.B., Liu, Y., Zhang, Z., Ge, J., Zhao, H., Felde, V.A., Deng, C., Cai, M., Li, H., Ren, W., Wei, H., Yang, H., Zhang, J., Yu, Z., Guo, Z., 2020b. Evolution of vegetation and climate variability on the Tibetan Plateau over the past 1.74 million years. *Sci. Adv.* 6, eaay6193.
- Zhu, J., Poulsen, C.J., Tierney, J.E., 2019. Simulation of Eocene extreme warmth and high climate sensitivity through cloud feedbacks. *Sci. Adv.* 5, eaax1874.
- Zizka, A., Silvestro, D., Andermann, T., Azevedo, J., Ritter, C.D., Edler, D., Farooq, H., Herdean, A., Ariza, M., Scharn, R., Svantesson, S., Wengström, N., Zizka, V., Antonelli, A., 2019. CoordinateCleaner: Standardized cleaning of occurrence records from biological collection databases. *Methods Ecol. Evol.* 10, 744–751.