UvA-DARE (Digital Academic Repository)

Augmented reality-based remote family visits in nursing homes

Abels, E.A.M.; Toet, A.; Stokking, H.; Klunder, T.; van Berlo, Z.M.C.; Smeets, B.; Niamut, O.

DOI

<https://doi.org/10.1145/3452918.3465502>

Publication date

2021

Document Version

Final published version

Published in

IMX '21: ACM international conference on interactive media experiences

License

CC BY

[Link to publication](#)

Citation for published version (APA):

Abels, E. A. M., Toet, A., Stokking, H., Klunder, T., van Berlo, Z. M. C., Smeets, B., & Niamut, O. (2021). Augmented reality-based remote family visits in nursing homes. In *IMX '21: ACM international conference on interactive media experiences* (pp. 258–263). ACM.
<https://doi.org/10.1145/3452918.3465502>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Augmented Reality-Based Remote Family Visits in Nursing Homes

Eva A.M. Abels
TNO, the Hague, Netherlands
eamabels@gmail.com

Alexander Toet
TNO, the Hague, Netherlands
lex.toet@tno.nl

Hans Stokking
TNO, the Hague, Netherlands
hans.stokking@tno.nl

Tessa Klunder
TNO, the Hague, Netherlands
tessa.klunder@tno.nl

Zeph M.C. van Berlo
University of Amsterdam,
Amsterdam, Netherlands
z.m.c.vanberlo@uva.nl

Bram Smeets
MeanderGroep Zuid-Limburg,
Heerlen, Netherlands
bramsmeets@mgzl.nl

Omar Niamut
TNO, the Hague, Netherlands
omar.niamut@tno.nl

ABSTRACT

During the COVID-19 pandemic, many nursing homes had to restrict visitations. This had a major negative impact on the wellbeing of residents and their family members. In response, residents and family members increasingly resorted to mediated communication to maintain social contact. To facilitate high-quality mediated social contact between residents in nursing homes and remote family members, we developed an augmented reality (AR)-based communication tool. In this study, we compared the user experience (UX) of AR-communication with that of video calling, for 10 pairs of residents and family members. We measured enjoyment, spatial presence and social presence, attitudes, behavior and conversation duration. In the AR-communication condition, residents perceived a 3D projection of their remote family member onto a chair placed in front of them. In the video calling condition, the family member was shown using 2D video. In both conditions, the family member perceived the resident in the video calling mode on a 2D screen. While residents reported no differences in their UX between both conditions, family members reported higher spatial presence for the AR-communication condition compared to video-calling. Conversation durations were significantly longer during AR-communication than during video calling. We tentatively suggest that there may be (unconscious) differences in UX during AR-based communication compared to video calling.

CCS CONCEPTS

• **Human-centered computing** → Human computer interaction (HCI); Interaction paradigms; Mixed / augmented reality.

Permission to make digital or hard copies of part or all of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for third-party components of this work must be honored. For all other uses, contact the owner/author(s).
IMX '21, June 21–23, 2021, Virtual Event, USA
© 2021 Copyright held by the owner/author(s).
ACM ISBN 978-1-4503-8389-9/21/06.
<https://doi.org/10.1145/3452918.3465502>

KEYWORDS

Augmented Reality, Social XR, User Experience, Social Presence, Spatial Presence, Nursing Home

ACM Reference Format:

Eva A.M. Abels, Alexander Toet, Hans Stokking, Tessa Klunder, Zeph M.C. van Berlo, Bram Smeets, and Omar Niamut. 2021. Augmented Reality-Based Remote Family Visits in Nursing Homes. In *ACM International Conference on Interactive Media Experiences (IMX '21)*, June 21–23, 2021, Virtual Event, USA. ACM, New York, NY, USA, 6 pages. <https://doi.org/10.1145/3452918.3465502>

1 INTRODUCTION

Loneliness is one of the most predominant feelings among residents of nursing homes and is associated with negative health outcomes and reduced quality of life [1]. Visits by family members can typically reduce the residents' feelings of loneliness [1]. The importance of these visits became painfully clear during the COVID-19 pandemic, when visitation restrictions resulted in increased feelings of loneliness and social disconnection among residents and their family members [2]. But even before the pandemic, family members often could not visit their relatives living in nursing homes as often as they would like, due to a lack of time and long travel distances [3].

Next to in-person visits, another way for nursing home residents to maintain contact with family and friends is via mediated communication, like phone- or video calling. While this may be a practical way to stay in touch [4], it is often experienced as less fulfilling than face-to-face interactions [3]. A main reason for this is that using current audio-visual communication tools often does not allow for experiencing true feelings of social presence or togetherness, since these tools do not reliably convey essential social and spatial cues [3, 5].

Augmented reality (AR)-based and virtual-reality (VR)-based communication systems may alleviate this problem, by providing a more complete and immersive representation of the social and contextual cues of the communication setting [6]. Thus, AR-based communication may enhance the quality of the mediated contact between residents and their remote family members, which may ultimately positively affect the residents' mental health [7, 8].

The independent research organization TNO (www.tno.nl) recently developed a new AR-based communication tool, based on previous work on social VR [9]. The goal of the current study was to investigate to what extent this tool can facilitate social contact between remote family members and their relatives in nursing homes. First, we conducted focus group sessions with residents, family members, and caretakers, to assess their communication needs and wishes. Based on the outcomes of these sessions, we derived three indicators for the user experience (UX) of mediated (AR or video calling) communication: enjoyment, spatial presence (*'being there'* [10, 11]) and social presence (*'being together with an intelligent other'* [6]). Next, we conducted an experiment with a two-group (AR communication vs video calling) within-subjects design to compare the UX of AR-communication to that of video calling. The sample consisted of 10 participant pairs of residents and family members. In the video calling condition, the resident saw the remote family member projected on a flat (2D) screen. In the AR-communication condition, a 3D representation of the remote family member was projected onto a chair placed in front of the resident (creating the impression that the family member was actually present in the room). In both conditions, the family member saw the resident in the video calling mode on a 2D screen. We hypothesized that the residents would evaluate the AR-communication condition more positively than video calling, while family members would not evaluate both conditions differently.

2 RELATED WORK

The UX of immersive communication systems has been extensively investigated, focusing on the technology's capability to promote the senses of spatial and social presence. Other factors that determine UX of these systems are the naturalness of interaction, conversational engagement, comfortableness of the environment and enjoyment [12]. However, such research has been mainly focused on VR technology, while much less is known about AR-based systems [13]. Since VR and AR affordances are fundamentally different [14], it is unclear whether (social) interaction measures in VR apply to AR. Augmented reality systems supplement the real world with 3D-registered virtual objects [13] and do not replace the user's surroundings as VR systems would do [8]. This leads to different conceptualizations of immersion and spatial presence. For VR, these concepts refer to the medium's capability to deliver an illusion of reality of the virtual world and the user's perception of physically *'being there'* [10, 11]. For AR however, they refer to the plausibility and consistency of the virtual content in the user's own surroundings, such that virtual objects seem part of the physical world [15, 16]. While UX assessment techniques developed for VR can probably be adapted to AR, there is currently no standard evaluation methodology to assess UX of AR-mediated social communication [8]. In this study we therefore used an evaluation methodology based on indicators derived from focus group sessions.

3 METHOD

3.1 Participants and procedure

Ten pairs of participants were included in our study, with each pair consisting of one resident (aged between 80 and 95 years) and one family member (aged between 47 and 68 years). To mitigate

potential novelty effects, the AR system was introduced to the residents before the experiment. Each pair participated in both conditions (i.e., AR and video calling) in counterbalanced order within a single testing day. Family members were informed of the resident's communication condition.

We evaluated the UX of the participants directly after each session. For each participant, we measured their perceived enjoyment, spatial presence and social presence, attitudes, behaviors and conversation duration. The UX of the residents was evaluated in a semi-structured interview and by behavioral observations by a caretaker familiar with the resident, and the UX of the family members was assessed using questionnaires.

For evaluation of the UX indicators of enjoyment, spatial presence and social presence, we asked the residents and family members to grade their experiences on a scale from 0 to 10. Additionally, the family members filled in a Dutch adaptation of the Networked Minds Questionnaire (NMQ) [17]. To investigate attitudes, we asked the residents if they were interested to use the technology again in the future. For the family members, questionnaire items were included about the technology's suitability for repeated use for the residents and for themselves. Caretakers scored the residents' behavior using an adapted Dutch version of the Music in Dementia Assessment Scales (MiDAS) [18], which was accommodated to examine the well-being of a person during remote family visits. We also recorded the duration of each session, with a maximum of 20 minutes. This experimental protocol was approved by the local ethical review board and written informed consent was obtained from all participants.

3.2 Technical setup

The resident was seated in front of a 12.9" iPad Pro tablet, while being recorded by a Logitech webcam that was attached on top of the iPad (Figure 1). The family member was displayed on the iPad, which supported AR placement and rendering of 3D volumetric objects. The iPad was mounted vertically on a stand, positioned in between the resident's chair and an empty chair (Figure 2, left). On the other side, the family member was seated in front of a 43" TV screen on which the resident was displayed (Figure 1 and 2, right), while being registered by a color-plus-depth (RGB-D) camera, the Azure Kinect, that was positioned in front of the TV.

Two computers were used to connect the devices on both sides via the internet (Figure 1). Microsoft Teams was used to deliver the video images from the resident to the family member and to transmit the audio signals in both directions. Audio was recorded and presented using Jabra 750 speakerphones to ensure good audio quality. A dedicated local area network was used for video and audio data transmission, ensuring a high-performance and uninterrupted connection.

In the video calling (Microsoft Teams) condition, the resident was filmed by the iPad front camera while the family member was presented in 2D on the iPad (Figure 3, right). In the AR-communication condition, the family member was projected through the iPad in 3D onto a chair in front of the resident, such that it seemed that the family member was physically sitting opposite the resident (Figure 3, left). In both conditions, the resident was displayed in 2D video to the family member. This means that the difference between both

Figure 1: Schematic overview of the AR-based communication system. The resident (left) viewed the family member as a 3D projection in AR on the 12.9" iPad Pro and was recorded by a Logitech webcam. The family member (right) viewed the resident as a 2D presentation on a 43" TV screen and was recorded by a Kinect camera. Two computers were used to connect the devices on both sides via the internet.

Figure 2: Rooms and set up of the experiment. On the resident's side (left): a = chair onto which the family member would be projected; b = Logitech webcam; c = iPad; d = Jabra 750 speakerphone; e = position of the resident. On the family member's side (right): f = Azure Kinect camera; g = TV screen; h = Jabra 750 speakerphone; i = position of the family member.

conditions was only visible for the resident, while the family member always saw a 2D image of the resident. The communication mode was only manipulated on the side of the resident, because the AR tool was primarily intended for their use.

Everything else was held constant between conditions. The same screens were used, and the audio signals were similar since the same Jabra 750 speakerphones and Microsoft Teams connections were used for audio transmission. This offered good lip synchronization in both conditions, which was confirmed by expert opinions. The user's self-view in Microsoft Teams was disabled during video calling, since the AR tool also lacked a self-view.

3.3 Statistical analysis

IBM SPSS Statistics 26 (www.ibm.com) for MacOS was used to perform all statistical analyses. A significance level of 0.05 was used for all hypothesis testing. The normality of the data was assessed with Shapiro-Wilk tests. Within-group differences were tested using paired samples *t*-tests for normally distributed data and Wilcoxon signed rank tests for skewed data. Between-group differences were tested using independent samples *t*-tests for normally distributed data and Mann-Whitney U tests for skewed data.

One-way repeated-measures analysis of variance (ANOVA) were conducted due to the within-subjects design of this study. We tested for interaction effects by adding the variable of interest to the model as a between-subjects factor. For UX indicator scores, we used two-way repeated measures ANOVA to test the interaction of group and condition.

4 RESULTS

4.1 UX indicators

There were no significant within-group or between-group differences for any UX indicator scores and neither did we find interaction effects of group and condition (Table 1). The difference in spatial presence for family members was only marginally significant ($z = -1.90, p = 0.06$), with higher scores for the AR-communication condition compared to video calling.

For the family members' responses on the NMQ, we did not find significant differences between conditions either, for total score and subscales (Table 2). The difference in perceived message understanding was only marginally significant ($z = -1.91, p = 0.06$), with

Figure 3: Examples of the AR (left) and video calling (right) conditions on the resident's side, with close-ups of the iPad screen.

Table 1: Mean scores of the UX indicators

		AR-communication	Video calling	Within-group (<i>p</i> value)
Enjoyment	Residents	8.50 ± 1.27	8.50 ± 1.27	1.00
	Family members	8.80 ± 1.14	8.20 ± 0.92	0.11 ^a
	Between-group (<i>p</i> value)	0.56 ^a	0.55	0.26
Spatial presence	Residents	7.60 ± 2.01	7.50 ± 1.96	0.34
	Family members	6.60 ± 3.75	5.10 ± 3.67	0.06 ^a
	Between-group (<i>p</i> value)	0.91 ^a	0.08	0.11
Social presence	Residents	7.60 ± 2.07	7.60 ± 1.96	1.00
	Family members	8.00 ± 2.45	7.90 ± 1.66	0.85 ^a
	Between-group (<i>p</i> value)	0.51 ^a	0.72	0.86

Note. Data are mean ± SD. ^aNonparametric tests were used due to skewed data.

Table 2: Networked Minds Questionnaire scores

	AR-communication	Video calling	<i>p</i> value
Total score	182.60 ± 17.28	178.90 ± 22.64	0.52
Subscales			
Co-presence	6.65 ± 0.73	6.30 ± 0.87	0.40 ^a
Attentional allocation	5.67 ± 0.92	5.23 ± 0.81	0.17 ^a
Perceived message understanding	6.33 ± 0.88	6.08 ± 1.11	0.06 ^a
Perceived affective understanding	5.87 ± 1.07	5.80 ± 1.03	0.86 ^a
Perceived emotional interdependence	3.15 ± 1.90	3.08 ± 1.60	0.88
Perceived behavioral interdependence	4.88 ± 1.74	5.35 ± 1.46	0.24

Note. Data are mean ± SD. For the subscales, total scores are divided by the number of items belonging to that scale due to the number of items per subscale being uneven. All subscales included six items, except perceived message understanding, which included four.

^aNonparametric tests were used due to skewed data.

higher scores for the AR-communication condition compared to video calling.

4.2 Attitudes

For both the AR tool and video calling, 70% of the residents indicated that they were interested in using the technology again. 90% of the family members found repeated use of the AR tool suitable both

Table 3: Behavioral observation scores on the MiDAS

	AR-communication	Video calling	<i>p</i> value
Interest	75.00 ± 17.16	71.50 ± 11.07	0.34 ^a
Response	72.50 ± 17.20	73.00 ± 12.74	0.86
Initiation	71.50 ± 15.28	73.00 ± 7.89	0.59 ^a
Involvement	76.50 ± 14.15	73.50 ± 14.15	0.28
Enjoyment	74.00 ± 15.06	69.00 ± 9.94	0.32

Note. Data are mean ± SD. ^aNonparametric tests were used due to skewed data.

Figure 4: Mean session duration for the AR-communication (left, dark blue) and video calling (right, light blue) conditions. * $p < 0.05$.

for the residents and for themselves. For video calling, this fraction was 70%.

4.3 Behavior

For the behavioral observation scores, there were no significant differences between conditions for all items of the MiDAS (Table 3).

4.4 Use

On average, the sessions with the AR solution lasted significantly longer than those with video calling, $t(9) = 2.52$, $p = 0.03$ (Figure 4). The mean difference in duration was 2.50 minutes ($SD = 3.14$; 95% $CI = [0.26, 4.74]$).

5 DISCUSSION

In this study we compared the UX of communication between nursing home residents and their family members via a newly developed AR-based tool to that of video calling. Family members reported slightly higher levels of spatial presence and perceived message understanding for AR-communication compared to video calling. This is surprising, given the fact that family members perceived the same (2D) image of their relative. This finding may reflect either the family members' positive expectations (since they were aware

of the condition) or subtle changes in the behavior of the residents in the AR condition. Family members preferred the AR tool over video calling for future use, while residents had a positive attitude towards both technologies. The average session duration for AR-communication was significantly longer than for video calling. Contrary to our expectations, we did not find differences between AR-communication and video calling for the residents' UX. Only family members reported slight differences in their UX between both conditions. This may have been due to the residents being less able to verbalize their UX. The difficulty in understanding of and responding to questionnaire items has previously been identified in presence research [19], specifically among nursing home residents in the context of video calling [4]. In this study, it became apparent that some of the residents found it difficult to articulate their experiences in grades. Some residents, for example, attributed the same score to all UX indicators for both conditions, even when their verbal expressions indicated otherwise, while again others tended to give socially desirable scores. Ultimately, the residents reported enjoying the sessions and experiencing feelings of togetherness and connectedness.

It appears that residents did not (consciously) experience differences in UX between conditions. The residents may have been too focused on their family members to notice these differences. In contrast, family members perceived slightly higher spatial presence and message understanding in the AR condition.

5.1 Strengths and limitations

The main limitation of this study (in terms of generalizability and power) was its smaller sample size. Inclusion of participants was restricted because of the COVID-19 pandemic. Future studies should further investigate spatial presence and perceived message understanding during AR-based remote family visits, as these findings were only marginally significant in our study.

A limitation of the AR system was the visibility of the edges of the screen, which may have prevented a convincing seamless experience. For future development, the option of removing visible screen edges should be explored.

Also, this study only involved a one-time use of the AR solution. A suggestion for future research is to study long(er) term use. This will allow for gaining deeper insight into the actual suitability of the AR solution for remote family visits in nursing homes. For example, this can involve studying effects of repeated use of the AR solution on the residents' (mental) well-being and loneliness.

Lastly, as mentioned above, the evaluation methodology to assess the UX of the residents may need to be tailored to this population

to be able to assess their experiences in more detail. Indicators pertaining to duration, frequency and consistency of use may be useful in this context.

6 CONCLUSION

In this study, we compared the UX of a new AR-based communication tool with video calling for social interaction between residents in nursing homes and their remote family members. For residents, we found no differences in their UX with both communication modes. Family members reported marginally higher spatial presence in the AR-communication condition compared to video calling. Also, the average duration of AR-communication sessions was longer compared to video calling.

In this study, we set out to determine whether AR-based communication would be overall advantageous over video calling. In this regard, the evidence we presented here, unfortunately, remains inconclusive. Nevertheless, it should be noted that the results do suggest that AR-based communication can be considered a viable alternative to video calling, for maintaining contact between residents of nursing homes and their friends and family.

ACKNOWLEDGMENTS

This research was part of a collaboration between The Netherlands Organisation for applied scientific research (TNO) and Meander-Groep, a healthcare provider in South-Limburg, the Netherlands. This project was partially funded by TNO's Early Research Project 'Social eXtended Reality'.

REFERENCES

- [1] Kristel Paque, Hilde Bastiaens, Peter Van Bogaert, & Tinne Dilles. 2018. Living in a nursing home: a phenomenological study exploring residents' loneliness and other feelings. *Scandinavian Journal of Caring Sciences*, 32 (4), 1477-1484. <https://doi.org/10.1111/scs.12599>.
- [2] C. Noone, J. McSharry, M. Smalle, A. Burns, K. Dwan, D. Devane, & E. C. Morrissey. 2020. Video calls for reducing social isolation and loneliness in older people: a rapid review. *Cochrane Database of Systematic Reviews*, (5), Art. No.: CD013632. <https://doi.org/10.1002/14651858.CD013632>.
- [3] Ryoko Furukawa, & Martha Driessnack. 2013. Video-mediated communication to support distant family connectedness. *Clinical Nursing Research*, 22 (1), 82-94. <https://doi.org/10.1177/1054773812446150>.
- [4] Sonam Zamir, Catherine Hagan Hennessy, Adrian H. Taylor, & Ray B. Jones. 2018. Video-calls to reduce loneliness and social isolation within care environments for older people: an implementation study using collaborative action research. *BMC Geriatrics*, 18 (1), 62. <https://doi.org/10.1186/s12877-018-0746-y>.
- [5] Robby Nadler. 2020. Understanding "Zoom fatigue": Theorizing spatial dynamics as third skins in computer-mediated communication. *Computers and Composition*, 58, 102613. <https://doi.org/10.1016/j.compcom.2020.102613>.
- [6] Catherine S. Oh, Jeremy N. Bailenson, & Gregory F. Welch. 2018. A systematic review of social presence: definition, antecedents, and implications. *Frontiers in Robotics and AI*, 5 (114) <https://doi.org/10.3389/frobt.2018.00114>.
- [7] Nathan M. D'Cunha, Dung Nguyen, Nenad Naumovski, Andrew J. McKune, Jane Kellett, Ekavi N. Georgousopoulou, Jane Frost, & Stephen Isbel. 2019. A mini-review of virtual reality-based interventions to promote well-being for people living with dementia and mild cognitive impairment. *Gerontology*, 65 (4), 430-440. <https://doi.org/10.1159/000500040>.
- [8] Li Na Lee, Mi Jeong Kim, & Won Ju Hwang. 2019. Potential of augmented reality and virtual reality technologies to promote wellbeing in older adults. *Applied Sciences*, 9 (17), 3556. <https://doi.org/10.3390/app9173556>.
- [9] Simon N.B. Gunkel, Martin Prins, Hans Stokking, & Omar Niamut. 2017. Social VR platform: Building 360-degree shared VR spaces. In: *Adjunct Publication of the 2017 ACM International Conference on Interactive Experiences for TV and Online Video. Association for Computing Machinery*: 3089914, 83-84. <https://doi.org/10.1145/3084289.3089914>.
- [10] Frank Biocca, Chad Harms, & Jenn Gregg. 2001. The Networked Minds measure of social presence: Pilot test of the factor structure and concurrent validity. In: *4th Annual International Workshop on Presence*, 1-9.
- [11] Mel Slater, & Sylvia Wilbur. 1997. A framework for immersive virtual environments (FIVE): Speculations on the role of presence in virtual environments. *Presence: Teleoperators and Virtual Environments*, 6 (6), 603-616. <https://doi.org/10.1162/pres.1997.6.6.603>.
- [12] Jie Li, Yiping Kong, Thomas Röggl, Francesca De Simone, Swamy Ananthanarayan, Huib de Ridder, Abdallah El Ali, & Pablo Cesar. 2019. Measuring and understanding photo sharing experiences in social virtual reality. In: *Conference on Human Factors in Computing Systems (CHI 2019)* New York, USA. <https://doi.org/10.1145/3290605.3300897>.
- [13] Mark Roman Miller, Hanseul Jun, Fernanda Herrera, Jacob Yu Villa, Greg Welch, & Jeremy N. Bailenson. 2019. Social interaction in augmented reality. *PLOS ONE*, 14 (5), e0216290. <https://doi.org/10.1371/journal.pone.0216290>.
- [14] Kuo Ting Huang, Christopher Ball, Jessica Francis, Rabindra Ratan, Josephine Boumis, & Joseph Fordham. 2019. Augmented versus virtual reality in education: An exploratory study examining science knowledge retention when using augmented reality/virtual reality mobile applications. *Cyberpsychology, Behavior, and Social Networking*, 22 (2), 105-110. <https://doi.org/10.1089/cyber.2018.0150>.
- [15] Tim Hilken, Ko de Ruyter, Mathew Chylinski, Dominik Mahr, & Debbie I. Keeling. 2017. Augmenting the eye of the beholder: exploring the strategic potential of augmented reality to enhance online service experiences. *Journal of the Academy of Marketing Science*, 45 (6), 884-905. <https://doi.org/10.1007/s11747-017-0541-x>.
- [16] Anne R. Smink, Eva A. van Reijmersdal, Guda van Noort, & Peter C. Neijens. 2020. Shopping in augmented reality: The effects of spatial presence, personalization and intrusiveness on app and brand responses. *Journal of Business Research*, 118, 474-485. <https://doi.org/10.1016/j.jbusres.2020.07.018>.
- [17] Chad Harms, & Frank Biocca. 2004. Internal consistency and reliability of the Networked Minds measure of social presence. In: M. Alcaniz, & B. Rey (Eds.), *Seventh Annual International Workshop: Presence 2004*. Universidad Politecnica de Valencia: Valencia, Spain, 246-251.
- [18] Oriei McDermott, Vasiliki Orgeta, Hanne Mette Ridder, & Martin Orrell. 2014. A preliminary psychometric evaluation of Music in Dementia Assessment Scales (MiDAS). *International Psychogeriatrics*, 26 (6), 1011-1019. <https://doi.org/10.1017/S1041610214000180>.
- [19] Simone Grassini, & Karin Laumann. 2020. Questionnaire measures and physiological correlates of presence: A systematic review. *Frontiers in Psychology*, 11, Article 349. <https://doi.org/10.3389/fpsyg.2020.00349>.