



UvA-DARE (Digital Academic Repository)

Search for resonant and non-resonant Higgs boson pair production in the $b\tau^+\tau^-$ decay channel in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector

Aaboud, M.; The ATLAS Collaboration

DOI

[10.1103/PhysRevLett.121.191801](https://doi.org/10.1103/PhysRevLett.121.191801)

Publication date

2018

Document Version

Other version

Published in

Physical Review Letters

License

CC BY

[Link to publication](#)

Citation for published version (APA):

Aaboud, M., & The ATLAS Collaboration (2018). Search for resonant and non-resonant Higgs boson pair production in the $b\tau^+\tau^-$ decay channel in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. *Physical Review Letters*, 121(19), [191801].
<https://doi.org/10.1103/PhysRevLett.121.191801>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Erratum: Search for Resonant and Nonresonant Higgs Boson Pair Production in the $b\bar{b}\tau^+\tau^-$ Decay Channel in pp Collisions at $\sqrt{s} = 13$ TeV with the ATLAS Detector [Phys. Rev. Lett. **121, 191801 (2018)]**

M. Aaboud *et al.**

(ATLAS Collaboration)



(Received 5 February 2019; published 27 February 2019)

DOI: [10.1103/PhysRevLett.122.089901](https://doi.org/10.1103/PhysRevLett.122.089901)

One correction is noted, which does not affect the results or limits reported. The acceptance times efficiency numbers quoted for the nonresonant signal in the Letter (but not those used in the analysis) did not take into account the effect of the MADGRAPH5_aMC@NLO Monte Carlo event weights and are therefore incorrect. In the paragraph discussing the final fit and limit-setting procedure, the corrected sentence should read as follows: The acceptance times efficiency for the NR HH signal is 3.2% (1.9%) in the combined SLT and LTT $\tau_{\text{lep}}\tau_{\text{had}}$ ($\tau_{\text{had}}\tau_{\text{had}}$) channel over the full BDT distribution, decreasing to 1.8% (1.4%) for the two most sensitive BDT bins.

*Full author list given at the end of the original Letter.

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published articles title, journal citation, and DOI.