



UvA-DARE (Digital Academic Repository)

De novo induction of resistance against voriconazole in *Aspergillus fumigatus*

Händel, N.; de la Sayette, S.; Verweij, P.E.; Brul, S.; ter Kuile, B.H.

DOI

[10.1016/j.jgar.2015.01.001](https://doi.org/10.1016/j.jgar.2015.01.001)

Publication date

2015

Document Version

Final published version

Published in

Journal of Global Antimicrobial Resistance

License

Article 25fa Dutch Copyright Act (<https://www.openaccess.nl/en/policies/open-access-in-dutch-copyright-law-taverne-amendment>)

[Link to publication](#)

Citation for published version (APA):

Händel, N., de la Sayette, S., Verweij, P. E., Brul, S., & ter Kuile, B. H. (2015). De novo induction of resistance against voriconazole in *Aspergillus fumigatus*. *Journal of Global Antimicrobial Resistance*, 3(1), 52-53. <https://doi.org/10.1016/j.jgar.2015.01.001>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.



Letter to the Editor

De novo induction of resistance against voriconazole in *Aspergillus fumigatus*

Sir,

The opportunistic pathogen *Aspergillus fumigatus* causes invasive aspergillosis in patients with reduced immune function [1]. Infections caused by this organism are difficult to treat, particularly in critically ill patients. Therapy prospects become very dismal when the pathogen is resistant to the various azole compounds that are the drugs of choice against this organism.

The same resistance mechanism, designated TR₃₄/L98H, was uncovered in clinical isolates from patients who had not been treated with azoles previously as in *A. fumigatus* exposed to azole fungicides used for crop protection and material preservation [2]. This observation suggests that the origin of this resistance mechanism may lie in the use of azoles in the environment, since resistant conidia can easily reach the human population transported by air or in any other way. What is not understood at present is whether azole resistance in *A. fumigatus* can be induced de novo by exposure to sublethal drug concentrations, as was shown for *Escherichia coli* [3], or whether horizontal transfer of resistance genes is essential. The development of resistance during patient therapy suggests that de novo emergence of resistance is possible [4,5], but since genetic exchange cannot entirely be ruled out, this needs to be proven under conditions that do not allow gene transfer.

An azole-susceptible strain of *A. fumigatus*, designated AZN 8196, was grown on plates as described previously [2] with sublethal concentrations of voriconazole and itraconazole. Following 7 days of incubation, the strains were transferred to (i) a plate with the same drug concentration and (ii) a plate with a two times higher level. When the plate with the higher drug concentration showed growth within 1 week, the plate with the lower level was discarded and the procedure was repeated. At every transfer, the minimum inhibitory concentration (MIC) of the strain was measured by following growth during 48 h in a set of liquid cultures having drug concentrations stepwise increasing by a factor of 2.

Exposure of *A. fumigatus* to stepwise increasing concentrations of voriconazole starting at 0.125 µg/mL resulted in a rapid increase of the MIC once the concentration of the drug exceeded the initial MIC (Fig. 1). After 5 weeks, the MIC reached 8 µg/mL, even though the concentration in the plate was only 2 µg/mL. A similar attempt to induce itraconazole resistance was not successful. The MIC for itraconazole went up only by a factor of 2, from 0.125 µg/mL to 0.25 µg/mL. The sharp increase in MIC for voriconazole suggests that a genetic mutation may play a role. This idea was investigated by sequencing PCR products of the *cyp51A* and *cyp51B* genes that contained known hotspots for azole resistance in *A. fumigatus* [5]. The wild-type and voriconazole-resistant sequence were unchanged compared with the published sequences of the *cyp51A* and *cyp51B* genes as recorded in GenBank under accession nos.




Fig. 1. Minimum inhibitory concentrations (MICs) of *Aspergillus fumigatus* for voriconazole and itraconazole as a function of culture time in days. The concentration of voriconazole in the substrate increased stepwise by a factor of 2 each time from 0.125 µg/mL to 4 µg/mL during the experiment.

AF338659 and AF338660, respectively. The only mutations found were in the *cyp51B* gene of the strains grown on plates that had an itraconazole concentration of 0.25 µg/mL; these were two silent mutations (S274S and P394P). Hence, if any mutations were involved in developing this de novo azole resistance, they were not in the *cyp51A* and *cyp51B* genes.

On the one hand it is possible that *A. fumigatus* adjusts itself by differentially regulating expression levels to induce, for example efflux pumps, as was found for de novo acquisition of antibiotic resistance in *E. coli* [6]. Even this type of adaptation has lasting effects and is not readily reversed when the drug is no longer present. On the other hand it is equally likely that resistance to voriconazole is caused by different mutations than the well known ones in *cyp51A* and *cyp51B*. Regardless, in both cases the main conclusion is that azole resistance in *A. fumigatus* is readily induced by exposure to sublethal concentrations of these drugs. Use of a single strain may limit the general applicability of this conclusion, but the proof of principle remains valid. Especially in agricultural practice, where the application is often not overly precise, sublethal concentrations might be encountered. It is therefore essential that unnecessary exposure of *A. fumigatus* to these essential drugs for human medicine is prevented.

Funding

Funding was provided by The Netherlands Food and Consumer Product Safety Authority (Utrecht, The Netherlands).

Competing interests

BHtK is employed by The Netherlands Food and Consumer Product Safety Authority. All other authors declare no competing interests.

Ethical approval

Not required.

Acknowledgment

The expert technical assistance of Mr. A. Rijs is greatly appreciated.

References

- [1] Vermeulen E, Lagrou K, Verweij PE. Azole resistance in *Aspergillus fumigatus*: a growing public health concern. *Curr Opin Infect Dis* 2013;26:493–500.
- [2] Verweij PE, Snelders E, Kema GH, Mellado E, Melchers WJ. Azole resistance in *Aspergillus fumigatus*: a side-effect of environmental fungicide use? *Lancet Infect Dis* 2009;9:789–95.
- [3] van der Horst MA, Schuurmans JM, Smid MC, Koenders BB, ter Kuile BH. De novo acquisition of resistance to three antibiotics by *Escherichia coli*. *Microb Drug Resist* 2011;17:141–7.
- [4] Camps SM, van der Linden JW, Li Y, Kuijper EJ, van Dissel JT, Verweij PE, et al. Rapid induction of multiple resistance mechanisms in *Aspergillus fumigatus* during azole therapy: a case study and review of the literature. *Antimicrob Agents Chemother* 2012;56:10–6.
- [5] Escribano P, Recio S, Peláez T, González-Rivera M, Bouza E, Guinea J. In vitro acquisition of secondary azole resistance in *Aspergillus fumigatus* isolates after prolonged exposure to itraconazole: presence of heteroresistant populations. *Antimicrob Agents Chemother* 2012;56:174–8.
- [6] Arendrup MC, Mavridou E, Mortensen KL, Snelders E, Frimodt-Møller N, Khan H, et al. Development of azole resistance in *Aspergillus fumigatus* during azole therapy associated with change in virulence. *PLoS ONE* 2010;5: e10080.

Nadine Händel

Sarah de la Sayette

University of Amsterdam, Swammerdam Institute of Life Sciences,
Science Park 904, 1098 XH Amsterdam, The Netherlands

Paul E. Verweij

Department of Medical Microbiology,

Radboud University Medical Centre, Nijmegen, The Netherlands

Stanley Brul

University of Amsterdam, Swammerdam Institute of Life Sciences,
Science Park 904, 1098 XH Amsterdam, The Netherlands

Benno H. ter Kuile^{a,b,*}

^aUniversity of Amsterdam, Swammerdam Institute of Life Sciences,
Science Park 904, 1098 XH Amsterdam, The Netherlands

^bThe Netherlands Food and Consumer Product Safety Authority,
Catharijnesingel 59, 3511 GG Utrecht, The Netherlands

*Corresponding author at: University of Amsterdam,
Swammerdam Institute of Life Sciences, Science Park 904, 1098 XH
Amsterdam, The Netherlands. Tel.: +31 64 659 6684.

E-mail addresses: B.H.terKuile@uva.nl,
b.h.terkuile@NVWA.nl (B.H. ter Kuile).

9 June 2014