



UvA-DARE (Digital Academic Repository)

Deep-learning based reconstruction of the shower maximum X_{\max} using the water-Cherenkov detectors of the Pierre Auger Observatory

Pierre Auger Collaboration

DOI

[10.1088/1748-0221/16/07/P07019](https://doi.org/10.1088/1748-0221/16/07/P07019)

Publication date

2021

Document Version

Final published version

Published in

Journal of Instrumentation

License

CC BY

[Link to publication](#)

Citation for published version (APA):

Pierre Auger Collaboration (2021). Deep-learning based reconstruction of the shower maximum X_{\max} using the water-Cherenkov detectors of the Pierre Auger Observatory. *Journal of Instrumentation*, 16(07), Article P07019. <https://doi.org/10.1088/1748-0221/16/07/P07019>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Deep-learning based reconstruction of the shower maximum X_{\max} using the water-Cherenkov detectors of the Pierre Auger Observatory



PIERRE
AUGER
OBSERVATORY

The Pierre Auger collaboration

E-mail: spokespersons@auger.org

ABSTRACT: The atmospheric depth of the air shower maximum X_{\max} is an observable commonly used for the determination of the nuclear mass composition of ultra-high energy cosmic rays. Direct measurements of X_{\max} are performed using observations of the longitudinal shower development with fluorescence telescopes. At the same time, several methods have been proposed for an indirect estimation of X_{\max} from the characteristics of the shower particles registered with surface detector arrays. In this paper, we present a deep neural network (DNN) for the estimation of X_{\max} . The reconstruction relies on the signals induced by shower particles in the ground based water-Cherenkov detectors of the Pierre Auger Observatory. The network architecture features recurrent long short-term memory layers to process the temporal structure of signals and hexagonal convolutions to exploit the symmetry of the surface detector array. We evaluate the performance of the network using air showers simulated with three different hadronic interaction models. Thereafter, we account for long-term detector effects and calibrate the reconstructed X_{\max} using fluorescence measurements. Finally, we show that the event-by-event resolution in the reconstruction of the shower maximum improves with increasing shower energy and reaches less than 25 g/cm^2 at energies above $2 \times 10^{19} \text{ eV}$.

KEYWORDS: Data analysis; Pattern recognition, cluster finding, calibration and fitting methods; Large detector systems for particle and astroparticle physics; Particle identification methods

ARXIV EPRINT: [2101.02946](https://arxiv.org/abs/2101.02946)



Contents

1	Introduction	1
2	Data sets and their preparation	3
2.1	Simulation libraries	4
2.2	Hybrid dataset	4
2.3	Pre-processing of data	5
2.4	Augmentation of simulated data	6
3	Deep neural network for reconstructing the shower maximum	6
3.1	Architecture	7
3.2	Training	9
4	Performance on simulations	9
4.1	Training and evaluation of the network using EPOS-LHC simulated events	11
4.2	Evaluation of the network on QGSJetII-04 and Sibyll 2.3 simulated events	11
4.3	Zenith dependency of the reconstruction	12
4.4	Distribution of the reconstructed shower maxima	14
5	Application to hybrid data	15
5.1	Corrections for detector-aging effects	15
5.2	Evaluation and calibration of the first moment of the distribution of shower maxima	16
5.3	Potential for determining the second moment of the distribution of shower maxima	18
6	Summary	18
	The Pierre Auger collaboration	24

1 Introduction

The depth of the air shower maximum X_{\max} induced by ultra-high energy cosmic rays is a key observable to estimate the mass of primary cosmic particles (see e.g. [1] for a review). In this work we introduce a new method to reconstruct X_{\max} from signals recorded by water-Cherenkov detectors (WCDs), whose duty cycle is almost 100%, unlike fluorescence telescopes, which can only operate during moonless nights with a duty cycle of about 15%.

The Pierre Auger Observatory [2] was constructed to study cosmic rays at the highest energies with high accuracy. With its instrumented area of 3000 km² the observatory is sufficiently large to measure the rare high-energy events up to 10²⁰ eV with adequate statistics. Designed as a hybrid instrument, on the one hand the development of the air showers is directly followed by observing the fluorescence light emitted by nitrogen molecules in the air with 27 telescopes,

constituting the fluorescence detector (FD). As measurements using the fluorescence technique are only possible during moonless nights it is used to collect data for an in-depth understanding of the air showers [3–6] with reduced statistics. On the other hand, secondary particles of the air showers are detected by 1600 water-Cherenkov Detectors (WCDs) located on the ground, which form the surface detector (SD). The detectors are located at a distance of 1.5 km from each other in a hexagonal configuration (figure 1(a)) and measure the time-dependent density of secondary shower particles. The measurement is based on Cherenkov light emitted by the relativistic particles as they pass through each water-filled detector. This light is recorded by three photomultiplier tubes (PMTs) attached to the top of the tank and processed using flash analog-to-digital converters with a sampling rate of 40 MHz, translating to bins with a width of 25 ns.

For the determination of the air shower direction and energy with the WCDs there are already established precise reconstruction methods [7] that use the arrival times of the shower particles and the spatial extent of the air shower. Reconstructing the depth of maximum of the shower, on the other hand, is a major challenge. In contrast to the fluorescence telescopes, there is no direct observation of the shower development in the atmosphere. Instead, the shower properties are encoded in the signal traces induced by the secondary particles traversing the WCDs.

During the longitudinal development of the air shower, the relative abundances of hadrons, muons, electrons and photons change in addition to geometric dependencies such as the distance to the shower core or the zenith angle. The signal traces generated by these individual components in the three PMTs of the WCDs are different and thus provide the possibility to extract information about the depth of the shower maximum X_{\max} . Whereas muons do not interact with the atmosphere and reach the detector usually in earlier times, the electromagnetic component is strongly shielded and reach the detector later and more spread in time. See figure 1(b) for a simulated example trace with muons and electromagnetic shower particles.

Experimental proof of the successful use of the signal characteristics to determine the depth of maximum of the shower has already been achieved. It was demonstrated [8, 9] that using shower universality [10, 11] X_{\max} can be measured with good accuracy by decomposing the signal traces using templates of typical waveforms induced by the different shower components using the universality of signals with respect to the shower development stage. Furthermore, the correlation between the risetimes of the signals in the WCDs was used to determine X_{\max} [12]. With this method the average composition of UHECRs was determined up to 100 EeV. The measurement up to such high energies has been only made possible using the large exposure and high reconstruction efficiency of the SD, increasing the statistics by a factor 25 above 3 EeV (factor 12 above 30 EeV) when compared to the FD [13].

Our new method aims to measure, beside the average composition, the mixing of the composition $\sigma(X_{\max})$ with high statistics by exploiting mass-sensitive information on an event-by-event basis. This opens up possibilities for event-based anisotropy studies using information on the cosmic-ray mass.

This method to reconstruct X_{\max} using the measured signals of the WCDs is based on a deep neural network (DNN), whose architecture and optimization was developed specifically for the measurement conditions at the Pierre Auger Observatory. The network is a further development of [14] and takes into account current developments in machine learning, especially elements from speech recognition and utilization of spatial symmetries.




Figure 1. (a) Simulated signal pattern measured by the surface detector. The marker sizes indicate the amount of measured signal and the colors represent the arrival time of the shower at a given station (yellow=early, red=late). The arrow denotes the projection of the shower axis on the surface and its tip the shower core. (b) Simulated signal trace of a cosmic-ray event measured at a surface-detector station at a distance of about 1000 m to the shower core. Different colors indicate signals from different shower components.

In the hybrid network architecture, all time intervals of the signal traces of the PMTs are first scanned using a recurrent subnetwork. Thereby each signal trace is characterized by a total of 10 “machine-learned features“. In the following network layers these observables are merged with the particle arrival times at the respective WCD locations. When processing the signals at the given positions, the hexagonal symmetry of the observatory is generically mapped into the network architecture. The network parameters for forming the 10 observables to characterize the signal traces and for combining all available information are adjusted in a training procedure. The most important tools of this optimization are the training data and the objective function, which is minimized during network training.

This work is structured as follows. First, we specify the data sets for both the simulation studies and measured Auger hybrid data, which include information from the FD for validation purposes. We explain in detail how the simulated data are prepared and augmented for the optimization of the network parameters and the reconstruction of X_{max} . After that, we describe in detail the architecture and training of the deep network. Then we show the X_{max} reconstruction performance of the network on simulated data as a function of energy, zenith angle, mass of the primary particle, and the effect of using two hadronic interaction models different from the one used in the training. Finally, we verify the capabilities of the network by direct comparison of the measured maximum shower depth X_{max} of the network and of the FD. We correct for detector-aging effects resulting from long-term operation of the observatory. Subsequently, we calibrate the absolute X_{max} value of the network output, and determine the X_{max} resolution of the network as a function of the primary energy.

2 Data sets and their preparation

The measured air shower footprint consists of a characteristic pattern of several triggered WCDs arranged in a hexagonal grid (see figure 1(a)). Using three PMTs each triggered station measures

the time-dependent density of particles encoded in three signal traces. An example of a simulated signal trace is shown in figure 1(b).

The basic idea is to provide the network as input the raw data of a measured cosmic-ray event. The raw information for each triggered station consists of three signal time traces, the station position and the time of the first shower particles arriving at the station.

For successful adjustment of the network parameters, careful preparations of the data sets from simulation campaigns used for the optimization process are crucial. For example, the parameters can be set more easily if the numerical values of the input variables do not vary considerably. Therefore, both the amplitudes and the time values of the WCD-signal traces are re-scaled prior to their input into the network. Also, typical conditions when operating the observatory need to be included in the simulated training data used for adjusting the parameters.

Below we specify the data sets from simulation campaigns, and data with information from the FD used to validate the X_{\max} reconstruction of the deep neural network.

2.1 Simulation libraries

For generating training and validation data we use CORSIKA 7.6400 [15] for the simulation of extensive air showers. The Pierre Auger Offline software [16] is used for the simulation of the detector responses, which is based on GEANT4 [17], and the reconstruction of the air shower parameters [2]. The CORSIKA showers are reused 5–30 times at other locations of the observatory along with a re-simulation of the detector responses. The simulation library consists of events for primary hydrogen, helium, oxygen and iron, uniformly distributed in azimuth angle and following a zenith angle distribution flat in $\cos^2 \theta$ with $0^\circ \leq \theta < 65^\circ$. The energy range covers 1–160 EeV ($1 \text{ EeV} = 10^{18} \text{ eV}$) and follows a spectrum of E^{-1} . For the training of the network we use exclusively simulations with the EPOS-LHC hadronic interaction model [18]. Overall, the training set contains around 400,000 events.

For testing the performance we prepare a test set with EPOS-LHC showers containing 50,000 events and two sets of showers simulated using the hadronic interaction model QGSJetII-04 [19] and Sibyll 2.3 [20] containing 450,000 events each. In contrast to the training data for the test sets we limit the energy range to 3–160 EeV and the zenith angle range to $0\text{--}60^\circ$, where the standard SD reconstruction shows no significant reconstruction biases [2]. The extended phase space during training prevents incorrectly reconstructed energies and zenith angles at the edges of the phase space.

2.2 Hybrid dataset

The accuracy of the deep-learning based reconstruction of X_{\max} is examined using a high-quality set of hybrid data where nearly unbiased X_{\max} measurements are performed using the fluorescence technique. A complete description of the hybrid reconstruction and the high-quality event selection can be found in [3].

Here, events are only kept if X_{\max} falls into the FD field of view and the fraction of the shower profile inside the field of view is large enough to allow an unbiased measurement of X_{\max} . Additionally, we discard events below the full trigger efficiency of the surface detector at roughly 3.16 EeV to prevent biased reconstructions. Further, we remove events (less than $< 0.5\%$) with area over peak (A/P) values [21], given in units of 25 ns, which are very low (< 2.7) or high

(> 3.5) and cannot be corrected using a linear function (see section 5.1). To guarantee a precise SD reconstruction, we reject events with broken or non-existent stations in the first hexagon formed around the station with the largest signal [22]. After the selection 3109 events, collected from 1 January 2004 to 31 December 2017, remain for the evaluation of the DNN performance.

2.3 Pre-processing of data

Footprints of air showers on the ground can reach sizes of several tens of square kilometers. To reduce the memory consumption and increase generalization capacities of the DNN, we use only the information from a fixed size of the SD consisting of a sub-array of 13×13 WCDs around the WCD with the highest signal.

For air showers with zenith angles below $\theta = 60^\circ$ more than 99% of the triggered stations are contained within this sub-array. We expect the effect of enlarging the sub-grid to be negligible as only stations with very low signals lie outside the sub-grid.

As the positions of the WCDs feature a hexagonal grid several representations of the local neighborhood are possible [14]. Our algorithm is based on the so-called axial representation.

Signal traces. The total number of particles measured in each detector is expected to approximate a power law as a function of the distance to the shower axis. Hence, we use a logarithmic re-scaling of the signal traces $S_i(t)$:

$$\tilde{S}_i(t) = \frac{\log_{10}[S_i(t)/\text{VEM} + 1]}{\log_{10}[S_{\text{norm}}/\text{VEM} + 1]}. \quad (2.1)$$

Here VEM (vertical equivalent muon) is a unit equal to the signal induced by a muon traversing the WCD vertically and $S_{\text{norm}} = 100$ VEM normalizes signals of 100 VEM to unity. To allow for positive values only and to leave contributions with $S = 0$ unchanged we use $S(t) + 1$.

Particle arrival times. The curvature of the shower front is reflected in the arrival times $t_{0,i}$ of the first shower particles at the SD stations. For each triggered station in an event, the particle arrival time is normalized with respect to the arrival time τ_{center} measured at the station with the largest signal and the standard deviation $\sigma_{t,\text{train data}}$ of the arrival times of the complete training set:

$$\tilde{t}_{0,i} = \frac{t_{0,i} - \tau_{\text{center}}}{\sigma_{t,\text{train data}}}. \quad (2.2)$$

Station states. Air showers falling close to the edges of the detector grid can have footprints with a substantial fraction lying outside the surface-detector grid. Also, in rare cases, a WCD within the 13×13 sub-grid may not provide a signal owing to technical problems. To provide the information in the network training if the absence of a signal originates from air shower physics or from detector effects, we add a feature map for the given sub-grid encoding the station states as additional input (1 = ready to measure, 0 = broken or missing in the grid).

Network prediction. During the training of the network the prediction (output) of the neural network is compared to the *true* labels of the dataset in each iteration. As the input values of the network lie approximately in the range of $[-1; 1]$ and the weights of the model are properly initialized, the expected output values of the network follow approximately a Gaussian with standard deviation $\sigma = 1$ and mean $\mu = 0$.

To speed-up the learning process we also normalize the label values to lie in the same range using standard normalization ($\sigma = 1, \mu = 0$). Instead of normalizing the label values directly, the normalization is implemented as the final layer in our network for the X_{\max} output allowing us to monitor the learning process in physics quantities.

2.4 Augmentation of simulated data

Depending on the sub-grid, the detector states of the WCDs vary on an event-by-event basis. Furthermore, there are differences between detector states in data and simulation. Even small deviations between simulation and data can have a non-negligible influence on the network prediction if there are differences in the phase space. For example, a broken WCD would create a hole in the measured footprint which then distracts the reconstruction if the model is not trained on such detector conditions. To make the algorithm robust to distortions and fluctuations, we implemented an on-the-fly augmentation. Hence during the training every particular simulated event will feature in each repeated iteration different detector states. Note that this augmentation is exclusively used in the network training procedure and is not applied when evaluating the network.

Photomultiplier tubes without response. During detector operation, one or more of the three photomultiplier tubes (PMTs) of a WCD may not respond. To make the network robust to missing signal traces, we randomly mask signal traces during the training. The number of broken PMTs is approximated as Poisson distribution and was tuned to follow the data distribution.

Water-Cherenkov detectors without response. As already stated above, a map of detector stations without response is used describing the current station states. During the network training, detector stations are randomly marked as broken, by setting all measured properties to zero, which is additionally accounted for in the map of stations without response. The number of broken stations is approximated as Poisson distribution and was tuned to follow the distribution of measured data.

Saturated signal traces. Shower geometries with a shower core close to a detector station can exceed the maximum signal that can be quantitatively recorded. This effect is even more frequent at the highest energies, where 50% events of air showers with 50 EeV exhibit typically one saturated detector station. Each surface-detector station is calibrated for the value of 1 VEM in units of the hardware every 60 s. This value changes for each detector station and PMT owing to individual hardware properties and other external influences. Since the number of units is limited, the saturation value is time dependent and differs for each detector station [7].

To make the network robust to different saturation values, we sample the maximum measurable values from a Gaussian distribution adapted by typical saturation values observed in data. In detail we make use of an one-sided truncated Gaussian to prevent very small and negative saturation values. Independently for each PMT, we finally clip the unsaturated trace provided by the simulation according to the sampled saturation value.

3 Deep neural network for reconstructing the shower maximum

Two challenges are to be mastered for reconstructing X_{\max} from the signal traces of the water-Cherenkov Detectors:

1. To develop a network architecture that is optimally suited for the situation of the Pierre Auger Observatory and exploits all symmetries in the recorded data,
2. To adapt about 10^6 network parameters in a way that the correct shower depth is reconstructed under realistic operation conditions of the observatory.

In the following both the network architecture and the network training are described in detail.

3.1 Architecture

To reconstruct the shower maximum X_{\max} using a deep neural network we designed an architecture featuring hexagonal convolutions [23, 24] and recurrent layers illustrated in figure 2. We observed that using an architecture which supports the symmetry of the hexagonal grid of the detectors not only boosts the performance, but further makes the network robust to small fluctuations in the data which results in an improved generalization performance.




Figure 2. Architecture of the deep neural network used to reconstruct X_{\max} . Convolutional operations are shown with their kernel sizes, e.g. a “ 3×3 ” hexagonal convolution translates to a kernel covering one hexagon (7 WCD locations). The numbers in brackets denote the output shapes of the operations.

The input for the neural network is a tensor containing the signal time traces, the map of the arrival times and a map of the station states. For connecting the different physics quantities and extracting the maximum information from the data, we adapted state-of-the-art methods of computer vision and speech recognition.

We introduce a prior on physics relationships, hence learning first to extract information of the shower development using the signal traces before forming features of space and time. The architecture can be split into three parts, discussed in the following.

Detector-wise recurrent processing of signal traces. In the first part of the architecture, the signal trace is processed. To resolve long- and short-term correlations in the signal trace we make use of 1 bi-directional layer and 1 one-directional layer of LSTM cells [25]. Using bidirectional layers allows to connect not only the past time steps with the current step but additionally all future steps.

The most important concept is the station-wise sharing of the recurrent layers. As the physics of the measured traces is the same for each station, we share the trace network over all stations in the 13×13 sub-grid. Hence, exactly the same parameters are used in each station.

Note that this concept still allows the network to extract many very different features and not only a distinct characteristic of the trace. Although the same feature is extracted in each station (e.g. slope of the rising edge) each respective feature value will vary (e.g. slope differs in each station) within one event.

In this sub-network, the tensor of signal traces with a length of 120 time steps is reduced to 10 feature maps encoding the longitudinal shower development. Note that these 10 features, which are extracted for each station, are not hand-designed but are learned by the network to be particularly useful for the reconstruction of X_{\max} . As every station uses the same network, the resulting 10 feature maps share the same physics meaning, allowing to use convolutions in the spatial dimensions in the following part.

Densely-connected hexagonal convolutions. After analyzing the signal trace, the additional information, consisting of the arrival times of the first shower particles at the WCDs and of detector stations without response, is concatenated with the tensor already processed.

In this second part of the architecture, features relating space and time are extracted using convolutional operations on the WCD positions. Before applying hexagonal convolutions the representation needs to be transformed into the space of hexagonal rotations (P6), which is done individually for each feature map. In this space each filter holds 6 feature maps, one for each orientation [24].

The main concept of this block are densely-connected convolutions [26], which provide already extracted features in each subsequent layer. The resulting connections stabilize the training process by feature-reuse and an improved propagation of gradients. The block consists of 3 layers with ELUs [27] (exponential linear units) as activations and with kernels of size 7 covering one hexagon.

Residual reconstruction modules. The last part of the network features residual modules introduced by ResNet [28]. In the standard layout of the AixNet architecture [29] for each reconstruction task (shower axis, shower core, energy and X_{\max}) an individual subpart exists. Each part consists of 2 residual modules with 2 layers each, ReLUs (rectified linear units) as activation functions

and pooling operations between the blocks. The final layer of the X_{\max} block predicts the shower maximum after rescaling back to physics quantities as described in section 2.3.

3.2 Training

For training of the network we use 400,000 simulated air shower events with a mixed composition of hydrogen, helium, oxygen and iron (in equal parts). During the training we use the data augmentation as described in section 2.4 to increase the amount of data and to improve the generalization capacities of the network.

To penalize composition biases we use an element-wise mean squared error (MSE) as objective function:

$$\mathcal{L} = \sum_{Z=H,He,O,Fe} \langle \text{MSE}(\hat{X}_{\max,Z}) \rangle = \sum_{Z=H,He,O,Fe} \langle (\hat{X}_{\max,Z} - X_{\max,Z})^2 \rangle \quad (3.1)$$

$$= \sum_{Z=H,He,O,Fe} \text{Var}(\hat{X}_{\max,Z} - X_{\max,Z}) + \langle \hat{X}_{\max,Z} - X_{\max,Z} \rangle^2. \quad (3.2)$$

Here $\hat{X}_{\max,Z}$ is the predicted shower maximum and $X_{\max,Z}$ the label, for the respective element Z . $\text{Var}(\cdot)$ represents the batch-wise variance and $\langle \cdot \rangle$ the batch-wise expectation value. The network and the training is implemented using Keras [30] and the TensorFlow [31] backend.

We train our network using an Nvidia Geforce 1080 GTX GPU for 150 epochs with a batch size of 48, which takes about 60 h. We further stop the training if the objective function applied to validation data does not decrease after 10 epochs. We use the ADAM optimizer [32] with an initial learning rate of 0.001 and a decay of $2.5 \cdot 10^{-6}$. Furthermore, we reduce the learning rate by multiplying by a factor of 0.7 if the objective function evaluated on the validation data did not decrease in the recent 4 epochs.

As part of the work, a wide variety of network architectures was tested. These tests included various changes of each of the three network parts. In particular, different types of recurrent layers and various convolutional approaches were tested. The final model which utilizes LSTMs for the processing of signals and hexagonal convolutions for analyzing the spatial relations showed the best performance and stability. In addition, we found that this design is relatively robust to small changes in the architecture. A reasonable change of the hyperparameters, e.g. the number of layers or the number of features extracted by the trace network, had no significant impact on the final result.

4 Performance on simulations

In this section we study the performance of the deep neural network on the simulated data. Although none of the hadronic interaction models included in the simulations describe the measurements correctly, especially with respect to the muon deficit compared to data [33], we can at least investigate differences arising in the X_{\max} reconstruction when using different hadronic interaction models that predict different muon numbers. As latest results [34] indicate that EPOS-LHC is able to describe measurements of X_{\max} better than QGSJetII-04, in this work the network was trained using the EPOS-LHC interaction model exclusively.

In the following we first investigate the results of the deep neural network when reconstructing X_{\max} of events also simulated with the EPOS-LHC model. Next, to study the influence when




Figure 3. Correlation between X_{\max} reconstructed using the deep neural network and the true values simulated using EPOS-LHC in the energy range $19.1 < \log_{10}(E/\text{eV}) < 19.2$ for proton (a) and iron (b) showers.




Figure 4. EPOS-LHC simulation study: energy-dependent (a) bias and (b) resolution of the X_{\max} reconstruction *evaluated* on EPOS-LHC showers with the deep neural network also *trained* using EPOS-LHC showers. Different colors indicate different primaries.

modifying the hadronic interaction model, we evaluate the neural network on other hadronic interaction models than those used for the training. We use simulations based on QGSJetII-04 and Sibyll 2.3 as crosscheck in the following. We further tested the inverse setting, not presented in this work, observing that our findings are equivalent when flipping the sign of the biases. Subsequently, we discuss the combined energy and zenith dependency of the X_{\max} reconstruction of the network, indicating the phase space region that can be used for a high quality reconstruction. Finally, we investigate the reconstructed X_{\max} distributions.

4.1 Training and evaluation of the network using EPOS-LHC simulated events

In the following we show the results of the X_{\max} reconstruction with the deep neural network as trained on simulations based on the EPOS-LHC interaction model. The events for evaluating the network originate from the separated test set and were also simulated using the EPOS-LHC model. In figure 3 the event-by-event correlation of the reconstructed and true X_{\max} is shown for a benchmark bin. The Pearson correlation coefficient is above 0.5 for all elements. Comparing figure 3(b) and figure 3(a) implies that element-specific shower-to-shower fluctuations are taken into account in the deep-learning based reconstruction. This results in very different X_{\max} distributions for proton and iron showers reconstructed by the DNN.

In figure 4(a) we show the X_{\max} reconstruction bias $\mu = \langle X_{\max, \text{DNN}} - X_{\max, \text{MC}} \rangle$ as a function of the cosmic ray energy. Above 10 EeV the bias is below $\pm 9 \text{ g/cm}^2$ independent of the composition of the simulated event set used for evaluation. The statistical errors obtained via bootstrapping are mostly hidden by the markers due to the large statistics of the test set (50,000 events). Thus, the expected precision in the determination of the first moment of X_{\max} at the highest energies is excellent if the data were to look like the EPOS-LHC simulated data.

In figure 4(b) we show the event-by-event resolution σ of the X_{\max} reconstruction as a function of the cosmic ray energy. The resolution improves with increasing energy, but exhibits a clear composition dependency. This dependency is expected as showers initiated by lighter nuclei exhibit larger shower-to-shower fluctuations. At 10 EeV the X_{\max} resolution for proton-induced showers reaches 38 g/cm^2 which is below the X_{\max} fluctuations of 40 g/cm^2 as measured by the Pierre Auger Observatory [34]. For iron induced showers the X_{\max} resolution at 10 EeV is at the level of 20 g/cm^2 only.

4.2 Evaluation of the network on QGSJetII-04 and Sibyll 2.3 simulated events

In order to investigate the dependency of the deep neural network on the hadronic interaction model used for simulations, we evaluate our network trained on showers simulated using the EPOS-LHC model on simulations using QGSJetII-04 and Sibyll 2.3. In figure 5 we show the event-by-event resolution σ and bias μ of the X_{\max} reconstruction as a function of the cosmic ray energy. In figure 5(a) we show the reconstruction bias μ as a function of the cosmic ray energy for QGSJetII-04. Above 10 EeV the bias is below $\pm 10 \text{ g/cm}^2$, however with a shift of approximately -5 g/cm^2 at larger energies. The shift observed for simulations using Sibyll 2.3 (see figure 5(c)) is slightly larger and amounts to roughly -15 g/cm^2 . We assign this effect to the hadronic interaction model as it was not visible in figure 4(a). Besides the total shift, a slight energy-dependency can be observed when averaging among the compositions of Sibyll 2.3 and QGSJetII-04.

Tracing back the differences to individual characteristics of the hadronic interaction models is highly complex since the method is based on time-dependent signals. Therefore, not only the abundance of the individual shower components but also their respective time-dependent shower development needs to be considered.

The resolution of the network consistently improves with increasing energy, but exhibits a clear composition dependency which is the same for both interaction models. At 10 EeV the X_{\max} resolution for proton-induced showers reaches 38 g/cm^2 and for iron showers 20 g/cm^2 as visible




Figure 5. Energy-dependent bias (left) and resolution (right) of the X_{max} reconstruction *evaluated* on simulated showers using the hadronic interaction models QGSJetII-04 (top) and Sibyll 2.3 (bottom). The network was *trained* using EPOS-LHC showers. Different colors indicate different primaries.

in figure 5(b) and figure 5(d). These values are close to the result shown in figure 4(b), so we conclude that the resolution of X_{max} is almost independent of the hadronic interaction model.

The element-wise event-by-event correlations are very similar for all elements and energies. This is not surprising since the correlation is connected to the resolution which was found to be independent of the hadronic interaction model.

4.3 Zenith dependency of the reconstruction

To investigate the zenith dependency of the reconstruction, the evaluation of the neural network on air showers simulated with EPOS-LHC is presented. For the zenith angle in figure 6(a), we observe a moderately varying, composition-dependent reconstruction bias of up to $\pm 10 \text{ g/cm}^2$ at $\theta = 60^\circ$. Also the X_{max} resolution in figure 6(b) exhibits moderate variations with the zenith angle. For proton-induced showers they are around 30 g/cm^2 , and for iron around 20 g/cm^2 . For all elements a particular good reconstruction is visible at $\theta \approx 45^\circ$ and a worsening for angles $\theta > 55^\circ$ and $\theta < 20^\circ$. The reason for this behavior is the concurrence of larger footprints and increasing




Figure 6. EPOS-LHC simulation study: zenith-dependent (a) bias and (b) resolution of the X_{\max} reconstruction *evaluated* on EPOS-LHC showers with the deep neural network also *trained* using EPOS-LHC showers. Different colors indicate different primaries.




Figure 7. Energy- and zenith-dependent bias of the deep neural network trained and evaluated using EPOS-LHC for proton (a) and iron (b) showers.

absorption effects of the atmosphere for larger zenith angles, leading to more triggered WCDs but reducing the number of particles measured by an individual detector station. Further, for very vertical showers the shower maximum can be close to the ground or even underground which makes the reconstruction more difficult.

Note that the results shown in figure 6 are dominated by events with low energies. To verify a phase-space region in energy and zenith angle for high-quality reconstruction of X_{\max} we present in figure 7 the zenith-energy dependency of the X_{\max} bias separately for protons and iron nuclei which show the largest bias. To obtain the same number of events in each bin, we plot $\sin^2 \theta$ and use logarithmic bins in energy. It is evident that a high-quality phase space at moderate zenith angles exists already at smaller energies, its size increases with energy and allows for reconstructions

with a small bias. Outside this region for very vertical and horizontal showers, the zenith bias is particularly apparent for proton primaries, but remains mostly in the order of 15 g/cm^2 per energy bin over the complete energy range. For iron nuclei the bias is even smaller. In addition, vertical proton showers with high energies show an increased bias as for such shower geometries shower maxima below ground are more likely. These found dependencies can also be observed with showers simulated using QGSJetII-04 and Sibyll 2.3.

Overall the reconstruction of X_{max} with respect to its bias and resolution performs well on simulations. Above cosmic ray energies of 10 EeV, the expected X_{max} reconstruction bias is below $\pm 10 \text{ g/cm}^2$ and the X_{max} resolution becomes better than 35 g/cm^2 even for the lightest particles. When investigating events with saturated stations, we do not find noteworthy differences in the network performance.

Although we observe slight differences caused by using different hadronic interaction models, the network reconstruction exhibits only minor dependencies on the interaction models.

4.4 Distribution of the reconstructed shower maxima

For estimating the cosmic-ray composition the correct shape of the X_{max} distribution is essential. Due to the fact that the DNN is trained using the mean squared error, the predicted distributions are not broadened by the resolution of the method, but tend to be truncated. This bias towards the mean of the true X_{max} distribution depends on the performance and creates a dependency on X_{max} . Consequently, the quality of the estimator cannot be determined by the resolution only and is therefore examined in detail. As a result, even if the resolution for iron showers is in the range of the shower-to-shower fluctuations, a distribution with correct shape can be predicted if the estimator is sufficiently precise. Precise means in this case that the DNN assigns proton-like values of X_{max} to proton-like showers and iron-like values of X_{max} to iron-like showers etc.

To investigate the reconstructed X_{max} distribution of the network for different energies we show in figure 8 the results of the X_{max} reconstruction with the deep neural network for two example bins. In figure 8(a) the reconstructed distribution at 20 – 30 EeV for showers simulated with EPOS-LHC is shown. The solid histograms represent the distribution of reconstructed X_{max} for proton (red) and iron (blue) showers. The simulated X_{max} values are denoted by dashed histograms. It is apparent that both, the proton distribution and the iron distribution, are in good agreement. In figure 8(b) the reconstructed X_{max} distribution for showers simulated using Sibyll 2.3 at the same energy is shown. Excepting a total bias of around 15 g/cm^2 which we expect from the observed hadronic-model bias (see figure 5(c)) the simulated and reconstructed distributions match well. The successful reconstruction of the X_{max} distributions can be generalized for energies $\geq 10 \text{ EeV}$. At lower energies the reconstruction biases at low and high zenith angles (compare figure 7) superimpose the physics fluctuations. A fiducial selection could potentially reduce this effect.

The observation that X_{max} distributions similar to the true values are reproduced for all elements and hadronic interaction models (despite an absolute offset), implies that the neural network can be used to gain insights beyond the first moment of the distribution. Beyond the measurement of $\sigma(X_{\text{max}})$, this opens up possibilities for in-depth analyses of the mass-fractions of UHECRs [4].




Figure 8. Comparison of the X_{\max} distributions as reconstructed with the network (solid histograms) for proton (red) and iron (blue) showers and the simulated distribution (dashed histograms) for (a) EPOS-LHC and (b) Sibyll 2.3 for energies between 20 and 30 EeV.

5 Application to hybrid data

In this section we evaluate the performance of the deep neural network on events which include measurements of the surface detector and the fluorescence detector. First, we correct for long-term aging effects of the WCDs, which are moderate over the many years of operation, but are still noticeable in the network prediction. Then we calibrate the absolute value of the reconstructed X_{\max} using hybrid measurements. Finally, we will determine the X_{\max} resolution to verify the potential for obtaining new information about the cosmic-ray composition at the highest energies from the first two moments of the X_{\max} distribution.

5.1 Corrections for detector-aging effects

The exact shape of the signal traces has a decisive influence on the network predictions (see figure 1(b)). Besides the slightly varying response of each PMT (see section 2.4), over the long operating time of the observatory, aging effects in the signals of the WCDs occur as a combined effect of the water quality, the reflective Tyvek coating, the PMTs, and the electronics. The influence of these effects is continuously monitored by the local calibration using atmospheric muons that constantly pass through the detectors [22]. This allows to quantify measured signals in the unit of vertical equivalent muons (VEM). Every 60 s the calibration parameters Q_{VEM} and I_{VEM} are determined using the signal pulses induced by atmospheric muons. Here, I_{VEM} corresponds to the pulse-height induced by a vertical traversing muon and Q_{VEM} to the integrated pulse (bins of 25 ns).

To correct for slightly different pulse-shapes and effects of aging, the *area over peak* (A/P) variable has demonstrated its effectiveness [21]. It describes the ratio $A/P = Q_{\text{VEM}}/I_{\text{VEM}}$, and hence is a measure of the pulse shape by quantifying the relation between the signal height and decay. Due to the FADC sampling rate of 40 MHz, A/P is given in units of 25 ns. As first order approximation the area over peak is averaged for each event over all PMTs of triggered stations, which results to a characteristic and event-wise $\langle A/P \rangle$.




Figure 9. (a) Calibration of detector and aging effects. Describing the observed X_{\max} difference with respect to measured events with calibration parameters close to the simulation $(A/P)_{\text{sim}} = 3.2$, plotted versus the averaged area over peak $\langle A/P \rangle$ variable. (b) Stable long-term dependency of the average X_{\max} after the A/P calibration.

The gradual aging effects of the detectors are measurable in the detector monitoring and the predictions of the neural network. Since the beginning of the operation $\langle A/P \rangle$ decayed from roughly 3.2 to 2.95. During that time, the predicted $\langle X_{\max} \rangle$ of the DNN decayed by about 1 g/cm^2 per year. As in simulations all detectors show the same $(A/P)_{\text{sim}} = 3.2$, we calibrate the predicted $X_{\max, A/P=3.2}$ of the network with respect to events with $\langle A/P \rangle$ close to the simulated values, where we expect a high-quality reconstruction. The calibrated predictions give

$$X_{\max} = X_{\max, A/P=3.2} + b(\langle A/P \rangle), \quad (5.1)$$

where b is the observed difference. The difference shows a clear correlation with $\langle A/P \rangle$ (see figure 9(a)) and is parameterized using: $b(\langle A/P \rangle) = a \cdot (\langle A/P \rangle - 3.2)$, with the fitted calibration constant $a = 52.2 \pm 3.8 \text{ g/cm}^2$. This removes the influence of the aging effects of the water-Cherenkov detectors on the network prediction of X_{\max} as shown in figure 9(b). Due to the difference of the A/P -values in data and simulations, the average X_{\max} prediction is increased by roughly 9 g/cm^2 using the A/P calibration.

5.2 Evaluation and calibration of the first moment of the distribution of shower maxima

To further quantify the reconstructions of the DNN and allow for an absolute calibration of X_{\max} we use hybrid events which offer high-quality fluorescence measurements. In figure 10 the event-by-event correlation between the X_{\max} reconstruction of the DNN and the FD is shown. The Pearson correlation coefficient amounts to 0.63 and remains above 0.6 even if X_{\max} is corrected for the elongation rate observed by the FD [4].

In figure 11 we show the event-by-event difference $\Delta X_{\max} = X_{\max, \text{DNN}} - X_{\max, \text{FD}}$ for 3 example energy intervals as reconstructed by the deep neural network and the FD. All distributions are rather narrow and follow a Gaussian. This observation indicates that the shower-to-shower fluctuations measured with the FD, which feature a Gaussian convolved with an exponential [35], are also




Figure 10. Event-by-event correlation of the X_{\max} reconstruction of the deep neural network and their counterparts reconstructed by the FD.




Figure 11. Difference of the X_{\max} values as reconstructed with the deep neural network and the fluorescence detector for three example SD-energy intervals.

covered in the SD-based reconstruction. The observed offset of around -30 g/cm^2 indicates that the predicted X_{\max} values are too small on average. Hence, the neural network predicts shallower shower maxima.

This shift is caused by slightly different shapes of the signal traces in simulations and data. One possible reason is the shower development. In fact, a bias was already present when evaluating the DNN using showers of hadronic interaction models different than used for the training. Also the size of the observed bias is not unexpected, since already previous analyses indicated problems of these models to precisely describe the lateral and longitudinal profiles of the muon component [12, 36]. In addition, differences in the detector simulation of the signal characteristics of the WCDs compared to data can contribute to the observed bias.

With modern methods, the differences between data and simulations can be quantitatively determined and eliminated [37]. In that data-driven method the simulation is refined to look more data-like using a semi-supervised learning approach. In contrast, in this work we eliminate differences with a calibration using the independent measurements of the FD.

The energy dependency of this reconstruction bias is summarized in figure 12(a). Within

the statistics of the hybrid data used, no significant energy dependency is observed. Only the lowest energy bin at 3 EeV exhibits a slightly smaller bias. Therefore, we calibrate the X_{\max} value reconstructed by the network with the average bias as measured with the FD:

$$X_{\max}^{\text{SD}} = X_{\max, \text{DNN}} + 30.0 \text{ g/cm}^2. \quad (5.2)$$

The offset was determined by a 1-parameter fit to $30.0 \pm 0.6 \text{ g/cm}^2$ and is shown by the horizontal fitted line in figure 12(a). Even if a constant gives a rather good fit ($\chi^2/\text{ndf} = 1$), the first bins show an increased deviation from the FD measurements, which can be explained by biased reconstructions at low energies (compare figure 7). If these bins are not taken into account, the fit remains stable and moves by roughly -1 g/cm^2 .

5.3 Potential for determining the second moment of the distribution of shower maxima

In this section we will evaluate the resolution in the X_{\max} reconstruction of the deep neural network using the hybrid events.

In figure 12(b) we show the standard deviation $\sigma(\Delta X_{\max})$ of the event-wise differences between the X_{\max} value as reconstructed by the deep neural network and the X_{\max} value as measured by the FD. The symbols are located at the average energy of all events within the corresponding energy bin as indicated by the grey horizontal bars.

The vertical error bars indicate the uncertainty of $\sigma(\Delta X_{\max})$ combined for both reconstruction methods of the network and the FD. In detail, this uncertainty was calculated by a bootstrapping method using 1000 random draws from the ΔX_{\max} distribution and calculating the standard deviation σ for each set.

The energy dependency of the combined X_{\max} resolution is obtained by fitting an exponential function $\sigma_{\Delta X_{\max}}(E) = a \cdot e^{-b \cdot (\log_{10} E/\text{eV} - 18.5)} + c$ to the data, which is shown as a red curve. The values of the coefficients are $a = 18.0 \pm 2.5 \text{ g/cm}^2$, $b = 2.9 \pm 1.2$ and $c = 27.7 \pm 2.6 \text{ g/cm}^2$. The X_{\max} resolution [3] of the FD is indicated by the dashed grey curve. To obtain an estimate of the X_{\max} resolution of the deep neural network, we perform a quadratic subtraction of the FD resolution (dashed grey curve) from the combined X_{\max} resolution (solid red curve). The resulting X_{\max} resolution of the deep neural network using only the measurements of the WCDs is shown as dashed red curve in figure 12(b). We find that the resolution on the measured data reaches less than 25 g/cm^2 above 20 EeV, which is in the same order of magnitude as predicted by our simulation studies (figure 4(b)) reinforcing the finding that the resolution seems to be independent of the hadronic model. This will enable new insights into the cosmic-ray composition at high energies.

6 Summary

In this work we presented a new approach for reconstructing the maximum shower depth X_{\max} using only the signal traces of the water-Cherenkov detectors (WCDs) placed on the ground, which record a tiny subset of the billions of shower particles. It was shown that the presented method is capable of exploiting the data measured by the WCDs more comprehensively than ever before by adapting deep learning techniques, resulting in an unprecedented performance for mass composition studies using the surface detector (SD). The new method yields both the large statistics of the SD and a measurement of X_{\max} on an event-by-event basis. In consequence, our method opens up the




Figure 12. (a) Energy-dependent bias of the deep neural network with respect to the reconstruction of the fluorescence detector. (b) Energy-dependent resolution of the deep neural network with respect to the reconstruction of the fluorescence detector.

possibility to measure the abundance of mass groups of UHECRs at 100 EeV and beyond for the first time on data.

As reconstruction method, we have developed an advanced deep neural network which is especially suited for the situation of the Pierre Auger Observatory. The signal traces of the WCDs are analyzed by the network using so-called LSTM cells and their measurements are combined according to the hexagonal symmetry of the detector grid.

A key issue to correctly adjust the network parameters is the proper preparation of the data used for the network training. In addition to re-scaling and normalization of the signal amplitudes and time measurements, we implement real operation-conditions in the simulation data as data augmentation during the training. This includes missing WCDs due to hardware failures or showers falling close to the edges of the detector grid, missing signal traces of single photomultipliers and detector stations with saturated signal traces owing to high-energy events or very close shower cores. By including such effects, we make the network robust against small differences between simulation and measured data, enhancing its generalization capacities and providing an accurate reconstruction of X_{\max} for zenith angles up to 60° and even for events with saturated station electronics.

Initially we evaluate the performance of the network on simulated data. When evaluating the network using disjunct data from the same simulation as used for training, we observe an almost bias free reconstruction of X_{\max} . The X_{\max} resolution improves with increasing cosmic ray energy and is composition dependent. For proton-induced showers the resolution is 38 g/cm^2 at 10 EeV and improves to 28 g/cm^2 at the highest energies. For iron primaries the resolution is better than 20 g/cm^2 above 20 EeV.

When changing the hadronic interaction model for the evaluation of the network, we find that only the absolute bias in the X_{\max} reconstruction changes. In contrast to this shift in X_{\max} , the resolution of X_{\max} appears to be essentially independent of the hadronic interaction model. Additionally, we found that the network is able to reproduce the correct shape of the X_{\max} distribution for all elements and hadronic interaction models at energies above 10 EeV.

Finally, we test the network performance using events which include measurements of the fluorescence detector. We first eliminated effects of detector aging from long-term operation of the observatory. Compared to the results of the fluorescence measurement a remaining shift of the reconstructed X_{\max} value of the network of about 30 g/cm^2 was found. The shift is independent of the cosmic ray energy allowing for a straightforward calibration. The observed shift is larger than the constant shift of up to -15 g/cm^2 observed in the application of different interaction models. However, since all these models are not able to precisely describe the muonic component of air showers, a shift in the observed magnitude is not unexpected. Additionally, inaccuracies in the detector simulation could contribute to the observed bias.

We then estimate the X_{\max} resolution of the network on data by subtracting the X_{\max} resolution of the fluorescence detector in quadrature. We obtain a resolution of about 25 g/cm^2 above 20 EeV which is well compatible with the resolution expected from our simulation studies.

Thus, a high statistics measurement of the first moment $\langle X_{\max} \rangle$ and the second moment $\sigma(X_{\max})$ using the deep neural network reconstruction of the WCD-signal traces has a great potential to provide new insights into the cosmic-ray composition at the highest energies.

Acknowledgments

The successful installation, commissioning, and operation of the Pierre Auger Observatory would not have been possible without the strong commitment and effort from the technical and administrative staff in Malargüe. We are very grateful to the following agencies and organizations for financial support.

Argentina — Comisión Nacional de Energía Atómica; Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT); Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET); Gobierno de la Provincia de Mendoza; Municipalidad de Malargüe; NDM Holdings and Valle Las Leñas; in gratitude for their continuing cooperation over land access; Australia — the Australian Research Council; Brazil — Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq); Financiadora de Estudos e Projetos (FINEP); Fundação de Amparo à Pesquisa do Estado de Rio de Janeiro (FAPERJ); São Paulo Research Foundation (FAPESP) Grants No. 2019/10151-2, No. 2010/07359-6 and No. 1999/05404-3; Ministério da Ciência, Tecnologia, Inovações e Comunicações (MCTIC); Czech Republic — Grant No. MSMT CR LTT18004, LM2015038, LM2018102, CZ.02.1.01/0.0/0.0/16_013/0001402, CZ.02.1.01/0.0/0.0/18_046/0016010 and CZ.02.1.01/0.0/0.0/17_049/0008422; France — Centre de Calcul IN2P3/CNRS; Centre National de la Recherche Scientifique (CNRS); Conseil Régional Ile-de-France; Département Physique Nucléaire et Corpusculaire (PNC-IN2P3/CNRS); Département Sciences de l'Univers (SDU-INSU/CNRS); Institut Lagrange de Paris (ILP) Grant No. LABEX ANR-10-LABX-63 within the Investissements d'Avenir Programme Grant No. ANR-11-IDEX-0004-02; Germany — Bundesministerium für Bildung und Forschung (BMBF); Deutsche Forschungsgemeinschaft (DFG); Finanzministerium Baden-Württemberg; Helmholtz Alliance for Astroparticle Physics (HAP); Helmholtz-Gemeinschaft Deutscher Forschungszentren (HGF); Ministerium für Innovation, Wissenschaft und Forschung des Landes Nordrhein-Westfalen; Ministerium für Wissenschaft, Forschung und Kunst des Landes Baden-Württemberg; Italy — Istituto Nazionale di Fisica Nucleare (INFN); Istituto Nazionale di Astrofisica (INAF); Ministero

dell'Istruzione, dell'Università e della Ricerca (MIUR); CETEMPS Center of Excellence; Ministero degli Affari Esteri (MAE); México — Consejo Nacional de Ciencia y Tecnología (CONACYT) No. 167733; Universidad Nacional Autónoma de México (UNAM); PAPIIT DGAPA-UNAM; The Netherlands – Ministry of Education, Culture and Science; Netherlands Organisation for Scientific Research (NWO); Dutch national e-infrastructure with the support of SURF Cooperative; Poland -Ministry of Science and Higher Education, grant No. DIR/WK/2018/11; National Science Centre, Grants No. 2013/08/M/ST9/00322, No. 2016/23/B/ST9/01635 and No. HARMONIA 5–2013/10/M/ST9/00062, UMO-2016/22/M/ST9/00198; Portugal — Portuguese national funds and FEDER funds within Programa Operacional Factores de Competitividade through Fundação para a Ciência e a Tecnologia (COMPETE); Romania — Romanian Ministry of Education and Research, the Program Nucleu within MCI (PN19150201/16N/2019 and PN19060102) and project PN-III-P1-1.2-PCCDI-2017-0839/19PCCDI/2018 within PNCDI III; Slovenia — Slovenian Research Agency, grants P1-0031, P1-0385, I0-0033, N1-0111; Spain — Ministerio de Economía, Industria y Competitividad (FPA2017-85114-P and PID2019-104676GB-C32, Xunta de Galicia (ED431C 2017/07), Junta de Andalucía (SOMM17/6104/UGR, P18-FR-4314) Feder Funds, RENATA Red Nacional Temática de Astropartículas (FPA2015-68783-REDT) and María de Maeztu Unit of Excellence (MDM-2016-0692); USA — Department of Energy, Contracts No. DE-AC02-07CH11359, No. DE-FR02-04ER41300, No. DE-FG02-99ER41107 and No. DE-SC0011689; National Science Foundation, Grant No. 0450696; The Grainger Foundation; Marie Curie-IRSES/EPLANET; European Particle Physics Latin American Network; and UNESCO.

References

- [1] K.-H. Kampert and M. Unger, *Measurements of the cosmic ray composition with air shower experiments*, *Astropart. Phys.* **35** (2012) 660 [[arXiv:1201.0018](#)].
- [2] PIERRE AUGER collaboration, *The Pierre Auger cosmic ray observatory*, *Nucl. Instrum. Meth. A* **798** (2015) 172 [[arXiv:1502.01323](#)].
- [3] PIERRE AUGER collaboration, *Depth of maximum of air-shower profiles at the Pierre Auger Observatory. I. Measurements at energies above $10^{17.8}$ eV*, *Phys. Rev. D* **90** (2014) 122005.
- [4] PIERRE AUGER collaboration, *Depth of maximum of air-shower profiles at the Pierre Auger Observatory. II. Composition implications*, *Phys. Rev. D* **12** (2014) 122006.
- [5] PIERRE AUGER collaboration, *Measurement of the proton-air cross-section at $\sqrt{s} = 57$ TeV with the Pierre Auger Observatory*, *Phys. Rev. Lett.* **109** (2012) 062002 [[arXiv:1208.1520](#)].
- [6] PIERRE AUGER collaboration, *Measurement of the depth of maximum of extensive air showers above 10^{18} eV*, *Phys. Rev. Lett.* **104** (2010) 091101 [[arXiv:1002.0699](#)].
- [7] PIERRE AUGER collaboration, *Reconstruction of events recorded with the surface detector of the Pierre Auger Observatory*, *2020 JINST* **15** P10021 [[arXiv:2007.09035](#)].
- [8] M. Ave, R. Engel, M. Roth and A. Schulz, *A generalized description of the signal size in extensive air shower detectors and its applications*, *Astropart. Phys.* **87** (2017) 23.
- [9] M. Ave, M. Roth and A. Schulz, *A generalized description of the time dependent signals in extensive air shower detectors and its applications*, *Astropart. Phys.* **88** (2017) 46.

- [10] A.M. Hillas, *Angular and energy distributions of charged particles in electron photon cascades in air*, *J. Phys. G* **8** (1982) 1461.
- [11] P. Lipari, *Concepts of “age” and “universality” in cosmic ray showers*, *Phys. Rev. D* **79** (2009) 063001.
- [12] PIERRE AUGER collaboration, *Inferences on mass composition and tests of hadronic interactions from 0.3 to 100 EeV using the water-Cherenkov detectors of the Pierre Auger Observatory*, *Phys. Rev. D* **96** (2017) 122003 [[arXiv:1710.07249](#)].
- [13] C.J.T. Peixoto, *Estimating the depth of shower maximum using the surface detectors of the Pierre Auger Observatory*, *PoS(ICRC2019)*440.
- [14] M. Erdmann, J. Glombitza and D. Walz, *A deep learning-based reconstruction of cosmic ray-induced air showers*, *Astropart. Phys.* **97** (2018) 46 [[arXiv:1708.00647](#)].
- [15] D. Heck et al., *CORSIKA: A Monte Carlo code to simulate extensive air showers*, Forschungszentrum Karlsruhe Report FZKA 6019 (1998).
- [16] S. Argiro et al., *The offline software framework of the Pierre Auger Observatory*, *Nucl. Instrum. Meth. A* **580** (2007) 1485 [[arXiv:0707.1652](#)].
- [17] GEANT4 collaboration, *GEANT4 — A simulation toolkit*, *Nucl. Instrum. Meth. A* **506** (2003) 250.
- [18] T. Pierog et al., *EPOS LHC: test of collective hadronization with LHC data*, *Phys. Rev. D* **92** (2015) 034906.
- [19] S. Ostapchenko, *QGSJET-II: towards reliable description of very high energy hadronic interactions*, *Phys. Rev. D* **72** (2006) 14.
- [20] F. Riehn, H.P. Dembinski, R. Engel, A. Fedynitch, T.K. Gaisser and T. Stanev, *The hadronic interaction model SIBYLL 2.3c and Feynman scaling*, *PoS(ICRC2017)*301 [[arXiv:1709.07227](#)].
- [21] PIERRE AUGER collaboration, *The Pierre Auger Observatory IV: operation and monitoring*, talk given at the 32nd International Cosmic Ray Conference, August 11–18, Beijing, China (2011) [[arXiv:1107.4806](#)].
- [22] PIERRE AUGER collaboration, *Trigger and Aperture of the Surface Detector Array of the Pierre Auger Observatory*, *Nucl. Instrum. Meth. A* **613** (2010) 29 [[arXiv:1111.6764](#)].
- [23] Y. Le Cun et al., *Handwritten digit recognition: applications of neural network chips and automatic learning*, *IEEE Commun. Mag.* **27** (1989) 11.
- [24] E. Hoogeboom, J.W.T. Peters, T.S. Cohen and M. Welling, *HexaConv*, [arXiv:1803.02108](#).
- [25] S. Hochreiter and J. Schmidhuber, *Long short-term memory*, *Neural Comput.* **9** (1997) 1735.
- [26] G. Huang et al., *Densely connected convolutional networks*, [arXiv:1608.06993](#).
- [27] D. Clevert, T. Unterthiner and S. Hochreiter, *Fast and accurate deep network learning by Exponential Linear Units (ELUs)*, [arXiv:1511.07289](#).
- [28] K. He, X. Zhang, S. Ren and J. Sun, *Deep residual learning for image recognition*, [arXiv:1512.03385](#).
- [29] J. Glombitza, *Air-shower reconstruction at the Pierre Auger Observatory based on deep learning*, *PoS(ICRC2019)*270.
- [30] F. Chollet, *Keras: deep learning for Python*, <https://github.com/fchollet/keras> (2015).
- [31] M. Abadi et al., *TensorFlow: large-scale machine learning on heterogeneous systems* <https://www.tensorflow.org> (2015).

- [32] D.P. Kingma and J. Ba, *Adam: a method for stochastic optimization*, [arXiv:1412.6980](#).
- [33] PIERRE AUGER collaboration, *Muons in air showers at the Pierre Auger Observatory: mean number in highly inclined events*, *Phys. Rev. D* **91** (2015) 032003 [Erratum *ibid.* **91** (2015) 059901] [[arXiv:1408.1421](#)].
- [34] A. Yushkov, *Mass composition of cosmic rays with energies above $10^{17.2}$ eV from the hybrid data of the Pierre Auger Observatory*, [PoS\(ICRC2019\)482](#).
- [35] C.J. Todero Peixoto, V. de Souza and J.A. Bellido, *Comparison of the moments of the X_{\max} distribution predicted by different cosmic ray shower simulation models*, *Astropart. Phys.* **47** (2013) 18.
- [36] Pierre Auger collaboration, *Muons in air showers at the Pierre Auger Observatory: Measurement of atmospheric production depth*, *Phys. Rev. D* **90** (2014) 012012.
- [37] M. Erdmann, L. Geiger, J. Glombitza and D. Schmidt, *Generating and refining particle detector simulations using the Wasserstein distance in adversarial networks*, *Comput. Softw. Big Sci.* **2** (2018) 4.

The Pierre Auger collaboration

A. Aab⁸⁰, P. Abreu⁷², M. Aglietta^{52,50}, J.M. Albury¹², I. Allekotte¹, A. Almela^{8,11}, J. Alvarez-Muñiz⁷⁹, R. Alves Batista⁸⁰, G.A. Anastasi^{61,50}, L. Anchordoqui⁸⁷, B. Andrada⁸, S. Andringa⁷², C. Aramo⁴⁸, P.R. Araújo Ferreira⁴⁰, J.C. Arteaga Velázquez⁶⁶, H. Asorey⁸, P. Assis⁷², G. Avila¹⁰, A.M. Badescu⁷⁵, A. Bakalova³⁰, A. Balaceanu⁷³, F. Barbato^{43,44}, R.J. Barreira Luz⁷², K.H. Becker³⁶, J.A. Bellido^{12,68}, C. Berat³⁴, M.E. Bertaina^{61,50}, X. Bertou¹, P.L. Biermann^b, T. Bister⁴⁰, J. Biteau³⁵, J. Blazek³⁰, C. Bleve³⁴, M. Boháčová³⁰, D. Boncioli^{55,44}, C. Bonifazi²⁴, L. Bonneau Arbeletche¹⁹, N. Borodai⁶⁹, A.M. Botti⁸, J. Brack^d, T. Bretz⁴⁰, P.G. Brichetto Orchera⁸, F.L. Briechle⁴⁰, P. Buchholz⁴², A. Bueno⁷⁸, S. Buitink¹⁴, M. Buscemi⁴⁵, K.S. Caballero-Mora⁶⁵, L. Caccianiga^{57,47}, F. Canfora^{80,82}, I. Caracas³⁶, J.M. Carceller⁷⁸, R. Caruso^{56,45}, A. Castellina^{52,50}, F. Catalani¹⁷, G. Cataldi⁴⁶, L. Cazon⁷², M. Cerda⁹, J.A. Chinellato²⁰, K. Choi¹³, J. Chudoba³⁰, L. Chytka³¹, R.W. Clay¹², A.C. Cobos Cerutti⁷, R. Colalillo^{58,48}, A. Coleman⁹³, M.R. Coluccia⁴⁶, R. Conceição⁷², A. Condorelli^{43,44}, G. Consolati^{47,53}, F. Contreras¹⁰, F. Convenga^{54,46}, D. Correia dos Santos²⁶, C.E. Covault⁸⁵, S. Dasso^{5,3}, K. Daumiller³⁹, B.R. Dawson¹², J.A. Day¹², R.M. de Almeida²⁶, J. de Jesús^{8,39}, S.J. de Jong^{80,82}, G. De Mauro^{80,82}, J.R.T. de Mello Neto^{24,25}, I. De Mitri^{43,44}, J. de Oliveira²⁶, D. de Oliveira Franco²⁰, F. de Palma^{54,46}, V. de Souza¹⁸, E. De Vito^{54,46}, M. del Río¹⁰, O. Deligny³², A. Di Matteo⁵⁰, C. Dobrigkeit²⁰, J.C. D'Olivo⁶⁷, R.C. dos Anjos²³, M.T. Dova⁴, J. Ebr³⁰, R. Engel^{37,39}, I. Epicoco^{54,46}, M. Erdmann⁴⁰, C.O. Escobar^a, A. Etchegoyen^{8,11}, H. Falcke^{80,83,82}, J. Farmer⁹², G. Farrar⁹⁰, A.C. Fauth²⁰, N. Fazzini^a, F. Feldbusch³⁸, F. Fenu^{52,50}, B. Fick⁸⁹, J.M. Figueira⁸, A. Filipčič^{77,76}, T. Fodran⁸⁰, M.M. Freire⁶, T. Fujii^{92,e}, A. Fuster^{8,11}, C. Galea⁸⁰, C. Galelli^{57,47}, B. García⁷, A.L. Garcia Vegas⁴⁰, H. Gemmeke³⁸, F. Gesualdi^{8,39}, A. Gherghel-Lascu⁷³, P.L. Ghia³², U. Giaccari⁸⁰, M. Giammarchi⁴⁷, M. Giller⁷⁰, J. Glombitza⁴⁰, F. Gobbi⁹, F. Gollan⁸, G. Golup¹, M. Gómez Berisso¹, P.F. Gómez Vitale¹⁰, J.P. Gongora¹⁰, J.M. González¹, N. González¹³, I. Goos^{1,39}, D. Góra⁶⁹, A. Gorgi^{52,50}, M. Gottowik³⁶, T.D. Grubb¹², F. Guarino^{58,48}, G.P. Guedes²¹, E. Guido^{50,61}, S. Hahn^{39,8}, P. Hamal³⁰, M.R. Hampel⁸, P. Hansen⁴, D. Harari¹, V.M. Harvey¹², A. Haungs³⁹, T. Hebbeker⁴⁰, D. Heck³⁹, G.C. Hill¹², C. Hojvat^a, J.R. Hörandel^{80,82}, P. Horvath³¹, M. Hrabovský³¹, T. Huege^{39,14}, J. Hulsman^{8,39}, A. Insolia^{56,45}, P.G. Isar⁷⁴, P. Janecek³⁰, J.A. Johnsen⁸⁶, J. Jurysek³⁰, A. Kääpä³⁶, K.H. Kampert³⁶, B. Keilhauer³⁹, J. Kemp⁴⁰, H.O. Klages³⁹, M. Kleifges³⁸, J. Kleinfeller⁹, M. Köpke³⁷, N. Kunka³⁸, B.L. Lago¹⁶, R.G. Lang¹⁸, N. Langner⁴⁰, M.A. Leigui de Oliveira²², V. Lenok³⁹, A. Letessier-Selvon³³, I. Lhenry-Yvon³², D. Lo Presti^{56,45}, L. Lopes⁷², R. López⁶², L. Lu⁹⁴, Q. Luce³⁷, A. Lucero⁸, J.P. Lundquist⁷⁶, A. Machado Payeras²⁰, G. Mancarella^{54,46}, D. Mandat³⁰, B.C. Manning¹², J. Manshanden⁴¹, P. Mantsch^a, S. Marafico³², A.G. Mariazzi⁴, I.C. Mariş¹³, G. Marsella^{59,45}, D. Martello^{54,46}, H. Martinez¹⁸, O. Martínez Bravo⁶², M. Mastrodicasa^{55,44}, H.J. Mathes³⁹, J. Matthews⁸⁸, G. Matthiae^{60,49}, E. Mayotte³⁶, P.O. Mazur^a, G. Medina-Tanco⁶⁷, D. Melo⁸, A. Menshikov³⁸, K.-D. Merenda⁸⁶, S. Michal³¹, M.I. Micheletti⁶, L. Miramonti^{57,47}, S. Mollerach¹, F. Montanet³⁴, C. Morello^{52,50}, M. Mostafá⁹¹, A.L. Müller⁸, M.A. Muller²⁰, K. Mulrey¹⁴, R. Mussa⁵⁰, M. Muzio⁹⁰, W.M. Namasaka³⁶, A. Nasr-Esfahani³⁶, L. Nellen⁶⁷, M. Niculescu-Oglinzanu⁷³, M. Niechciol⁴², D. Nitz⁸⁹, D. Nosek²⁹, V. Novotny²⁹, L. Nožka³¹, A. Nucita^{54,46}, L.A. Núñez²⁸, M. Palatka³⁰, J. Pallotta², P. Papenbreer³⁶, G. Parente⁷⁹, A. Parra⁶², M. Pech³⁰, F. Pedreira⁷⁹, J. Pękala⁶⁹, R. Pelayo⁶⁴, J. Peña-Rodríguez²⁸, E.E. Pereira Martins^{37,8}, J. Perez Armand¹⁹, C. Pérez Bertoli^{8,39}, M. Perlin^{8,39}, L. Perrone^{54,46}, S. Petrerá^{43,44}, T. Pierog³⁹, M. Pimenta⁷², V. Pirronello^{56,45}, M. Platino⁸, B. Pont⁸⁰, M. Pothast^{82,80}, P. Privitera⁹², M. Prouza³⁰, A. Puyleart⁸⁹, S. Querchfeld³⁶, J. Rautenberg³⁶, D. Ravignani⁸, M. Reininghaus^{39,8}, J. Ridky³⁰, F. Riehn⁷², M. Risse⁴², V. Rizi^{55,44}, W. Rodrigues de Carvalho¹⁹, J. Rodriguez Rojo¹⁰, M.J. Roncoroni⁸, M. Roth³⁹, E. Roulet¹, A.C. Rovero⁵, P. Ruehl⁴², S.J. Saffi¹², A. Saftoiu⁷³, F. Salamida^{55,44}, H. Salazar⁶², G. Salina⁴⁹, J.D. Sanabria Gomez²⁸, F. Sánchez⁸, E.M. Santos¹⁹, E. Santos³⁰, F. Sarazin⁸⁶, R. Sarmiento⁷², C. Sarmiento-Cano⁸, R. Sato¹⁰, P. Savina^{54,46,32}, C.M. Schäfer³⁹, V. Scherini⁴⁶, H. Schieler³⁹, M. Schimassek^{37,8}, M. Schimp³⁶, F. Schlüter^{39,8}, D. Schmidt³⁷, O. Scholten^{81,14}, P. Schovánek³⁰, F.G. Schröder^{93,39}, S. Schröder³⁶, J. Schulte⁴⁰,

S.J. Sciutto⁴, M. Scornavacche^{8,39}, A. Segreto^{51,45}, S. Sehgal³⁶, R.C. Shellard¹⁵, G. Sigl⁴¹, G. Silli^{8,39}, O. Sima^{73,f}, R. Šmída⁹², P. Sommers⁹¹, J.F. Soriano⁸⁷, J. Souhard³⁴, R. Squartini⁹, M. Stadelmaier^{39,8}, D. Stanca⁷³, S. Stanić⁷⁶, J. Stasielak⁶⁹, P. Stassi³⁴, A. Streich^{37,8}, M. Suárez-Durán²⁸, T. Sudholz¹², T. Suomijärvi³⁵, A.D. Supanitsky⁸, J. Šupík³¹, Z. Szadkowski⁷¹, A. Taboada³⁷, A. Tapia²⁷, C. Taricco^{61,50}, C. Timmermans^{82,80}, O. Tkachenko³⁹, P. Tobiska³⁰, C.J. Todero Peixoto¹⁷, B. Tomé⁷², A. Travaini⁹, P. Travnicek³⁰, C. Trimarelli^{55,44}, M. Trini⁷⁶, M. Tueros⁴, R. Ulrich³⁹, M. Unger³⁹, L. Vaclavěk³¹, M. Vacula³¹, J.F. Valdés Galicia⁶⁷, L. Valore^{58,48}, E. Varela⁶², V. Varma K.C.^{8,39}, A. Vásquez-Ramírez²⁸, D. Veberič³⁹, C. Ventura²⁵, I.D. Vergara Quispe⁴, V. Verzi⁴⁹, J. Vicha³⁰, J. Vink⁸⁴, S. Vorobiov⁷⁶, H. Wahlberg⁴, C. Watanabe²⁴, A.A. Watson^c, M. Weber³⁸, A. Weindl³⁹, L. Wiencke⁸⁶, H. Wilczyński⁶⁹, T. Winchen¹⁴, M. Wirtz⁴⁰, D. Wittkowski³⁶, B. Wundheiler⁸, A. Yushkov³⁰, O. Zapparrata¹³, E. Zas⁷⁹, D. Zavrtnik^{76,77}, M. Zavrtnik^{77,76}, L. Zehrer⁷⁶, A. Zepeda⁶³

- ¹ *Centro Atómico Bariloche and Instituto Balseiro (CNEA-UNCuyo-CONICET), San Carlos de Bariloche, Argentina*
- ² *Centro de Investigaciones en Láseres y Aplicaciones, CITEDEF and CONICET, Villa Martelli, Argentina*
- ³ *Departamento de Física and Departamento de Ciencias de la Atmósfera y los Océanos, FCEyN, Universidad de Buenos Aires and CONICET, Buenos Aires, Argentina*
- ⁴ *IFLP, Universidad Nacional de La Plata and CONICET, La Plata, Argentina*
- ⁵ *Instituto de Astronomía y Física del Espacio (IAFE, CONICET-UBA), Buenos Aires, Argentina*
- ⁶ *Instituto de Física de Rosario (IFIR) — CONICET/U.N.R. and Facultad de Ciencias Bioquímicas y Farmacéuticas U.N.R., Rosario, Argentina*
- ⁷ *Instituto de Tecnologías en Detección y Astropartículas (CNEA, CONICET, UNSAM), and Universidad Tecnológica Nacional — Facultad Regional Mendoza (CONICET/CNEA), Mendoza, Argentina*
- ⁸ *Instituto de Tecnologías en Detección y Astropartículas (CNEA, CONICET, UNSAM), Buenos Aires, Argentina*
- ⁹ *Observatorio Pierre Auger, Malargüe, Argentina*
- ¹⁰ *Observatorio Pierre Auger and Comisión Nacional de Energía Atómica, Malargüe, Argentina*
- ¹¹ *Universidad Tecnológica Nacional — Facultad Regional Buenos Aires, Buenos Aires, Argentina*
- ¹² *University of Adelaide, Adelaide, S.A., Australia*
- ¹³ *Université Libre de Bruxelles (ULB), Brussels, Belgium*
- ¹⁴ *Vrije Universiteit Brussels, Brussels, Belgium*
- ¹⁵ *Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, RJ, Brazil*
- ¹⁶ *Centro Federal de Educação Tecnológica Celso Suckow da Fonseca, Nova Friburgo, Brazil*
- ¹⁷ *Universidade de São Paulo, Escola de Engenharia de Lorena, Lorena, SP, Brazil*
- ¹⁸ *Universidade de São Paulo, Instituto de Física de São Carlos, São Carlos, SP, Brazil*
- ¹⁹ *Universidade de São Paulo, Instituto de Física, São Paulo, SP, Brazil*
- ²⁰ *Universidade Estadual de Campinas, IFGW, Campinas, SP, Brazil*
- ²¹ *Universidade Estadual de Feira de Santana, Feira de Santana, Brazil*
- ²² *Universidade Federal do ABC, Santo André, SP, Brazil*
- ²³ *Universidade Federal do Paraná, Setor Palotina, Palotina, Brazil*
- ²⁴ *Universidade Federal do Rio de Janeiro, Instituto de Física, Rio de Janeiro, RJ, Brazil*
- ²⁵ *Universidade Federal do Rio de Janeiro (UFRJ), Observatório do Valongo, Rio de Janeiro, RJ, Brazil*
- ²⁶ *Universidade Federal Fluminense, EEIMVR, Volta Redonda, RJ, Brazil*
- ²⁷ *Universidad de Medellín, Medellín, Colombia*
- ²⁸ *Universidad Industrial de Santander, Bucaramanga, Colombia*
- ²⁹ *Charles University, Faculty of Mathematics and Physics, Institute of Particle and Nuclear Physics, Prague, Czech Republic*
- ³⁰ *Institute of Physics of the Czech Academy of Sciences, Prague, Czech Republic*
- ³¹ *Palacky University, RCPTM, Olomouc, Czech Republic*
- ³² *CNRS/IN2P3, IJCLab, Université Paris-Saclay, Orsay, France*
- ³³ *Laboratoire de Physique Nucléaire et de Hautes Energies (LPNHE), Sorbonne Université, Université de Paris, CNRS-IN2P3, Paris, France*

- 34 *Univ. Grenoble Alpes, CNRS, Grenoble Institute of Engineering Univ. Grenoble Alpes, LPSC-IN2P3, 38000 Grenoble, France*
- 35 *Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France*
- 36 *Bergische Universität Wuppertal, Department of Physics, Wuppertal, Germany*
- 37 *Karlsruhe Institute of Technology (KIT), Institute for Experimental Particle Physics, Karlsruhe, Germany*
- 38 *Karlsruhe Institute of Technology (KIT), Institut für Prozessdatenverarbeitung und Elektronik, Karlsruhe, Germany*
- 39 *Karlsruhe Institute of Technology (KIT), Institute for Astroparticle Physics, Karlsruhe, Germany*
- 40 *RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany*
- 41 *Universität Hamburg, II. Institut für Theoretische Physik, Hamburg, Germany*
- 42 *Universität Siegen, Department Physik — Experimentelle Teilchenphysik, Siegen, Germany*
- 43 *Gran Sasso Science Institute, L'Aquila, Italy*
- 44 *INFN Laboratori Nazionali del Gran Sasso, Assergi (L'Aquila), Italy*
- 45 *INFN, Sezione di Catania, Catania, Italy*
- 46 *INFN, Sezione di Lecce, Lecce, Italy*
- 47 *INFN, Sezione di Milano, Milano, Italy*
- 48 *INFN, Sezione di Napoli, Napoli, Italy*
- 49 *INFN, Sezione di Roma "Tor Vergata", Roma, Italy*
- 50 *INFN, Sezione di Torino, Torino, Italy*
- 51 *Istituto di Astrofisica Spaziale e Fisica Cosmica di Palermo (INAF), Palermo, Italy*
- 52 *Osservatorio Astrofisico di Torino (INAF), Torino, Italy*
- 53 *Politecnico di Milano, Dipartimento di Scienze e Tecnologie Aerospaziali, Milano, Italy*
- 54 *Università del Salento, Dipartimento di Matematica e Fisica "E. De Giorgi", Lecce, Italy*
- 55 *Università dell'Aquila, Dipartimento di Scienze Fisiche e Chimiche, L'Aquila, Italy*
- 56 *Università di Catania, Dipartimento di Fisica e Astronomia, Catania, Italy*
- 57 *Università di Milano, Dipartimento di Fisica, Milano, Italy*
- 58 *Università di Napoli "Federico II", Dipartimento di Fisica "Ettore Pancini", Napoli, Italy*
- 59 *Università di Palermo, Dipartimento di Fisica e Chimica "E. Segrè", Palermo, Italy*
- 60 *Università di Roma "Tor Vergata", Dipartimento di Fisica, Roma, Italy*
- 61 *Università Torino, Dipartimento di Fisica, Torino, Italy*
- 62 *Benemérita Universidad Autónoma de Puebla, Puebla, México*
- 63 *Centro de Investigación y de Estudios Avanzados del IPN (CINVESTAV), México, D.F., México*
- 64 *Unidad Profesional Interdisciplinaria en Ingeniería y Tecnologías Avanzadas del Instituto Politécnico Nacional (UPIITA-IPN), México, D.F., México*
- 65 *Universidad Autónoma de Chiapas, Tuxtla Gutiérrez, Chiapas, México*
- 66 *Universidad Michoacana de San Nicolás de Hidalgo, Morelia, Michoacán, México*
- 67 *Universidad Nacional Autónoma de México, México, D.F., México*
- 68 *Universidad Nacional de San Agustín de Arequipa, Facultad de Ciencias Naturales y Formales, Arequipa, Peru*
- 69 *Institute of Nuclear Physics PAN, Krakow, Poland*
- 70 *University of Łódź, Faculty of Astrophysics, Łódź, Poland*
- 71 *University of Łódź, Faculty of High-Energy Astrophysics, Łódź, Poland*
- 72 *Laboratório de Instrumentação e Física Experimental de Partículas — LIP and Instituto Superior Técnico — IST, Universidade de Lisboa — UL, Lisboa, Portugal*
- 73 *"Horia Hulubei" National Institute for Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- 74 *Institute of Space Science, Bucharest-Magurele, Romania*
- 75 *University Politehnica of Bucharest, Bucharest, Romania*
- 76 *Center for Astrophysics and Cosmology (CAC), University of Nova Gorica, Nova Gorica, Slovenia*
- 77 *Experimental Particle Physics Department, J. Stefan Institute, Ljubljana, Slovenia*
- 78 *Universidad de Granada and C.A.F.P.E., Granada, Spain*
- 79 *Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain*
- 80 *IMAPP, Radboud University Nijmegen, Nijmegen, The Netherlands*

- 81 *KVI — Center for Advanced Radiation Technology, University of Groningen, Groningen, The Netherlands*
- 82 *Nationaal Instituut voor Kernfysica en Hoge Energie Fysica (NIKHEF), Science Park, Amsterdam, The Netherlands*
- 83 *Stichting Astronomisch Onderzoek in Nederland (ASTRON), Dwingeloo, The Netherlands*
- 84 *Universiteit van Amsterdam, Faculty of Science, Amsterdam, The Netherlands*
- 85 *Case Western Reserve University, Cleveland, OH, U.S.A.*
- 86 *Colorado School of Mines, Golden, CO, U.S.A.*
- 87 *Department of Physics and Astronomy, Lehman College, City University of New York, Bronx, NY, U.S.A.*
- 88 *Louisiana State University, Baton Rouge, LA, U.S.A.*
- 89 *Michigan Technological University, Houghton, MI, U.S.A.*
- 90 *New York University, New York, NY, U.S.A.*
- 91 *Pennsylvania State University, University Park, PA, U.S.A.*
- 92 *University of Chicago, Enrico Fermi Institute, Chicago, IL, U.S.A.*
- 93 *University of Delaware, Department of Physics and Astronomy, Bartol Research Institute, Newark, DE, U.S.A.*
- 94 *University of Wisconsin-Madison, Department of Physics and WIPAC, Madison, WI, U.S.A.*
- (a) *Fermi National Accelerator Laboratory, Fermilab, Batavia, IL, U.S.A.*
- (b) *Max-Planck-Institut für Radioastronomie, Bonn, Germany*
- (c) *School of Physics and Astronomy, University of Leeds, Leeds, United Kingdom*
- (d) *Colorado State University, Fort Collins, CO, U.S.A.*
- (e) *now at Hakubi Center for Advanced Research and Graduate School of Science, Kyoto University, Kyoto, Japan*
- (f) *also at University of Bucharest, Physics Department, Bucharest, Romania*