



UvA-DARE (Digital Academic Repository)

Mixed partition functions and exponentially bounded edge-connection rank

Regts, G.; Sevenster, B.

DOI

[10.4171/AIHPD/100](https://doi.org/10.4171/AIHPD/100)

Publication date

2021

Document Version

Submitted manuscript

Published in

Annales de l'Institut Henri Poincaré D

[Link to publication](#)

Citation for published version (APA):

Regts, G., & Sevenster, B. (2021). Mixed partition functions and exponentially bounded edge-connection rank. *Annales de l'Institut Henri Poincaré D*, 8(2), 179-200. <https://doi.org/10.4171/AIHPD/100>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Mixed partition functions and exponentially bounded edge-connection rank

Guus Regts* Bart Sevenster †

June 16, 2020

Abstract

We study graph parameters whose associated edge-connection matrices have exponentially bounded rank growth. Our main result¹ is an explicit construction of a large class of graph parameters with this property that we call *mixed partition functions*. Mixed partition functions can be seen as a generalization of partition functions of vertex models, as introduced by de la Harpe and Jones, [P. de la Harpe, V.F.R. Jones, Graph invariants related to statistical mechanical models: examples and problems, *Journal of Combinatorial Theory, Series B* 57 (1993) 207–227] and they are related to invariant theory of orthosymplectic supergroup. We moreover show that evaluations of the characteristic polynomial of a simple graph are examples of mixed partition functions, answering a question of de la Harpe and Jones.

Keywords. partition function, graph parameter, orthogonal group, symplectic group, orthosymplectic Lie super algebra, circuit partition polynomial, connection matrix.
M.S.C [2010] Primary 05C45, 15A72; Secondary 05C25, 05C31

1 Introduction

De la Harpe and Jones [12] introduced vertex models on graphs as a generalization of the Ising-Potts model of statistical physics. Partition functions of vertex models form a rich class of graph parameters including, for instance, the number of matchings, the permanent of the adjacency matrix of a simple graph and the number of graph homomorphisms into a fixed graph [32]. Closely related concepts appear in quantum information theory as *tensor network contractions* [19], in theoretical computer science as *Holant problems* [5] and in knot theory as *Lie algebra weight systems* [7]. In the combinatorics literature a vertex model is often called an edge-coloring model, cf. [32, 18] and we adopt this terminology here.

*University of Amsterdam, Sciencepark 105-107, 1098 XH Amsterdam, the Netherlands. Email: guusregts@gmail.com. Supported by a NWO Veni grant.

†University of Amsterdam. Email: blsevenster@gmail.com. Supported by ERC grant number 339109. The present paper is partly based on this author’s PhD thesis [31].

¹Some of the results of this paper were announced in an extended abstract [25]. Unfortunately [25] contains a mistake; we will comment on that below.

Partition functions of edge coloring models Before we give the definition of partition functions of edge-coloring models, we first introduce some useful terminology. All graphs considered are finite and may have loops or multiple edges. The set of all graphs is denoted by \mathcal{G} . We will consider a single edge with an empty vertex set as a connected graph and we call this graph the *circle* and denote it by \bigcirc . For the purposes of this paper we let $\mathbb{N} = \mathbb{Z}_{\geq 0}$.

We next turn to the definition of partition function of edge-coloring models. We do this in a slightly unconventional way. This will however turn out to be convenient later. Let $k \in \mathbb{N}$ and let V_k be a vector space over \mathbb{C} with basis $\{e_1, \dots, e_k\}$. Let SV_k be the symmetric algebra of V_k . Recall that the symmetric algebra is the quotient of the tensor algebra TV_k by the ideal generated by $\{x \otimes y - y \otimes x \mid x, y \in V_k\}$. The image of $e_{i_1} \otimes \dots \otimes e_{i_n}$ in SV_k under this quotient map is denoted by $e_{i_1} \odot \dots \odot e_{i_n}$. A basis of SV_k is given by the elements $\{\odot_{i \in S} e_i\}$, where S is a multiset consisting of elements in $[k] := \{1, \dots, k\}$. An element $h \in (SV_k)^*$ is called a *k-color edge coloring model*. The *partition function* p_h of the edge coloring model h is the graph parameter over \mathbb{C} defined, for a graph $G = (V, E)$, by

$$p_h(G) := \sum_{\phi: E \rightarrow [k]} \prod_{v \in V} h(\odot_{a \in \delta(v)} e_{\phi(a)}), \quad (1)$$

where $\delta(v)$ is the set of edges incident with v . Note that $p_h(\bigcirc) = k$. If a graph parameter is the partition function of a *k-color edge coloring model* for some $k \in \mathbb{N}$, then we will refer to it as an *ordinary partition function*. A graph parameter f is called *multiplicative* if for two graphs G and H we have that $f(G \cup H) = f(G)f(H)$. We note that the partition function of an edge-coloring model h is multiplicative.

Example 1. Let us show that the number of matchings in a graph is an ordinary partition function. Take $k = 2$. Let $h \in (SV_2)^*$ be defined by $h(e_1^{\odot n_1} \odot e_2^{\odot n_2}) = 1$ if $n_2 \leq 1$ and zero otherwise. Then the set of edges mapped to 2 by a map $\phi : E \rightarrow [2]$ form a matching if and only if each vertex sees color 2 at most once. Hence these edges form a matching if and only if ϕ contributes 1 to the sum (1).

We have just defined (ordinary) partition functions for graphs, but they can also be defined for directed graphs [12], virtual link diagrams [23], chord diagrams [29], and several other combinatorial structures [30]. Instead of coloring the edges, one can also color the vertices. This then gives rise to the partition function of a *spin model* cf. [12], or *vertex coloring model* cf. [11, 18]. See Subsection 3.2 for details.

Edge connection matrices Several types of connection matrices have played an important role in characterizing partition functions [11, 32, 28, 29, 30, 22, 23]. Connection matrices also play an important role in definability of graph parameters [14]. We refer to the book of Lovász [18] and to [21] for further background.

The connection matrices we consider in this paper originate in [32]. To define them we need the concept of a fragment. For $t \in \mathbb{N}$, a *t-fragment* is a graph which has t vertices of degree one labeled $1, 2, \dots, t$. We will refer to an edge incident with a labeled vertex as an *open end*. Each open end is labeled with the label of the labeled vertex incident with it. Let \mathcal{F}_t denote the collection of all *t-fragments*. Then \mathcal{F}_0 can be considered as the collection of all graphs \mathcal{G} . For $i = 1, 2$, let F_i be a *t_i-fragment* and let o_i be an open end of F_i . The procedure of removing the labeled vertices incident to the o_i and gluing the edges o_1 and o_2 together is referred to as the *gluing* of o_1 and o_2 . This way a $t_1 + t_2 - 2$ -fragment is obtained.

Define a *gluing operation* $*$: $\mathcal{F}_t \times \mathcal{F}_t \rightarrow \mathcal{G}$, where, for two t -fragments F_1, F_2 , $F_1 * F_2$ is the graph obtained from $F_1 \cup F_2$ by gluing the open ends with the same labels together. Note that by gluing two edges of which both the endpoints are labeled one creates a circle. For a graph parameter $f : \mathcal{G} \rightarrow \mathbb{C}$, the t -th *edge-connection matrix* $M_{f,t}$ is an infinite matrix whose rows and columns are indexed by t -fragments and which is defined by, for $F_1, F_2 \in \mathcal{F}_t$,

$$M_{f,t}(F_1, F_2) := f(F_1 * F_2).$$

Edge-rank connectivity We say that a graph parameter $f : \mathcal{G} \rightarrow \mathbb{C}$ has *exponentially bounded edge-connection rank* if there exists a constant $r > 0$ such that $\text{rk}(M_{f,t}) \leq r^t$ for all $t \in \mathbb{N}$. Following [11], the minimal such r we call the *edge-rank connectivity* of f . By Fekete's Lemma this minimal r is equal to $\sup_{t \in \mathbb{Z}_+} (\text{rk}(M_{f,t}))^{1/t}$. This definition is additionally motivated from an algorithmic perspective. Using a variation of an idea of Lovász and Welsh (see [18, Theorem 6.48]) one can show that a graph parameter f with edge-rank connectivity r can be computed in time $r^{O(\Delta(G)\text{tw}(G))}(|V(G)|)^{O(1)}$ for any graph G . (Here $\Delta(G)$ denotes the maximum degree, $\text{tw}(G)$, the treewidth, see e.g. [8], and $V(G)$ the vertex set of G .)

It is not difficult to show that if f is the partition function of a k -color edge coloring model, then the edge-rank connectivity of f is at most k . In fact, Schrijver [28] characterized partition functions of edge-coloring models over \mathbb{C} using the edge-rank connectivity:

Theorem 2 ([28]). *A graph parameter $f : \mathcal{G} \rightarrow \mathbb{C}$ is the partition function of an edge-coloring model over \mathbb{C} if and only if*

$$f(\emptyset) = 1, f(\bigcirc) \in \mathbb{R} \text{ and } \text{rk}(M_{f,t}) \leq f(\bigcirc)^t \text{ for all } t \in \mathbb{N}. \quad (2)$$

In [9] an example of a graph parameter with finite edge rank connectivity was given that is not the partition function of an edge-coloring model. In [24], the authors of the present paper generalized this example by introducing skew-partition functions and showing that these have finite edge-connection rank connectivity.

Our contributions In the present paper we introduce a class of graph parameters that may be seen as a mix of ordinary partition functions and skew-partition functions, that we call *mixed partition functions*. Their definition is a little involved and will be given in Section 2. Mixed partition functions are connected to invariant theory of the orthogonal group, the symplectic group and the orthosymplectic Lie super algebra.² While we say a few words about this connection in Section 5, we will focus on combinatorial aspects of mixed partition functions in the present paper and consider the connection to invariant theory in a subsequent paper [26].

One of our main results is that mixed partition functions have finite edge-rank connectivity and we in fact conjecture that mixed partition functions form the entire class of graph parameters (that take value 1 on the empty graph) of finite edge-rank connectivity³. We shall give various examples in Section 3 of mixed partition functions. In particular we shall

²In the extended abstract [25] we incorrectly announced that the orthosymplectic Lie super algebra does not play a role.

³In [25] we in fact announced this to be true. We hope to be able to prove this conjecture in [26]. As there are quite a number of technical challenges related to supersymmetry and the fact that the orthosymplectic Lie super algebra is generally not semisimple, we choose to call it a conjecture for now.

show that a certain evaluation of the characteristic polynomial of simple graphs cannot be realized as the partition function of an edge-coloring model, answering a question of de la Harpe and Jones [12], while we show that any evaluation of the characteristic polynomial can be described as a mixed partition function.

2 Definitions

We now turn to the definition of the relevant edge-coloring models and their mixed partition functions. We first recall the construction of skew-partition functions from [24], which we modify slightly.

2.1 Skew partition functions

An *Eulerian graph* $G = (V, E)$ is a graph such that each vertex has even degree (we do not assume G to be connected). A *local pairing* κ of G is at each vertex $v \in V$ a decomposition κ_v of the edges incident with v into ordered pairs, i.e., $\kappa_v = \{(a_1, a_2), \dots, (a_{d(v)-1}, a_{d(v)})\}$. If $(a_i, a_{i+1}) \in \kappa_v$, then we say that a_i and a_{i+1} are *paired* at v . Let ω be an Eulerian orientation of the edges of G and κ a local pairing of G . We say that κ and ω are *compatible* if for each vertex v and for each $(a_1, a_2) \in \kappa_v$ the arc a_1 is incoming at v and the arc a_2 is outgoing at v under ω .

Recall that a circuit is a closed walk where vertices may occur multiple times, but edges may not. Fix an Eulerian orientation ω . A compatible local pairing κ decomposes E into circuits that, after choosing a starting point v_0 and a direction, are of the form $(v_0, e_1, \dots, e_i, v_i, e_{i+1}, \dots, v_n)$, where $v_0 = v_n$ and such that e_i and e_{i+1} are paired at v_i for each $i \in [n]$. We consider the indices modulo n . We will refer to a circuit in this decomposition as a κ -*circuit*. We define $c(\kappa)$ to be the number of circuits in this decomposition.

Let $V_{2\ell} := \mathbb{C}^{2\ell}$ and let $\{f_1, \dots, f_{2\ell}\}$ be the standard basis of $V_{2\ell}$. The *exterior algebra* $\wedge V_{2\ell}$ of $V_{2\ell}$ is the quotient of the tensor algebra $TV_{2\ell}$ by the ideal generated by $\{x \otimes y + y \otimes x \mid x, y \in V_{2\ell}\}$. The image of $f_{i_1} \otimes \dots \otimes f_{i_n}$ under this quotient map is denoted by $f_{i_1} \wedge \dots \wedge f_{i_n}$.

Define $g_i \in V_{2\ell}$ for $i = 1, \dots, 2\ell$ by

$$g_i := \begin{cases} -f_{i+\ell} & \text{if } i \leq \ell, \\ f_{i-\ell} & \text{if } i > \ell. \end{cases} \quad (3)$$

Let $G = (V, E)$ be an Eulerian graph with Eulerian orientation ω and compatible local pairing κ . For $h = (h_v)_{v \in V}$ with $h^v \in (\wedge V_{2\ell})^*$ for each $v \in V$, we define

$$s_h(G, \omega, \kappa) := (-1)^{c(\kappa)} \sum_{\phi: E \rightarrow [2\ell]} \prod_{v \in V} h^v \left(\bigwedge_{(a_1, a_2) \in \kappa_v} f_{\phi(a_1)} \wedge g_{\phi(a_2)} \right). \quad (4)$$

By skew-symmetry this is independent of the order in which we take the wedge over the elements of κ_v . We see that $s_h(G, \omega, \kappa) = 0$ if G contains a vertex of degree larger than 2ℓ , as $\wedge^n V_{2\ell} = \{0\}$ if $n > 2\ell$.

The following proposition follows immediately from [24, Proposition 1].

Proposition 3. *Let G be an Eulerian graph with Eulerian orientation ω and compatible local pairing κ and let $h = (h^v)_{v \in V}$ with $h^v \in (\wedge V_{2\ell})^*$ for each $v \in V$. Then $s_h(G, \omega, \kappa)$ is independent of the choice of ω and κ and we write $s_h(G) := s_h(G, \omega, \kappa)$.*

The proposition allows us to make the following definition.

Definition 4. Let $G = (V, E)$ be a graph. For $h = (h^v)_{v \in V}$ with $h^v \in (\wedge V_{2\ell})^*$ for each $v \in V$, the partition function p_h of h is defined as

$$p_h(G) := \begin{cases} s_h(G) & \text{if } G \text{ is Eulerian,} \\ 0 & \text{otherwise.} \end{cases} \quad (5)$$

We note that if h^v only depends on $\deg(v)$ and not on the actual vertex v , then p_h is a graph invariant, which, following [24], we call a *skew partition function*. We moreover note that $p_h(\bigcirc) = -2\ell$.

We now turn to the definition of mixed partition functions.

2.2 Mixed partition functions

For $k, \ell \in \mathbb{N}$, let V_k be a vector space of dimension k over \mathbb{C} with basis $\{e_1, \dots, e_k\}$ and let $V_{2\ell}$ be a vector space of dimension 2ℓ over \mathbb{C} with basis $\{f_1, \dots, f_{2\ell}\}$. We call any $h \in (SV_k \otimes \wedge V_{2\ell})^*$ a $(k, 2\ell)$ -color edge coloring model.

For a graph $G = (V, E)$ and $H \subseteq E$, the subgraph (V, H) of G is denoted by $G(H)$. If $G(H)$ is Eulerian, then we say that H is Eulerian. For $F \subseteq E$ Eulerian, let ω be an Eulerian orientation of the edges of $G(F)$ and let κ be a compatible local pairing of $G(F)$. For $v \in V$, let $\delta_{E \setminus F}(v)$ be the set of edges incident with v that are not in F . For $h \in (SV_k \otimes \wedge V_{2\ell})^*$, we now define

$$s_h(G, F, \omega, \kappa) := (-1)^{c(\kappa)} \sum_{\substack{\phi: F \rightarrow [2\ell] \\ \psi: E \setminus F \rightarrow [k]}} \prod_{v \in V} h \left(\bigodot_{a \in \delta_{E \setminus F}(v)} e_{\psi(a)} \otimes \bigwedge_{(a_1, a_2) \in \kappa_v} f_{\phi(a_1)} \wedge g_{\phi(a_2)} \right). \quad (6)$$

Fix one coloring $\psi : E \setminus F \rightarrow [k]$ and define for $v \in V$, $h_\psi^v \in (\wedge V_{2\ell})^*$ by

$$h_\psi^v(f_{i_1} \wedge \dots \wedge f_{i_{\deg_F(v)}}) := h \left(\bigodot_{a \in \delta_{E \setminus F}(v)} e_{\psi(a)} \otimes f_{i_1} \wedge \dots \wedge f_{i_{\deg_F(v)}} \right)$$

for $i_1, \dots, i_{\deg_F(v)} \in [2\ell]$ and let h_ψ^v evaluate to zero on $\wedge^d V_{2\ell}$ for all $d \neq \deg_F(v)$. Then we observe that

$$s_h(G, F, \omega, \kappa) = \sum_{\psi: E \setminus F \rightarrow [k]} p_{h_\psi}(G(F)),$$

and hence $s_h(G, F, \omega, \kappa)$ does not depend on the choice of ω and κ and we may define $s_h(G, F) := s_h(G, F, \omega, \kappa)$ for any choice of Eulerian orientation ω and compatible labelling κ .

Definition 5. For $h \in (SV_k \otimes \wedge V_{2\ell})^*$ the partition function of h is defined by, for a graph $G = (V, E)$,

$$p_h(G) := \sum_{\substack{F \subseteq E \\ F \text{ Eulerian}}} s_h(G, F). \quad (7)$$

We sometimes refer to the partition function just defined as a *mixed partition function* so as to distinguish it from ordinary partition functions and skew partition functions. We note that $p_h(\bigcirc) = k - 2\ell$.

We can now state our main theorem concerning the edge-rank connectivity of mixed partition function.

Theorem 6. *If f is the partition function of an element $h \in (SV_k \otimes \wedge V_{2\ell})^*$, then*

$$\text{rk}(M_{f,t}) \leq (k + 2\ell)^t \text{ for each } t \in \mathbb{N}. \quad (8)$$

We will prove this theorem in Section 4. First we give several examples of mixed partition functions in the next section.

3 Examples of mixed partition functions

We start with some basic examples.

3.1 Basic examples

Example 7. Let h be a $(k,0)$ -color edge coloring model. For a graph $G = (V, E)$ and $F \subseteq E$, we have that $s_h(G, F) = 0$ unless $F = \emptyset$. So $p_h(G)$ reduces to

$$s_h(G, \emptyset) = \sum_{\psi: E \rightarrow [k]} \prod_{v \in V} h(\bigodot_{a \in \delta(v)} e_{\psi(a)}).$$

So we see that p_h is an ordinary partition function as in (1). We similarly see that if h is a $(0, 2\ell)$ -color edge coloring model that p_h reduces to $s_h(G, E)$. So p_h is a skew partition function as in (5).

Example 8. If $h_0 \in (SV_k)^*$ and $h_1 \in (\wedge V_{2\ell})^*$, then let $h = h_0 \otimes h_1 \in (SV_k \otimes \wedge V_{2\ell})^*$. For a graph $G = (V, E)$ and $F \subseteq E$ Eulerian, it follows directly from (6) that $s_h(G, F) = p_{h_0}(G(E \setminus F))p_{h_1}(G(F))$. So we find that

$$p_h(G) = \sum_{\substack{F \subseteq E \\ F \text{ Eulerian}}} p_{h_0}(G(E \setminus F))p_{h_1}(G(F)). \quad (9)$$

We now move to a few more involved examples of mixed partition functions.

3.2 The characteristic polynomial

In this subsection we assume that our graphs do not have the circle, \bigcirc , as a connected component. The *adjacency matrix* A of G is the $V \times V$ matrix such that for $i, j \in V$ with $i \neq j$, $A(i, j)$ is the multiplicity of the edge $\{i, j\}$ in E and such that for $i \in V$, $A(i, i)$ is twice the number of loops at the vertex i . The *characteristic polynomial* $p(G)$ of G is defined as $p(G; t) := \det(tI - A)$. We show below that for each t there exists an $(2, 2)$ -edge-coloring model $h(t)$ such that $p(G; t) = p_{h(t)}(G)$ for all graphs G , see Proposition 10 below. First we turn to a question of de la Harpe and Jones concerning spin models and the characteristic polynomial. To this end give the definition of the partition function of a spin model. We follow the definition of de la Harpe and Jones [12].

Let $n \in \mathbb{N}$ and let B be a symmetric $n \times n$ matrix with values in some commutative ring R . We call B a *spin model over R* . The partition function, p_B , of the spin model B is defined, for a graph $G = (V, E)$, by

$$p_B(G) := \sum_{\kappa: V \rightarrow [n]} \prod_{\{v_1, v_2\} \in E} B(\kappa(v_1), \kappa(v_2)). \quad (10)$$

Note that this is well-defined as B is a symmetric matrix.

De la Harpe and Jones [12, Problem 1] asked about the existence of a spin model B over $\mathbb{C}[t]$, such that $p_B(G) = p(G; t)$ for each graph G . In the following proposition we shall show that the answer to this question is negative. In fact, we show something stronger.

Proposition 9. *There does not exist an edge coloring model h such that $p_h(G) = p(G; 0)$ for all graphs G .*

This proposition is indeed stronger than we need, since, by a result of Szegedy [32], the partition function of any spin model is equal to the partition function of an ordinary edge coloring model and hence Proposition 9 rules out the existence of a spin model of which the partition function equals the characteristic polynomial evaluated at 0 and therefore provides a negative answer to the question of de la Harpe and Jones. Proposition 10 below may serve as an alternative answer to the question of de la Harpe and Jones.

We now turn to a proof of Proposition 9.

Proof of Proposition 9. Let us abuse notation and write $\det(G)$ for the determinant of the adjacency matrix of G . Note that for a graph with an even number of vertices we have $p(G; 0) = \det(G)$. We will make use of the characterization of partition functions of edge coloring models as given in [9]. Fix k and consider the graph G consisting of $k + 1$ copies of the 6-cycle C_6 . Direct one edge in each cycle and label the endpoints of these arcs 1 up to $k + 1$. For a permutation $\pi \in S_{k+1}$, denote by G_π the graph obtained from G by letting π permute the endpoints of the directed edges. Note that if the permutation π can be written as the product of disjoint cycles π_1, \dots, π_t , then G_π is the graph consisting of t cycles, of length $6|\pi_1|, \dots, 6|\pi_t|$ respectively. Here $|\pi_i|$ denotes the length of the cycle π_i ; we include cycles of length 1. If $p(G; 0)$ is the partition function of a k -color edge coloring model, then, by [9, Theorem 1], it must satisfy

$$\sum_{\pi \in S_{k+1}} \operatorname{sgn}(\pi) p(G_\pi; 0) = 0. \quad (11)$$

It follows from, for example, [2, Section 1.4.3], that $\det(C_k) = 0$ if $k = 0 \pmod{4}$ and $\det(C_k) < 0$ if $k = 2 \pmod{4}$. This implies that for $\det(G_\pi)$ to be non-zero none of the cycles π_1, \dots, π_t may be of even length. However, if all cycles in the cycle decomposition of π are of odd length, then the parity of the number of these cycles is equal to the parity of $k + 1$. So in this case, $\det(G_\pi)$ is strictly positive if this parity is even and strictly negative if this parity is odd for all such permutations π . As all cycles of π are odd we have $\operatorname{sgn}(\pi) = 1$. Since $\det(G_\pi) = p(G_\pi; 0)$ for all π , this shows that $\sum_{\pi \in S_{k+1}} \operatorname{sgn}(\pi) p(G_\pi; 0)$ is either strictly positive or strictly negative. So it is non-zero. So we conclude that (11) is violated and hence that $p(\cdot; 0)$ cannot be the partition function of any edge coloring model. \square

Proposition 10. *For each $t \in \mathbb{C}$, there exists a $(2, 2)$ -color edge coloring model $h(t)$ such that $p_{h(t)}(G) = p(G; t)$ for all graphs G .*

Proof. Using the Leibniz expansion of the determinant, Sachs [27] gave an expression of the characteristic polynomial of a graph G in terms of certain subgraphs of G . The expression extends to graphs with multiple edges and loops. Let $G = (V, E)$ be a graph. Let \mathcal{H} be the set of $H \subseteq E$ such that each connected component of $G(H)$ is either a vertex, an edge or a cycle. For $H \in \mathcal{H}$, let $e^*(H)$ and $c(H)$ denote the number of connected components of

$G(H)$ that are edges and cycles respectively. Let $V[H] \subseteq V$ be the set of vertices of G that are incident with an edge of H . Then Sachs showed that

$$p(G; t) = \sum_{H \in \mathcal{H}} (-1)^{e^*(H)} (-2)^{c(H)} t^{|V| - |V[H]|}. \quad (12)$$

We now give a $(2, 2)$ -color edge coloring model $h = h(t)$ such that $p_h(G) = p(G; t)$ for each $t \in \mathbb{C}$ and graph G . Let h be defined as follows:

$$\begin{aligned} h(e_1^{\odot i} \otimes f_1 \wedge g_1) &= 1 \text{ for } i \in \mathbb{N}, \\ h(e_1^{\odot i} \odot e_2) &= \sqrt{-1} \text{ for } i \in \mathbb{N}, \\ h(e_1^{\odot i}) &= t \text{ for } i \in \mathbb{N}, \end{aligned}$$

and let h evaluate to 0 on basis elements of $SV_2 \otimes \wedge V_2$ that are not in the span of these elements. Now let $F \subseteq E$ be Eulerian. We compute $s_h(G, F)$. If $G(F)$ has a vertex that is not of degree 0 or 2, then $s_h(G, F) = 0$.

So let us assume that each vertex of $G(F)$ has degree 0 or 2. Let ω be an Eulerian orientation of F with a compatible local pairing κ of $G(F)$. Now let $\phi : F \rightarrow [2]$ and $\psi : E \setminus F \rightarrow [2]$. We first note that for the contribution of ϕ and ψ to $s_h(G, F, \omega, \kappa)$ to be non-zero, we need $\psi^{-1}(2)$ to be a matching in G that is not incident with any edge in F .

Now fix $\psi : E \setminus F \rightarrow [2]$ such that $\psi^{-1}(2)$ is a matching in G that is not incident with any edge in F . Let $H = F \cup \psi^{-1}(2) \subseteq E$. Note that at each vertex $v \in V$ that is not incident with H , we see $h(e_1^{\odot d(v)}) = t$. There are $|V| - |V[H]|$ such vertices v . If $v, u \in V$ are two vertices such that $\{u, v\}$ is an isolated edge of (V, H) , then at u we see $h(e_1^{\odot d(u)-1} \odot e_2) = \sqrt{-1}$ and at v we see $h(e_1^{\odot d(v)-1} \odot e_2) = \sqrt{-1}$. So two vertices u, v such that $\{u, v\}$ is an isolated edge of (V, H) contribute -1 to the partition function.

Now consider the colorings $\phi : F \rightarrow [2]$. Such a ϕ gives a non-zero contribution if and only if it is constant on the edges of each 2-regular connected component of (V, F) . So there are exactly $2^{c(F)}$ colorings $\phi : F \rightarrow [2]$ that have a non-zero contribution. As $c(\kappa) = c(F) = c(H)$, we see that the total contribution to the partition function of these colorings is exactly

$$(-1)^{e^*(H)} (-2)^{c(H)} t^{|V| - |V[H]|}. \quad (13)$$

Now summing over all F and corresponding ϕ and ψ , we find that $p_h(G)$ is indeed equal to $p(G; t)$ by (12). \square

3.3 Evaluations of the circuit partition polynomial

The circuit partition polynomial, introduced, in a slightly different form, by Martin in his thesis [20], is related to Eulerian walks in graphs and to the Tutte polynomial of planar graphs. Several identities for the circuit partition polynomial were established by Bollobás [1] and Ellis-Monaghan [10].

We say that two circuits are equivalent if one can be obtained from the other by possibly changing the starting vertex or the direction of the walk. For a graph $G = (V, E)$, let $X(G)$ be a set of representatives of this equivalence relation. Let $\mathcal{C}(G)$ be the collection of all partitions of E into circuits in $X(G)$. For $C \in \mathcal{C}(G)$, let $|C|$ be the number of circuits in the partition.

The circuit partition polynomial $J(G, x)$ is defined for a graph G by

$$J(G, x) := \sum_{C \in \mathcal{C}(G)} x^{|C|}.$$

So if G is not an Eulerian graph, then $J(G, x) = 0$. We clearly have that $J(G \cup H, x) = J(G, x)J(H, x)$ for two graphs G and H and it is natural to define $J(\bigcirc, x) = x$. We shall show that every integer evaluation of the circuit partition polynomial can be expressed as a mixed partition function. We first recall the result from [24] that positive and negative even integer evaluations can be expressed as ordinary partition functions and skew partition functions respectively, after which we show that odd negative evaluations can be expressed as mixed partition functions (with both k and 2ℓ positive).

For $k \in \mathbb{N}$, it was shown in [1, 10] that $J(G, k)$ can be expressed as

$$J(G, k) = \sum_A \prod_{v \in V} \prod_{i=1}^k (\deg_{A_i}(v) - 1)!!, \quad (14)$$

where A ranges over ordered partitions of E into k subsets A_1, \dots, A_k such that A_i is Eulerian for all $i \in [k]$. We use the convention that for a nonnegative integer d , $(2d - 1)!! = (2d - 1) \cdot (2d - 3) \cdots 3 \cdot 1$ and $(2d)!! = 0$.

We express (14) as the partition function of $h_0 \in (SV_k)^*$ as follows. For $(\alpha_1, \dots, \alpha_k) \in \mathbb{N}^k$, we set

$$h_0(\bigodot_{i \in [k]} e_i^{\odot \alpha_i}) := \prod_{i=1}^k (\alpha_i - 1)!!. \quad (15)$$

Using (14) it is not difficult to see that $p_{h_0}(G) = J(G, k)$ for each graph G , cf. [24].

Bollobás [1] showed that the evaluation of the circuit partition polynomial $J(G, x)$ of a graph at negative even integers -2ℓ can be expressed as

$$J(G, -2\ell) = \sum_{H_1, \dots, H_\ell} (-2)^{\sum_{i=1}^{\ell} c(H_i)}, \quad (16)$$

where this sum runs over all ordered partitions H_1, \dots, H_ℓ of the edge set of G such that for each $i \in [\ell]$ each vertex in (V, H_i) has degree 0 or degree 2 and where $c(H_i)$ is the number of 2-regular connected components of (V, H_i) . Now let $h_1 \in (\wedge V_{2\ell})^*$ be defined, for $S \subseteq [\ell]$, by

$$h_1(\bigwedge_{i \in S} f_i \wedge g_i) = 1; \quad (17)$$

h_1 evaluates to zero on all other basis elements not in the span of the elements above. In [24] it is shown, using (16), that $p_{h_1}(G) = J(G, -2\ell)$.

We will next show that mixed partition functions can also express evaluations of the circuit partition polynomial at negative odd integers. In [10], Ellis-Monaghan showed for a graph $G = (V, E)$ that

$$J(G, x + y) = \sum_{A \subseteq E} J(G(A), x) J(G(E \setminus A), y). \quad (18)$$

Now, for a negative odd integer $-2\ell + 1$, let $h_0 \in (SV_1)^*$ correspond to $k = 1$ in (15) and let $h_1 \in (\wedge V_{2\ell})^*$ be as in (17). Let $h = h_0 \otimes h_1 \in (SV_1 \otimes \wedge V_{2\ell})^*$. Then by (9) and (18) we find that $p_h(G) = J(G, -2\ell + 1)$, giving us an expression of $J(G, -2\ell + 1)$ as a mixed partition function.

Remark 1. Let us finally remark that the cycle partition polynomial evaluated at $x \notin \mathbb{Z}$ cannot be realized as the mixed partition function of any edge-coloring model. This follows from Theorem 6 and the fact that by [28, Proposition 2] the rank of the submatrix of $M_{J(\cdot, x), 2n}$, indexed by fragments in which each component consists of an unlabeled vertex with two open ends incident with it, has rank at least $n!$, cf. [24].

4 The edge-rank connectivity of mixed partition functions

In this section we prove Theorem 6.

We first show a lemma on matchings that will be useful later on. If M and N are directed perfect matchings on the same vertex set, then we denote by $o(M \cup N)$ the parity of the number of arcs in $M \cup N$ that need to be flipped to make $M \cup N$ into an Eulerian digraph. Note that we call a digraph Eulerian if for each vertex its in degree is equal to its out degree. Since each cycle in $M \cup N$ has even length this is well defined. Recall that $c(M \cup N)$ is the number of connected components of $M \cup N$.

Lemma 11. *Let M and N be two directed perfect matchings on $[2m]$ for $m \in \mathbb{N}$. Then the sign of any permutation in S_{2m} that sends N to M is equal to $(-1)^{c(M \cup N) + o(M \cup N)}$.*

Proof. Note that all permutations that send M to N have the same sign, as all permutations in S_{2m} that stabilize M have trivial sign. We may assume that $M \cup N$ consists of a single connected component. Let $\sigma_1, \sigma_2 \in S_{2m}$ be permutations that flip arcs of M and N respectively, such that $\sigma_1 M \cup \sigma_2 N$ has an Eulerian orientation. If the vertices of the cycle are given by v_1, v_2, \dots, v_{2m} in cyclic order, then the permutation $\tau = (v_1 v_2 \dots v_{2m})$ has the property that $\tau \sigma_1 M = \sigma_2 N$. As $2m$ is even, the sign of τ is -1 . So the permutation $\sigma_2^{-1} \tau \sigma_1$ sends M to N . Per construction we have that $\text{sgn}(\sigma_1) \text{sgn}(\sigma_2) = (-1)^{o(M \cup N)}$. This proves the lemma. \square

In the proof we will make use of some linear algebra that we will now define. Let (\cdot, \cdot) be the nondegenerate symmetric bilinear form on $V_k = \mathbb{C}^k$ given by $(x, y) := x^T y$ for $x, y \in \mathbb{C}^k$. Let $\langle \cdot, \cdot \rangle$ be the nondegenerate skew-symmetric bilinear form on $V_{2\ell} = \mathbb{C}^{2\ell}$ given by $\langle x, y \rangle = x^T J y$ for $x, y \in \mathbb{C}^{2\ell}$, where J is the $2\ell \times 2\ell$ matrix $\begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}$, with I the $\ell \times \ell$ identity matrix. We define $V_{k, 2\ell} := V_k \oplus V_{2\ell}$. We write an element w of $V_{k, 2\ell}$ as $w_0 + w_1$, where $w_0 \in V_k$ and $w_1 \in V_{2\ell}$. We equip this space with a nondegenerate bilinear form $[\cdot, \cdot]$ defined by

$$[x, y] := (x_0, y_0) + \langle x_1, y_1 \rangle, \quad (19)$$

for $x, y \in V_{k, 2\ell}$. We note that this form is often called a *super symmetric bilinear form*, cf. [6].

Proof of Theorem 6. Our goal is to show that for each $t \in \mathbb{N}$, we can write $M_{f, t}$ as a Gram matrix of vectors in $V_{k, 2\ell}^{\otimes t}$ with respect to the bilinear form $[\cdot, \cdot]$. This then immediately implies Theorem 6.

Let $t \in \mathbb{N}$ and let $F = (V, E)$ be a t -fragment. Recall that a t -fragment is a graph with t vertices of degree 1 labeled $1, \dots, t$. The set of unlabeled vertices of F is denoted by $V'(F)$. A subset $H \subseteq E$ is called *Eulerian* if the degree of each unlabeled vertex in $F(H)$ is even. Let $H \subseteq E$ be Eulerian. Let $S(H)$ be the set of labeled vertices incident with an edge in H . If H is chosen, we refer to $S(H)$ as S . Note that $|S|$ is even because H is Eulerian. We identify

the labeled vertices with $[t]$ according to the labeling. Through this identification we view S as a subset of $[t]$.

We now extend some of the definitions we gave for graphs to fragments. An *Eulerian orientation* ω of H is an orientation of the edges of H such that in $F(H)$, at each unlabeled vertex the number of incoming arcs is equal to the number of outgoing arcs. A *local pairing* κ of $F(H)$ is an assignment κ to each $v \in V'(F)$ of a decomposition κ_v of the edges in H incident with v into ordered pairs. The local pairing κ is called *compatible* with a Eulerian orientation ω if for each $v \in V'(F)$ and for each $(a_1, a_2) \in \kappa_v$ the arc a_1 is incoming under ω and the arc a_2 is outgoing under ω .

Now let κ be a local pairing of $F(H)$ compatible with an Eulerian orientation ω of H . Note that κ partitions the edge set of H into circuits and directed trails that begin and end in labeled vertices. We call this decomposition the κ -*decomposition* of H . Let $\hat{c}(\kappa)$ be the number of circuits in the κ -decomposition. Define $M(\omega, \kappa)$ to be the directed perfect matching on $S \subseteq [t]$ such that (i, j) is an arc of $M(\omega, \kappa)$ if there is a directed trail in the κ -decomposition from i to j . Write $S = \{i_1, \dots, i_{|S|}\}$ with $i_1 < \dots < i_{|S|}$. Let $\text{sgn}(M(\omega, \kappa))$ be the sign of a permutation that sends $M(\omega, \kappa)$ to the directed perfect matching with arcs $(i_1, i_2), \dots, (i_{|S|-1}, i_{|S|})$. This is well-defined by Lemma 11.

Let $\chi = (\chi_0, \chi_1)$ with $\chi_0 : [t] \setminus S \rightarrow [k]$ and $\chi_1 : S \rightarrow [2\ell]$. Such a pair $\chi = (\chi_0, \chi_1)$ is called *consistent* with S . We say that a coloring $\psi : E \setminus H \rightarrow [k]$ extends χ_0 if, for each $i \in [t] \setminus S$, we have $\chi_0(i) = \psi(a)$, where $a \in E \setminus H$ is the edge incident with i . We denote this by $\psi \sim \chi_0$. Similarly, we say that $\phi : H \rightarrow [2\ell]$ extends χ_1 if, for each $i \in S$, we have $\chi_1(i) = \phi(a)$, where $a \in H$ is the edge incident with i . Again, we denote this by $\phi \sim \chi_1$.

For $i \in [t] \setminus S$, let $c_{\chi, \omega, i} = e_{\chi_0(i)}$, and for $i \in S$, let $c_{\chi, \omega, i} = f_{\chi_1(i)}$ if the edge incident with i is incoming at i under ω and let $c_{\chi, \omega, i} = g_{\chi_1(i)}$ if the edge incident with i is outgoing at i under ω . We define the tensor $t'_{h, \chi}(F, H, \omega, \kappa)$ in $V_{k, 2\ell}^{\otimes t}$ by

$$t'_{h, \chi}(F, H, \omega, \kappa) := (-1)^{\hat{c}(\kappa)} \sum_{\substack{\psi \sim \chi_0 \\ \phi \sim \chi_1}} \prod_{v \in V'(F)} h \left(\bigodot_{a \in \delta_{E \setminus H}(v)} e_{\psi(a)} \otimes \bigwedge_{(a_1, a_2) \in \kappa_v} f_{\phi(a_1)} \wedge g_{\phi(a_2)} \right) \bigotimes_{i \in [t]} c_{\chi, \omega, i}$$

where the sum runs over all $\psi : E \setminus H \rightarrow [k]$ with $\psi \sim \chi_0$ and all $\phi : H \rightarrow [2\ell]$ with $\phi \sim \chi_1$. We define

$$t'_h(F, H, \omega, \kappa) := \sum_{\substack{\chi \text{ consistent} \\ \text{with } S}} t'_{h, \chi}(F, H, \omega, \kappa),$$

and finally we define

$$t_h(F, H, \omega, \kappa) := (-1)^{|S|/4} \text{sgn}(M(\omega, \kappa)) t'_h(F, H, \omega, \kappa).$$

We first make an important observation. Let ω' be obtained from ω by inverting the arcs in a directed trail P in the κ -decomposition and let κ' be obtained from κ by inverting all the pairings in the directed trail P (hence κ' is compatible with ω'). Note that $\text{sgn}(M(\omega, \kappa)) = -\text{sgn}(M(\omega', \kappa'))$, as $M(\omega', \kappa')$ is obtained from $M(\omega, \kappa)$ by inverting the direction of one arc. The total number of pairings and arcs in the directed trail P is odd. So similar to what we have seen in the proof of [24, Lemma 2], we find that $t'_h(F, H, \omega, \kappa) = -t'_h(F, H, \omega', \kappa')$. This shows that

$$t_h(F, H, \omega, \kappa) = t_h(F, H, \omega', \kappa'). \quad (20)$$

Now let $F_1 = (V_1, E_1)$ and $F_2 = (V_2, E_2)$ be two t -fragments with Eulerian subsets $H_1 \subseteq E_1$ and $H_2 \subseteq E_2$ such that $S(H_1) = S(H_2) = S$. Let $G = (V, E) = F_1 * F_2$. Note that H_1 and H_2 induce an Eulerian subset of E . We denote this set by $H_1 * H_2$. For $i = 1, 2$, let ω_i be an Eulerian orientation of H_i with a compatible local pairing κ_i of $F_i(H_i)$. We next show that

$$[t_h(F_1, H_1, \omega_1, \kappa_1), t_h(F_2, H_2, \omega_2, \kappa_2)] = s_h(G, H_1 * H_2). \quad (21)$$

By (20) we may assume that $\omega_1, \kappa_1, \omega_2$ and κ_2 are chosen in such a way that $(S, M(\omega_1, \kappa_1) \cup M(\omega_2, \kappa_2))$ is an Eulerian digraph. By Lemma 11 we see that

$$\text{sgn}(M(\omega_1, \kappa_1))\text{sgn}(M(\omega_2, \kappa_2)) = (-1)^{c(M(\omega_1, \kappa_1) \cup M(\omega_2, \kappa_2))},$$

as $o(M(\omega_1, \kappa_1) \cup M(\omega_2, \kappa_2)) = 0$. Furthermore, ω_1 and ω_2 induce an Eulerian orientation ω of $H_1 * H_2$ and the local pairing κ of $G(H_1 * H_2)$ induced by κ_1 and κ_2 is compatible with ω . So we find that

$$\text{sgn}(M(\omega_1, \kappa_1))\text{sgn}(M(\omega_2, \kappa_2))(-1)^{\hat{c}(\kappa_1)}(-1)^{\hat{c}(\kappa_2)} = (-1)^{c(\kappa)}. \quad (22)$$

Now let $\chi = (\chi_0, \chi_1)$ and $\chi' = (\chi'_0, \chi'_1)$ both be consistent with S . We consider

$$[t'_{h, \chi}(F_1, H_1, \omega_1, \kappa_1), t'_{h, \chi'}(F_2, H_2, \omega_2, \kappa_2)]. \quad (23)$$

Note that this is equal to 0 if χ_0 and χ'_0 do not agree. Furthermore, as the orientations of ω_1 and ω_2 are opposite at a labeled vertex in S , we see that χ_1 and χ'_1 also have to agree for (23) to be non-zero. So let us assume that $\chi = \chi'$. Note that as the orientation ω is Eulerian, at half of the vertices in S the arc of H_1 is incoming and the arc of H_2 is outgoing. So at such a vertex i the bilinear form becomes $\langle f_{\chi_1(i)}, g_{\chi_1(i)} \rangle = -1$. At the other half of the vertices in S the arc of H_2 is incoming and the arc of H_1 is outgoing. So at such a vertex i the bilinear form becomes $\langle g_{\chi_1(i)}, f_{\chi_1(i)} \rangle = 1$. These contributions cancel with $(-1)^{|S(H_1)|/4}(-1)^{|S(H_2)|/4}$. Together with (22) this shows (21).

Now, for $i = 1, 2$, let $H_i \subseteq E_i$ and let ω_i be an Eulerian orientation of H_i with a compatible local pairing κ_i of $F_i(H_i)$. Suppose that $S(H_1) \neq S(H_2)$. Then it follows that

$$[t_h(F_1, H_1, \omega_1, \kappa_1), t_h(F_2, H_2, \omega_2, \kappa_2)] = 0, \quad (24)$$

because at i in the symmetric difference of $S(H_1)$ and $S(H_2)$ there occurs an element of V_k at one side of the bilinear form and an element of $V_{2\ell}$ at the other side.

Note that as H_1 and H_2 run over all Eulerian subsets of F_1 and F_2 , we have that $H_1 * H_2$ runs over all Eulerian subsets of G . So it follows from (21) and (24) that

$$\left[\sum_{\substack{H_1 \subseteq E_1 \\ H_1 \text{ Eulerian}}} t_h(F_1, H_1, \omega_1, \kappa_1), \sum_{\substack{H_2 \subseteq E_2 \\ H_2 \text{ Eulerian}}} t_h(F_2, H_2, \omega_2, \kappa_2) \right] = \quad (25)$$

$$\sum_{\substack{H \subseteq E \\ H \text{ Eulerian}}} s_h(G, H, \omega, \kappa) = f(G), \quad (26)$$

where, for $i = 1, 2$, κ_i is a local pairing of $F_i(H_i)$ compatible with an Eulerian orientation ω_i of H_i . This shows that $M_{f,t}$ indeed is the Gram matrix of a set of vectors in $V_{k,2\ell}^{\otimes t}$ with respect to the bilinear form $[\cdot, \cdot]$. So the rank of $M_{f,t}$ is bounded by $(k + 2\ell)^t$. This proves Theorem 6. \square

5 Connections to invariant theory

In this section we will indicate how one might prove the conjectured converse to Theorem 6 by saying how mixed partition functions connect to the invariant theory of the orthogonal and symplectic groups and the orthosymplectic Lie superalgebra.

Following [9, 28, 24], to prove a converse to Theorem 6 one essentially needs to prove two statements. First one needs an algebraic characterization of mixed partition functions. Second one needs to show that a multiplicative graph parameter with finite edge-rank connectivity satisfies these algebraic conditions. Below we comment on how such a possible algebraic characterization is deeply connected to invariant theory.

For $k, \ell \in \mathbb{N}$, recall that V_k is a vector space of dimension k over \mathbb{C} with basis $\{e_1, \dots, e_k\}$ and recall that $V_{2\ell}$ is a vector space over \mathbb{C} of dimension 2ℓ with basis $\{f_1, \dots, f_{2\ell}\}$. We furthermore defined $V_{k,2\ell}$ as $V_k \oplus V_{2\ell}$. The *orthogonal group* O_k is the group of $k \times k$ matrices that preserve the symmetric bilinear form; i.e., for $g \in \mathbb{C}^{k \times k}$, $g \in O_k$ if and only if $\langle gx, gy \rangle = \langle x, y \rangle$ for all $x, y \in V_k$. The *symplectic group* $Sp_{2\ell}$ is the group of $2\ell \times 2\ell$ matrices that preserve the skew-symmetric bilinear form; i.e., for $g \in \mathbb{C}^{2\ell \times 2\ell}$, $g \in Sp_{2\ell}$ if and only if $\langle gx, gy \rangle = \langle x, y \rangle$ for all $x, y \in V_{2\ell}$. The group of $(k + 2\ell) \times (k + 2\ell)$ -matrices that preserve the form (19) can be shown to be the direct product of O_k and $Sp_{2\ell}$.

Consider a basis element $f = f_{i_1} \wedge \dots \wedge f_{i_t} \in \wedge V_{2\ell}$. We call f even if t is even and odd otherwise. Let us for $x \in \mathbb{Z}/2\mathbb{Z} = \{\bar{0}, \bar{1}\}$ denote by $(\wedge V_{2\ell})_x$ the subspace of $\wedge V_{2\ell}$ spanned by the basis elements $f_{i_1} \wedge \dots \wedge f_{i_t}$ for which $x = t \pmod 2$. We define

$$R = R(V_{k,2\ell}) := \text{Sym}(SV_k \otimes (\wedge V_{2\ell})_{\bar{0}}).$$

Through the canonical isomorphisms $V_k \cong (V_k^*)^*$ and $V_{2\ell} \cong (V_{2\ell}^*)^*$, we can view R as the space of regular functions on $(SV_k \otimes (\wedge V_{2\ell})_{\bar{0}})^*$. By \mathcal{CG} we denote the space of formal linear combinations of elements of \mathcal{G} with complex coefficients. Analogous to [9] and [24], we can define a map $p : \mathcal{CG} \rightarrow R$ such that for each $(k, 2\ell)$ -color edge coloring model $h \in (SV_k \otimes \wedge V_{2\ell})^*$ we have $p(G)(h) = p_h(G)$ for each graph G . (We refer to [26] for the explicit construction.)

To characterize mixed partition functions, following [9, 24], the idea is to consider the ideal generated by $p(G) - f(G)$ for all graphs G in R , where f is a graph parameter. Provided certain conditions are satisfied, Hilbert's Nullstellensatz can be used to find a common 'zero' for this ideal. Such a common zero h is exactly an edge-coloring model h such that $p_h(G) = p(G)(h) = f(G)$ for all graphs G .

Now in case $\ell = 0$, (resp. $k = 0$), the algebraic characterization of these respective partition function reads that the graph parameter f should map the kernel of the map p to zero (i.e. we first extend f linearly to a map $f : \mathcal{CG} \rightarrow \mathbb{C}$; this map should satisfy $\ker p \subseteq \ker f$). The image of this map p turns out to be the space of polynomials in R that are invariant under a natural action of the orthogonal group [9] (resp. the symplectic group [24]). The kernel of p can be described using the second fundamental theorem of invariant theory for the orthogonal (resp. symplectic) group. Using reductivity of the orthogonal and symplectic group, it can be shown that this yields a valid characterization of these respective partition functions.

When both k and 2ℓ are positive, we conjecture that the multiplicative graph parameters f that satisfy $\ker p \subseteq \ker f$ are exactly the mixed partition functions.

The kernel of p can still be described [26] and is related to the second fundamental theorem of invariant theory of the orthosymplectic supergroup cf. [15]. So to prove this

conjecture along the same lines of [9, 24] we would need a good understanding of the image of the map p .

There is a natural action of $O_k \times Sp_{2\ell}$ on R and the image of the map p consists of $O_k \times Sp_{2\ell}$ -invariants. However it is **not** true that the image of p is equal to the space of elements of R that are invariant under the group $O_k \times Sp_{2\ell}$. To say more about the image of p , we need some definitions from supersymmetry. We refer to [4, 6] for background on supersymmetry and for notation that we use here.

The space $SV_k \otimes \wedge V_{2\ell}$ has the structure of a super vector space. Its even part is $SV_k \otimes (\wedge V_{2\ell})_{\bar{0}}$ and its odd part is $SV_k \otimes (\wedge V_{2\ell})_{\bar{1}}$. The tensor algebra $T = T(SV_k \otimes \wedge V_{2\ell})$ then naturally carries the structure of a super algebra. The *super symmetric algebra*, S , is the quotient of T by the ideal generated by $x \otimes y - (-1)^{|x||y|} y \otimes x$ with $x, y \in SV_k \otimes \wedge V_{2\ell}$ homogeneous elements. (Here $|x| = 0$ if x is even and $|x| = 1$ if x is odd for any homogeneous element x of a super vector space.) Then R is a subalgebra of S and we have a natural projection $\Pi : S \rightarrow R$.

The orthosymplectic Lie superalgebra $\mathfrak{osp}(V_{k,2\ell})$ is the Lie superalgebra preserving the form (19), i.e., for each $X \in \mathfrak{osp}(V_{k,2\ell})$, we have that $[Xv, w] = (-1)^{|X||v|} [v, Xw]$, where we assume all elements involved to be homogenous and we view X as an element of $\text{End}(V_{k,2\ell})$. (Both $V_{k,2\ell}$ and $\text{End}(V_{k,2\ell})$ naturally carry the structure of a super vector space.) It turns out that the Lie super algebra $\mathfrak{osp}(V_{k,2\ell})$ and the group $O_k \times Sp_{2\ell}$ have a natural action on S and that

$$\text{im}(p) = \Pi(S^{O_k \times Sp_{2\ell}, \mathfrak{osp}(V_{k,2\ell})}),$$

the projection onto R of the space of elements of S that are invariant under the action of $O_k \times Sp_{2\ell}$ and the action of $\mathfrak{osp}(V_{k,2\ell})$. (This is proved in [26] using results of Berele and Regev [3] and recent results of Leher and Zhang [16].)

Since the Lie super algebra $\mathfrak{osp}(V_{k,2\ell})$ is not reductive, it is not clear how to continue the proof outline from [9, 24]. We expect that with a thorough understanding of invariant theory in this super symmetric setting one may finish the proof outline sketched here.

6 Concluding remarks

In this paper we have introduced mixed partition functions and given several examples of mixed partition functions, shown that they have finite edge-rank connectivity and briefly discussed connections with invariant theory. In [26] we will focus on the algebraic and invariant-theoretic aspects of mixed partition functions. By exploiting recent developments in the invariant theory of the orthosymplectic supergroup [15, 16, 17], we aim to prove that any multiplicative graph parameter with finite edge-connection rank is a mixed partition function.

It would be interesting to find more examples of mixed partition functions. Given the fact that supersymmetry originated in physics, it would be interesting to explore if there are natural statistical physics models that give rise to interesting combinatorial parameters, similar to how the Potts model partition function is related to the Tutte polynomial for example.

Acknowledgements

The research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013) / ERC grant agreement n° 339109.

We thank Lex Schrijver for useful comments on an earlier version of this paper. We moreover thank the anonymous referees for their constructive feedback.

References

- [1] B. Bollobás, Evaluations of the circuit partition polynomial, *Journal of Combinatorial Theory, Series B* **85.2** (2002) 261-268.
- [2] A.E. Brouwer, W.H. Haemers, *Spectra of Graphs*. Springer Science & Business Media, 2011.
- [3] A. Berele, A. Regev, Hook Young diagrams with applications to combinatorics and to representations of Lie superalgebras, *Advances in Mathematics* **64** (1987) 118–175.
- [4] C. Carmeli, Caston, and Rita Fiorese, *Mathematical foundations of supersymmetry*, Vol. 15, European Mathematical Society, 2011.
- [5] J. Cai, P. Lu, M. Xia, Computational complexity of Holant problems, *SIAM Journal on Computing* **40** (2011) 1101–1132.
- [6] S.J. Cheng, S. W. Wang, *Dualities and representations of Lie superalgebras*, American Mathematical Soc., 2012.
- [7] S. Chmutov, S. Duzhin, J. Mostovoy, *Introduction to Vassiliev Knot Invariants*, Cambridge University Press, 2012.
- [8] R. Diestel, *Graph theory*, Graduate Texts in Mathematics, Vol. 173, Springer Verlag, Heidelberg, 2000.
- [9] J. Draisma, D. Gijswijt, L. Lovász, G. Regts, A. Schrijver, Characterizing partition functions of the vertex model, *Journal of Algebra* **350** (2012) 197–206.
- [10] J.A. Ellis-Monaghan, Identities for circuit partition polynomials, with applications to the Tutte polynomial, *Advances in Applied Mathematics* **32** (2004) 188–197.
- [11] M. Freedman, L. Lovász, A. Schrijver, Reflection positivity, rank connectivity, and homomorphisms of graphs, *Journal of the American Mathematical Society* **20** (2007) 37–51.
- [12] P. de la Harpe, V.F.R. Jones, Graph invariants related to statistical mechanical models: examples and problems, *Journal of Combinatorial Theory, Series B* **57** (1993) 207–227.
- [13] M. Kontsevich, Vassiliev's knot invariants, *Adv. in Sov. Math* **16.2** (1993) 137–150.
- [14] T. Kotek, J. A. Makowsky, Connection matrices and the definability of graph parameters, in 26th International Workshop/21st Annual Conference of the EACSL, *Leibniz International Proceedings in Informatics (LIPIcs)*, pp. 411–425, Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik.

- [15] G.I. Lehrer, R.B. Zhang, The second fundamental theorem of invariant theory for the orthosymplectic supergroup, *Nagoya Mathematical Journal* (2014), 1–25.
- [16] G.I. Lehrer, R. B. Zhang, The first fundamental theorem of invariant theory for the orthosymplectic supergroup, *Communications in Mathematical Physics* **349** (2017), 661–702.
- [17] G.I. Lehrer, R. B. Zhang, Invariants of the orthosymplectic Lie superalgebra and super Pfaffians, *Mathematische Zeitschrift* **286** (2017), 893–917.
- [18] L. Lovász, *Large Networks and Graph Limits*, American Mathematical Society, Providence, Rhode Island, 2012.
- [19] I. L. Markov, Y. Shi, Simulating quantum computation by contracting tensor networks, *SIAM Journal on Computing* **38** (2008) 963–981.
- [20] P. Martin, *Enumérations eulériennes dans les multigraphes et invariants de Tutte-Grothendieck*, Diss. Institut National Polytechnique de Grenoble-INPG; Université Joseph-Fourier-Grenoble I, 1977.
- [21] G. Regts, *Graph Parameters and Invariants of the Orthogonal Group*, PhD thesis, University of Amsterdam, 2013.
- [22] G. Regts, A. Schrijver, B. Sevenster, On partition functions for 3-graphs, *Journal of Combinatorial Theory, Series B* **121** (2016) 421–431.
- [23] G. Regts, A. Schrijver, B. Sevenster, On the existence of real R-matrices for virtual link invariants, *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg* **87** (2017) 435–443.
- [24] G. Regts, B. Sevenster, Graph parameters from symplectic group invariants, *Journal of Combinatorial Theory, Series B* **122** (2017), 844–868.
- [25] G. Regts, B. Sevenster, Partition functions from orthogonal and symplectic group invariants, *Electronic Notes in Discrete Mathematics* **61** (2017), 1011–1017.
- [26] G. Regts, B. Sevenster, Mixed partition functions and invariants of the orthosymplectic group, manuscript.
- [27] H. Sachs, Über Teiler, Faktoren und charakteristische Polynome von Graphen. *Wiss. Z. Techn. Hochsch. Ilmenau* **13** (1967) 405–412.
- [28] A. Schrijver, Characterizing partition functions of the edge-coloring model by rank growth, *Journal of Combinatorial Theory, Series A* **136** (2015) 164–173.
- [29] A. Schrijver, Connection matrices and Lie algebra weight systems for multiloop chord diagrams, *Journal of Algebraic Combinatorics* **42** (2015) 896–905.
- [30] A. Schrijver, On traces of tensor representations of diagrams, *Linear Algebra and its Applications* **278** (2015) 28–41.
- [31] B.L. Sevenster, *New characterizations of partition functions using connection matrices*, PhD thesis, University of Amsterdam, 2018.

- [32] B. Szegedy, Edge-coloring models and reflection positivity, *Journal of the American Mathematical Society* **20** (2007) 969–988.