



## UvA-DARE (Digital Academic Repository)

### Modular Forms of Degree 2 and Curves of Genus 2 in Characteristic 2

Cléry, F.; van der Geer, G.

**DOI**

[10.1093/imrn/rnaa239](https://doi.org/10.1093/imrn/rnaa239)

**Publication date**

2022

**Document Version**

Final published version

**Published in**

International Mathematics Research Notices

**License**

CC BY

[Link to publication](#)

**Citation for published version (APA):**

Cléry, F., & van der Geer, G. (2022). Modular Forms of Degree 2 and Curves of Genus 2 in Characteristic 2. *International Mathematics Research Notices*, Article rnaa239. <https://doi.org/10.1093/imrn/rnaa239>

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

# Modular Forms of Degree 2 and Curves of Genus 2 in Characteristic 2

**Fabien Cléry<sup>1</sup> and Gerard van der Geer<sup>2,\*</sup>**

<sup>1</sup>Department of Mathematics, Loughborough University, England, and Institute of Computational and Experimental Research in Mathematics, 121 South Main Street, Providence, RI 02903, USA, <sup>2</sup>Korteweg-de Vries Instituut, Universiteit van Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands, and Université du Luxembourg, Unité de Recherche en Mathématiques, L-4364 Esch-sur-Alzette, Luxembourg

*\*Correspondence to be sent to: e-mail: [g.b.m.vandergeer@uva.nl](mailto:g.b.m.vandergeer@uva.nl)*

We describe the ring of modular forms of degree 2 in characteristic 2 using its relation with curves of genus 2.

## 1 Introduction

In the 1960s, Igusa determined the ring of Siegel modular forms of degree 2 using the close connection between the moduli of principally polarized abelian surfaces and the moduli of curves of genus 2, see [9]. Since curves of genus 2 in characteristic different from 2 can be described by binary sextics, he could use the invariant theory of binary sextics and establish the link with modular forms using Thomae's formulas that express theta constants in terms of cross ratios of zeros of binary sextics. Recently, in a joint work with Carel Faber, we exhibited in [2] a more direct version of this without the detour of Thomae's formulas and used it for describing vector-valued Siegel modular forms of degree 2.

Received March 10, 2020; Revised July 14, 2020; Accepted August 19, 2020

© The Author(s) 2020. Published by Oxford University Press.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

In positive characteristic, modular forms of degree 2 are described as sections of powers of the determinant bundle  $L$  of the Hodge bundle  $\mathbb{E}$  on the moduli space  $\mathcal{A}_2 \otimes \mathbb{F}_p$  of principally polarized abelian surfaces in characteristic  $p$ . It turns out that for  $p \geq 5$  the situation is very close to the case of characteristic 0. For  $p \geq 5$ , the graded ring

$$\mathcal{R}_2(\mathbb{F}_p) = \bigoplus_k H^0(\mathcal{A}_2 \otimes \mathbb{F}_p, L^k)$$

has generators of weight 4, 6, 10, 12, and 35 and there is a relation of degree 70 expressing the square of the odd weight generator in terms of the even weight generators, just as in characteristic 0. In fact, Igusa determined the ring  $\mathcal{R}_2(\mathbb{Z})$  of integral modular forms and the reduction map  $\mathcal{R}_2(\mathbb{Z}) \rightarrow \mathcal{R}_2(\mathbb{F}_p)$  is surjective for  $p \geq 5$ . We refer to [1, 10, 11, 15, 16].

The situation in characteristics 2 and 3 is different. The ring  $\mathcal{R}_2(\mathbb{F}_3)$  was determined in [7] using the relation with binary sextics.

In characteristic 2, curves of genus 2 can no longer be described by binary sextics. Nevertheless, as Igusa showed in [9], there is still a close relationship with the invariant theory of binary sextics. In this paper we use invariants of curves of genus 2 in characteristic 2 to determine the ring of modular forms of degree 2. The result is as follows.

**Theorem 1.1.** The ring  $\mathcal{R}_2(\mathbb{F}_2)$  is generated by modular forms of weights 1, 10, 12, 13, and 48 with one relation of weight 52:

$$\mathcal{R}_2(\mathbb{F}_2) = \mathbb{F}_2[\psi_1, \chi_{10}, \psi_{12}, \chi_{13}, \chi_{48}]/(R)$$

with  $R = \chi_{13}^4 + \psi_1^3 \chi_{10} \chi_{13}^3 + \psi_1^4 \chi_{48} + \chi_{10}^4 \psi_{12}$ . The ideal of cusp forms is generated by  $\chi_{10}$ ,  $\chi_{13}$  and  $\chi_{48}$ .

The modular form  $\psi_1$  is the Hasse invariant vanishing on the locus of non-ordinary abelian surfaces; the form  $\chi_{10}$  vanishes doubly on the locus of abelian surfaces that are products of elliptic curves. All the forms  $\psi_1, \chi_{10}, \psi_{12}, \chi_{13}$  and  $\chi_{48}$  are constructed using invariant theory, see Section 4.

We point out that we find non-zero (regular) vector-valued modular forms of weights not allowed in characteristic 0, like  $(3, -1)$  or  $(2, 0)$ .

We thank the referees for helpful remarks.

## 2 Modular Forms

We denote by  $\mathcal{A}_g$  the moduli stack of principally polarized abelian varieties of dimension  $g$  and by  $\mathbb{E}_g$  the Hodge bundle on  $\mathcal{A}_g$ . It extends to a Faltings–Chai-type toroidal compactification  $\tilde{\mathcal{A}}_g$ . We write  $L = \det(\mathbb{E}_g)$ . For  $g > 1$  sections of  $L^k$  over  $\mathcal{A}_g$  extend automatically to  $\tilde{\mathcal{A}}_g$ , a property known as the Koecher principle, see [5, Prop. 1.5, p. 140]. We write  $M_k(\Gamma_g) = H^0(\tilde{\mathcal{A}}_g \otimes \mathbb{F}_2, L^k)$  for the space of scalar-valued modular forms of weight  $k$ . Moreover, we set

$$\mathcal{R}_g(\mathbb{F}_2) = \bigoplus_k M_k(\Gamma_g),$$

the ring of scalar-valued modular forms of degree  $g$  in characteristic 2. It is a finitely generated  $\mathbb{F}_2$ -algebra. It is well-known that  $H^0(\tilde{\mathcal{A}}_g, L^k) = (0)$  for  $k < 0$ . For  $g = 1$  we know by Deligne [3] that  $\mathcal{R}_1(\mathbb{F}_2) = \mathbb{F}_2[c_1, \Delta]$  with  $c_1$  of weight 1 and  $\Delta$  a cusp form of weight 12.

We are interested in the case  $g = 2$ . In the following we shall simply write  $\mathbb{E}$  for the Hodge bundle  $\mathbb{E}_2$ .

In the Chow group with rational coefficients of codimension 1 cycles of  $\tilde{\mathcal{A}}_2$  the class of the locus  $\mathcal{A}_{1,1}$  of products of elliptic curves satisfies the relation

$$2[\overline{\mathcal{A}}_{1,1}] + [D] = 10\lambda_1, \quad (1)$$

where  $\lambda_1$  is the 1st Chern class of  $\mathbb{E}$  and  $D$  the divisor added to compactify  $\mathcal{A}_2$ , see [14, (8.4)]. Therefore, there exists a non-zero modular form  $\chi_{10}$  of weight 10 over  $\mathbb{F}_2$  vanishing doubly on the divisor  $\overline{\mathcal{A}}_{1,1} \otimes \mathbb{F}_2$ . As the cycle relation shows  $\chi_{10}$  is a cusp form. The form is determined up to a non-zero multiplicative constant and we shall normalize  $\chi_{10}$  later.

Since  $\mathcal{A}_{1,1}$  is the image of a degree 2 morphism  $\mathcal{A}_1 \times \mathcal{A}_1 \rightarrow \mathcal{A}_{1,1}$  by restriction to the locus  $\mathcal{A}_{1,1} \otimes \mathbb{F}_2$  we find an exact sequence

$$0 \rightarrow M_{k-10}(\Gamma_2) \rightarrow M_k(\Gamma_2) \rightarrow \text{Sym}^2(M_k(\Gamma_1)). \quad (2)$$

Another divisor that we will use is the locus  $V_1$  of principally polarized abelian surfaces of 2-rank  $\leq 1$ . The cycle class of this locus is known; it is the vanishing locus of the map  $L \rightarrow L^{\otimes 2}$  induced by Verschiebung on the universal abelian surface and we thus find (cf. [6])

$$[V_1] = \lambda_1.$$

Hence, there exists a modular form  $\psi_1$  of weight 1, determined up to a non-zero multiplicative constant, with divisor  $V_1$ . It is called the Hasse invariant. This is not a cusp form since it restricts to the Hasse invariant  $c_1$  on the boundary component  $\mathcal{A}_1^* \otimes \mathbb{F}_2$  in the Satake compactification  $\mathcal{A}_2^* \otimes \mathbb{F}_2$ . We shall normalize  $\psi_1$  later.

The existence of  $\psi_1$  implies  $\dim M_k(\Gamma_2) \leq \dim M_{k+1}(\Gamma_2)$ . The exact sequence (2) and the structure of  $\mathcal{R}_1(\mathbb{F}_2)$  imply that

$$\dim M_k(\Gamma_2) = 1 \quad \text{for } k = 1, \dots, 9 \quad \text{and } \dim M_{10}(\Gamma_2) = 2. \tag{3}$$

In order to estimate  $\dim M_k(\Gamma_2)$  for larger  $k$  we can apply semi-continuity (cf. [8, Thm. 12.8]) to  $L$  defined over  $\tilde{\mathcal{A}}_2/\mathbb{Z}$  and deduce  $\dim H^0(\tilde{\mathcal{A}}_2 \otimes \mathbb{C}, L^k) \leq \dim H^0(\tilde{\mathcal{A}}_2 \otimes \mathbb{F}_2, L^k)$ . Using this and the exact sequence (2) we can deduce

$$\dim M_{11}(\Gamma_2) = 2, \quad \text{and } 3 \leq \dim M_k(\Gamma_2) \leq 4 \text{ for } 12 \leq k \leq 19, \tag{4}$$

but we will deduce this in another way later, see (7).

Similarly, we can consider vector-valued modular forms. For each irreducible representation  $\rho$  of  $\text{GL}(2, \mathbb{Q})$  there is a corresponding bundle  $\mathbb{E}_\rho$ . Since we are dealing with  $g = 2$  the  $\rho$  in question correspond to pairs  $(j, k)$  of integers with  $j \geq 0$  and  $\mathbb{E}_\rho = \text{Sym}^j(\mathbb{E}) \otimes \det(\mathbb{E})^{\otimes k}$ . We write  $M_{j,k}(\Gamma_2)$  for  $H^0(\mathcal{A}_2 \otimes \mathbb{F}_2, \text{Sym}^j(\mathbb{E}) \otimes \det(\mathbb{E})^{\otimes k})$  and  $S_{j,k}(\Gamma_2)$  for the space of cusp forms. Again, these sections extend over  $\tilde{\mathcal{A}}_2 \otimes \mathbb{F}_2$  by the Koecher principle. But note that  $\rho \otimes \mathbb{F}_2$  might be reducible.

**Lemma 2.1.** We have  $\dim S_{6,8}(\Gamma_2) \geq 1$  and every element of  $S_{6,8}(\Gamma_2)$  vanishes on  $\mathcal{A}_{1,1} \otimes \mathbb{F}_2$ . If  $\dim S_{6,8}(\Gamma_2) \geq 2$  then  $M_{6,-2}(\Gamma_2) \neq 0$ .

**Proof.** We know that over  $\mathbb{C}$  we have  $\dim S_{6,8}(\Gamma_2) = 1$ . Hence, by semi-continuity of dimensions there exists a cusp form of weight  $(6, 8)$  in characteristic 2. Note that the Hodge bundle  $\mathbb{E}$  pulled back via

$$\mathcal{A}_1 \times \mathcal{A}_1 \rightarrow \mathcal{A}_{1,1} \hookrightarrow \mathcal{A}_2$$

can be written as  $p_1^*(\mathbb{E}_1) \oplus p_2^*(\mathbb{E}_1)$  with  $\mathbb{E}_1$  the Hodge bundle of  $\mathcal{A}_1$  and  $p_1, p_2$  the two projections. The pullback of  $\chi_{6,8}$  to  $(\mathcal{A}_1 \times \mathcal{A}_1) \otimes \mathbb{F}_2$  must vanish because it lands in

$$\bigoplus_{j=0}^6 S_{14-j}(\Gamma_1) \otimes S_{8+j}(\Gamma_1)$$

and this is (0). If we develop a form  $\varphi \in S_{6,8}(\Gamma_2)$  vanishing on  $\mathcal{A}_{1,1} \otimes \mathbb{F}_2$  in the normal direction of the divisor  $\mathcal{A}_{1,1} \otimes \mathbb{F}_2$  we see that the 1st (“non-constant”) term in its Taylor expansion lands in

$$\bigoplus_{j=0}^6 S_{15-j}(\Gamma_1) \otimes S_{9+j}(\Gamma_1)$$

and the only non-zero factor here is  $S_{12}(\Gamma_1) \otimes S_{12}(\Gamma_1)$  and the image is symmetric under the interchange of factors. But  $\dim S_{12}(\Gamma_1) = 1$ . Therefore, if  $\varphi_1, \varphi_2$  are two linearly independent cusp forms of weight (6, 8) there exists a non-trivial linear combination that vanishes with order  $\geq 2$  along  $\mathcal{A}_{1,1} \otimes \mathbb{F}_2$ . This yields a modular form divisible by  $\chi_{10}$ , hence a regular section of  $\text{Sym}^6(\mathbb{E}) \otimes \det(\mathbb{E})^{-2}$ , not identically zero on  $\mathcal{A}_{1,1} \otimes \mathbb{F}_2$ . ■

We thus have at least one non-zero modular form  $f \in S_{6,8}(\Gamma_2)$  that vanishes with multiplicity 1 on  $\mathcal{A}_{1,1} \otimes \mathbb{F}_2$ . Thus, we also find a rational modular form  $f/\chi_{10}$  with possible poles along  $\mathcal{A}_{1,1} \otimes \mathbb{F}_2$ .

We can analyze the order of the poles of the coordinates of  $f/\chi_{10}$  near a generic point of  $\mathcal{A}_{1,1}$ . We consider  $f$  near a generic point of  $\mathcal{A}_{1,1} \otimes \mathbb{F}_2$  and write its Taylor expansion as

$$f = (\gamma_0, \dots, \gamma_6) \quad \text{with } \gamma_i = \sum_{r \geq 0} \gamma_{i,r} t^r,$$

where  $t$  is a local normal coordinate. If for fixed  $i$  we have  $\gamma_{i,r} = 0$  for  $r < r_0$  then

$$\gamma_{i,r_0} \in QS_{14+r_0-i}(\Gamma_1) \otimes QS_{8+r_0+i}(\Gamma_1),$$

where  $QS$  denotes quasi-modular cusp forms, see [17, Sec. 5] for the notion of quasi-modular form. From the dimensions of  $S_k(\Gamma_1)$  we see as in the proof of Lemma 2.1 that

$\text{ord}_{\mathcal{A}_{1,1}}(\gamma_3) \geq -1$  and we can even get the estimate

$$\text{ord}_{\mathcal{A}_{1,1}}(\gamma_0, \dots, \gamma_6) \geq (2, 1, 0, -1, 0, 1, 2),$$

but we will not use this.

### 3 Curves of Genus 2 in Characteristic 2

Let  $C$  be a smooth projective curve of genus 2 over a perfect field  $k$  of characteristic 2. Let  $K$  be the canonical divisor (class) of  $C$ . Then by Riemann–Roch  $\dim H^0(C, O(nK)) = 2n - 1$  for  $n \geq 2$ . If  $\xi_0, \xi_1$  is a basis of  $H^0(C, O(K))$ , then there is an element  $\eta \in H^0(C, O(3K))$  such that  $\xi_0^3, \xi_0^2\xi_1, \xi_0\xi_1^2, \xi_1^3, \eta$  form a basis of  $H^0(C, O(3K))$ . Then

$$\xi_0^6, \xi_0^5\xi_1, \dots, \xi_1^6, \xi_0^3\eta, \xi_0^2\xi_1\eta, \xi_0\xi_1^2\eta, \xi_1^3\eta$$

are linearly independent in  $H^0(C, O(6K))$  and since  $\eta^2 \in H^0(C, O(6K))$  there must be a linear relation

$$\eta^2 + c_3\eta + c_6 = 0$$

with  $c_3$  (resp.  $c_6$ ) homogeneous of degree 3 (resp. 6) in  $\xi_0, \xi_1$ . Setting  $x = \xi_1/\xi_0$  and  $y = \eta/\xi_0^3$  we can write the equation as

$$y^2 + ay = b \tag{5}$$

with  $a \in k[x]$  non-zero of degree  $\leq 3$  and  $b \in k[x]$  of degree  $\leq 6$ . The hyperelliptic involution of the curve  $C$  is given by  $y \mapsto y+a$ . The fixed points are given by the equation  $a = 0$  (or  $c_3 = 0$  in projective coordinates); hence, there are 3, 2 or 1 ramification points. This corresponds to the 2-rank of  $\text{Jac}(C)$  being equal to 2, 1 or 0.

We point out that a curve of genus 2 defined by an equation (5) comes with a basis  $dx/a, x dx/a$  of  $H^0(C, O(K))$ .

The choices we made for arriving at equation (5) are a basis of  $H^0(C, O(K))$  and an element of  $\eta$  in  $H^0(C, O(3K))$ . Clearly, another choice of  $\eta$  is of the form  $\epsilon\eta + \theta$  with  $\epsilon$  a unit in  $k$  and  $\theta$  homogeneous of degree 3 in  $\xi_0, \xi_1$ , or  $y \rightarrow uy + v$  for  $u \in k^*$  and  $v \in k[x]$  of degree  $\leq 3$ .

The induced action on the pair  $(a, b)$  is by

$$(a, b) \mapsto (a/u, (b + v^2 + av)/u^2) \quad \text{with } u \in k^*, v \in k[x] \text{ of degree } \leq 3.$$

Another choice of basis of  $H^0(C, \mathcal{O}(K))$  is given by an element of  $\mathrm{GL}(V)$  with  $V = \langle \xi_0, \xi_1 \rangle$ . The action on  $x, y$  is by

$$x \mapsto (\alpha x + \beta)/(\gamma x + \delta), \quad y \mapsto y/(\gamma x + \delta)^3.$$

Let  $Y$  be the algebraic stack of triples  $(\pi, \alpha, \beta)$  with  $\pi : C \rightarrow S$  a curve of genus 2,  $\alpha : \mathcal{O}_S^{\oplus 2} \xrightarrow{\sim} \pi_* \omega_\pi$  and  $\beta \in \pi_*(\omega^{\otimes 3})$  nowhere in the image of  $\mathrm{Sym}^3(\pi_*(\omega_\pi)) \rightarrow \pi_*(\omega_\pi^{\otimes 3})$ . We may view it as the stack of curves of genus 2 with a framed Hodge bundle and a section of  $\omega_\pi^{\otimes 3}$  satisfying the additional condition that it yields an equation as (5). There is an obvious action of  $\mathrm{GL}(2)$  and an action of  $\mathrm{Sym}^3(\mathcal{O}_S^{\oplus 2})$ .

We thus consider the stack

$$[Y/\mathrm{GL}(2) \times \mathrm{Sym}^3(\mathcal{O}_S^{\oplus 2})]$$

and we can identify it with the moduli space  $\mathcal{M}_2 \otimes \mathbb{F}_2$  of curves of genus 2 in characteristic 2.

We now describe a concrete form of this stack. Let  $V$  be a 2-dimensional  $k$ -vector space generated by  $x_1, x_2$ . Consider the subspace of  $\mathrm{Sym}^3(V) \times \mathrm{Sym}^6(V)$  of pairs  $(a, b)$  satisfying the condition that the pair defines a non-singular curve; in the affine version this amounts to  $a$  and  $(a')^2 b + (b')^2$  (with  $a'$  and  $b'$  the derivative) having no root in common.

We write  $V_{j,l}$  for  $\mathrm{Sym}^j(V) \otimes \det(V)^{\otimes l}$ . We let the semi-direct product  $\mathrm{GL}(V) \ltimes V_{3,-1}$  act on  $V_{3,-1} \times V_{6,-2}$  via twisting the two actions of  $\mathrm{GL}(V)$  and  $\mathrm{Sym}^3(V)$ . Without twisting an element  $M \in \mathrm{GL}(V)$  acts by

$$(a, b) \mapsto (a(\alpha x_1 + \beta x_2, \gamma x_1 + \delta x_2), b(\alpha x_1 + \beta x_2, \gamma x_1 + \delta x_2))$$

and  $v \in \mathrm{Sym}^3(V)$  by

$$(a, b) \mapsto (a, b + v^2 + va).$$

After twisting  $c \cdot \mathrm{Id}_V$  acts via  $c$  on  $V_{3,-1}$  and by  $c^2$  on  $V_{6,-2}$  and this is compatible with the equation  $y^2 + ay = b$  if we let  $c \cdot \mathrm{Id}_V$  act on  $y$  by  $y \mapsto cy$ . It is also compatible with  $b \mapsto b + v^2 + av$  for  $v \in V_{3,-1}$ .



We thus consider

$$\mathcal{X}^0 \subset \mathcal{X} = V_{3,-1} \times V_{6,-2},$$

where  $\mathcal{X}^0$  is the open substack given by the condition that  $y^2 + ay = b$  defines a smooth projective curve of genus 2.

The moduli space  $\mathcal{M}_2 \otimes \mathbb{F}_2$  can be identified with the stack quotient

$$[\mathcal{X}^0/\mathrm{GL}(V) \times V_{3,-1}].$$

Note that by our choice of twisting the stabilizer of a pair  $(a, b)$  contains  $(\mathrm{id}_V, a)$  as it should, since the generic curve has an automorphism group of order 2. The pullback of the Hodge bundle  $\mathbb{E}$  on  $\mathcal{M}_2 \otimes \mathbb{F}_2$  is the equivariant bundle  $V$ .

#### 4 Invariants and Modular Forms

We write elements  $a \in \mathrm{Sym}^3(V)$  and  $b \in \mathrm{Sym}^6(V)$  as

$$a = \sum_{i=0}^3 a_i x_1^{3-i} x_2^i, \quad b = \sum_{i=0}^6 b_i x_1^{6-i} x_2^i.$$

Following standard usage, by an invariant for the action of  $\mathrm{GL}(V) \times \mathrm{Sym}^3(V)$  we mean a polynomial in the coordinates  $a_0, \dots, a_3$  and  $b_0, \dots, b_6$  of the pair  $(a, b)$  in  $\mathrm{Sym}^3(V) \times \mathrm{Sym}^6(V)$  that is invariant under  $\mathrm{SL}(V) \times \mathrm{Sym}^3(V)$ . We define  $\mathcal{K}$  as the ring of  $\mathrm{SL}(V) \times \mathrm{Sym}^3(V)$ -invariants under the action on  $\mathcal{X}$ .

The simplest example of such an invariant is the expression

$$K_1 = a_0 a_3 + a_1 a_2,$$

the square root of the discriminant of  $a$ .

Since the pullback to  $\mathcal{X}^0$  of the Hodge bundle  $\mathbb{E}$  under the Torelli map  $\mathcal{M}_2 \rightarrow \mathcal{A}_2$  is the equivariant bundle  $V$ , pulling back of scalar-valued modular forms gives invariants. This provides us with a homomorphism

$$\mu : \mathcal{R}_2(\mathbb{F}_2) \rightarrow \mathcal{K}.$$

Since the image of the Torelli map  $\mathcal{M}_2 \hookrightarrow \mathcal{A}_2$  is the open set that is the complement of  $\mathcal{A}_{1,1}$ , the zero locus of  $\chi_{10}$ , we see that an invariant defines a rational modular form that

becomes regular after multiplication by a power of  $\chi_{10}$ . Therefore, we can extend the map  $\mu$  to

$$\mathcal{R}_2(\mathbb{F}_2) \xrightarrow{\mu} \mathcal{K} \xrightarrow{\nu} \mathcal{R}_2(\mathbb{F}_2)_{\chi_{10}}$$

such that  $\nu \cdot \mu = \text{id}$ ; here, the index  $\chi_{10}$  indicates localization at the multiplicative system generated by  $\chi_{10}$ .

Since we know the existence of a modular form  $\psi_1 \in M_1(\Gamma_2)$  we see that  $\mu(\psi_1)$  must equal  $K_1$  as  $K_1$  is the only invariant of weighted degree 3. (Here the weight of  $a_i$  (resp.  $b_j$ ) is  $i$  (resp.  $j$ .) This implies in particular that  $\nu(K_1)$  is a regular modular form of weight 1.

Similarly for covariants. By a covariant under the action of  $\text{GL}(V) \times \text{Sym}^3(V)$ , we mean a polynomial in the coefficients  $a_i$  and  $b_j$  and  $x_1, x_2$ , which is an invariant for the action of  $\text{SL}(V) \times \text{Sym}^3(V)$ .

The simplest example of a covariant is given by the polynomial  $a$ . It defines an a priori rational modular form of weight  $(3, -1)$ .

**Lemma 4.1.** The form  $\nu(a)$  is a regular modular form of weight  $(3, -1)$ .

**Proof.** The pullback of the  $\text{Sym}^3(\mathbb{E})$  to  $(\mathcal{A}_1 \times \mathcal{A}_1) \otimes \mathbb{F}_2$  decomposes as

$$\bigoplus_{j=0}^3 p_1^*(\mathbb{E}_1)^{3-j} \otimes p_2^*(\mathbb{E}_1)^j$$

and the coefficients of  $a$  correspond to the factors. In view of the symmetry we have  $\text{ord}_{\mathcal{A}_{1,1}}(a_0) = \text{ord}_{\mathcal{A}_{1,1}}(a_3)$  and  $\text{ord}_{\mathcal{A}_{1,1}}(a_1) = \text{ord}_{\mathcal{A}_{1,1}}(a_2)$ . As we saw above the form  $\nu(K_1) = \nu(a_0 a_3 + a_1 a_2)$  is regular. By coordinate changes we may assume that  $a_0 = 0$  or  $a_1 = 0$ , so it easily follows that  $\text{ord}_{\mathcal{A}_{1,1}}(a_i) \geq 0$  and the form is regular. Moreover, since its pullback lands in  $\bigoplus_{j=0}^3 M_{2-j}(\Gamma_1) \otimes M_{j-1}(\Gamma_1)$  and  $M_2(\Gamma_1) \otimes M_{-1}(\Gamma_1) = (0)$  it follows that  $\text{ord}_{\mathcal{A}_{1,1}}(a_0) \geq 1$  and  $\text{ord}_{\mathcal{A}_{1,1}}(a_3) \geq 1$ . ■

**Corollary 4.2.** We have  $\text{ord}_{\mathcal{A}_{1,1}}(a_i) = 0$  for  $i = 1, 2$  and  $\text{ord}_{\mathcal{A}_{1,1}}(a_i) \geq 1$  for  $i = 0, 3$ .

**Proof.** We note that  $\psi_1$  does not vanish on  $\mathcal{A}_{1,1} \otimes \mathbb{F}_2$ . ■

**Remark 4.3.** The restriction of the modular form  $\nu(a)$  to  $\mathcal{A}_{1,1} \otimes \mathbb{F}_2$  is the form  $c_1 \otimes 1 + 1 \otimes c_1 \in M_1(\Gamma_1) \otimes M_0(\Gamma_1) \oplus M_0(\Gamma_1) \otimes M_1(\Gamma_1)$ .

Another example is the expression

$$(a_0 a_2 + a_1^2) x_1^2 + (a_0 a_3 + a_1 a_2) x_1 x_2 + (a_1 a_3 + a_2^2) x_2^2,$$

which then defines a regular modular form of weight  $(2, 0)$ . In fact, this lives in the 2nd factor of the decomposition  $\text{Sym}^2(V_{3,-1}) = V_{6,-2} \oplus V_{2,0}$ . Note that the map  $V_{2,0} \rightarrow V_{0,0}$  given by  $\alpha x_1^2 + \beta x_1 x_2 + \gamma x_2^2 \mapsto \beta$  defines an invariant in characteristic 2.

### 5 Modular Forms from Invariants

Igusa used in [9] invariants of binary sextics to construct modular forms and to construct a coarse moduli space of curves of genus 2. The  $\mathbb{Z}$ -algebra of even degree invariants of binary sextics describes a coarse moduli space

$$\mathcal{Y} = \text{Proj} \left( \mathbb{Z}[J_2, J_4, J_6, J_8, J_{10}] / (J_4^2 - J_2 J_6 + 4 J_8) \right)$$

for curves of genus 2 over  $\mathbb{Z}$ .

To get characteristic 2 invariants (or covariants) one lifts the curve given by  $y^2 + ay = b$  to the Witt ring, say defined by  $(\tilde{a}, \tilde{b})$ , and takes an invariant (or covariant) of the binary sextic defined by  $\tilde{a}^2 + 4\tilde{b}$ , divides these by an appropriate power of 2 and reduces these modulo 2. This defines invariants (or covariants).

If  $f = \sum_{i=0}^6 c_i x^{6-i}$  is the (universal) binary sextic then the invariant

$$J_2 = 2^{-2}(-120 c_0 c_6 + 20 c_1 c_5 - 8 c_2 c_4 + 3 c_3^2)$$

gives by the substitution

$$c_i = 4 b_i + \sum_{0 \leq r, s \leq 3, r+s=i} a_r a_s \tag{6}$$

the invariant  $K_2 = (a_0 a_3 + a_1 a_2)^2$ , that is,  $K_2 = K_1^2$ . For the formula for  $J_2$  and the formulas of the invariants  $J_{2i}$  ( $i = 2, 3, 5$ ) used below we refer to [13, pp. 139–140] and [12, p. 204].

Similarly, the invariant  $J_4$  given by

$$2^{-7}(2640 c_0^2 c_6^2 - 880 c_0 c_1 c_5 c_6 + \dots)$$

yields via  $I_4 = J_2^2 - 24J_4$  an invariant  $K_4$  in characteristic 2 that turns out to be reducible and divisible by  $K_1$ . We thus set  $K_3 = K_4/K_1$  and get an invariant

$$\begin{aligned} K_3 = & (a_0a_3 + a_1a_2)b_3^2 + (a_0^2a_3^2 + a_0a_2^3 + a_1^3a_3 + a_1^2a_2^2)b_3 + \\ & (a_0a_3 + a_1a_2)a_1^2b_4 + (a_0a_3 + a_1a_2)a_2^2b_2 + \\ & (a_0^2a_1a_3 + a_0^2a_2^2 + a_0a_1^2a_2 + a_1^4)b_5 + \\ & (a_0a_2a_3^2 + a_3^2a_1^2 + a_1a_2^2a_3 + a_2^4)b_1 + \\ & (a_0a_3 + a_1a_2)a_0^2b_6 + (a_0a_3 + a_1a_2)a_3^2b_0. \end{aligned}$$

Similarly,  $J_6$  yields the invariant  $K_3^2$  in characteristic 2. The invariant  $J_8$  yields the invariant

$$\begin{aligned} K_8 = & b_3^8 + (a_0a_3 + a_1a_2)^2b_3^6 + \\ & (a_0a_3 + a_1a_2)(a_0^2a_3^2 + a_0a_2^3 + a_1^3a_3 + a_1^2a_2^2)b_3^5 + \dots, \end{aligned}$$

and similarly from  $J_{10}$  we obtain

$$\begin{aligned} K_{10} = & b_3^6(a_0a_3)^4 + b_3^5(a_0^5a_3^5 + a_0^4a_1a_2a_3^4 + a_0^4a_2^3a_3^3 + a_0^3a_1^3a_3^4) + \\ & b_3^4(a_0^6a_3^6 + a_0^5a_1a_2a_3^5 + a_0^4a_1^2a_2^2a_3^4 + a_0^3a_1^3a_2^3a_3^3 + \\ & a_0^6a_3^4b_6 + a_0^5a_1a_3^4b_5 + a_0^5a_2^2a_3^3b_5 + a_0^4a_1^2a_3^4b_4 + \\ & a_0^4a_1a_2^3a_3^2b_5 + a_0^4a_2^6b_6 + a_0^4a_2^5a_3b_5 + a_0^4a_2^4a_3^2b_4 + \\ & a_0^4a_2^2a_3^4b_2 + a_0^4a_2a_3^5b_1 + a_0^4a_3^6b_0 + a_0^3a_1^2a_3^5b_1 + \\ & a_0^2a_1^4a_3^4b_2 + a_0^2a_1^3a_2a_3^4b_1 + a_0a_1^5a_3^4b_1 + a_1^6a_3^4b_0 + \\ & a_0^4a_2^4b_5^2 + a_1^4a_3^4b_1^2) + \dots. \end{aligned}$$

Finally, the invariant  $I_{15}$  defines an invariant  $K_{15}$ , but  $K_{15}$  can be expressed in terms of the invariants already obtained:

$$K_{15} = K_1^3K_3^4 + K_1^5K_{10} + K_1^4K_3K_8 + K_3^5.$$

**Lemma 5.1.** We have  $\text{ord}_{\mathcal{A}_{1,1}}(b_3) = -1$  and  $\text{ord}_{\mathcal{A}_{1,1}}(b_i) \geq 0$  if  $i \neq 3$ .

**Proof.** In characteristic 0 we have  $\dim S_{6,8}(\Gamma_2) = 1$ . The covariant  $\tilde{f}$ , the universal binary sextic, defines a rational modular form  $\nu(\tilde{f})$  such that  $\chi_{10}\nu(\tilde{f}) \in S_{6,8}(\Gamma_2)$ . A non-zero element  $\chi$  of  $S_{6,8}(\Gamma_2)$  vanishes on  $\mathcal{A}_{1,1}$  and the coordinates of  $\chi/\chi_{10}$  thus have order  $\geq -1$  along  $\mathcal{A}_{1,1}$ . Since the  $a_i$  are regular near  $\mathcal{A}_{1,1} \otimes \mathbb{F}_2$  we see using the formulas (6) that  $\text{ord}_{\mathcal{A}_{1,1}}(b_i) \geq -1$ . By an argument as in Lemma 2.1 if the vanishing order equals  $-1$ , the only non-zero term in the 1st (“non-constant”) term of the Taylor development sits in the middle coordinate and lands in  $S_{12}(\Gamma_1) \otimes S_{12}(\Gamma_1)$  and thus we see that  $\text{ord}_{\mathcal{A}_{1,1}}(b_i) \geq 0$  for  $i \neq 3$ . The invariant  $K_3$  defines a rational modular form  $\nu(K_3)$  of weight 3. Since it is not a multiple of  $\psi_1^3$  and  $\dim M_3(\Gamma_2) = 1$  it cannot be regular. Therefore,  $\text{ord}(b_3) = -1$ . ■

Using this lemma we obtain modular forms  $\psi_1 = \nu(K_1) \in M_1(\Gamma_2)$  and

$$\chi_{13} = \nu(K_3)\chi_{10} \in S_{13}(\Gamma_2)$$

as the form of  $K_3$  shows that  $\nu(K_3)$  has order  $-2$  along  $\mathcal{A}_{1,1}$  since  $K_1$  does not vanish on  $\mathcal{A}_{1,1}$ . We see that  $\chi_{13}$  is not divisible by  $\psi_1$ , hence  $\dim M_{12}(\Gamma_2) < \dim M_{13}(\Gamma_2)$ . In view of the estimates on the dimensions of  $M_k(\Gamma_2)$  implied by (2) we conclude that

$$\dim M_{12}(\Gamma_2) = 3 \quad \text{and} \quad \dim M_{13}(\Gamma_2) = 4. \tag{7}$$

By inspecting the expression of the invariant  $K_{10}$  in terms of the  $a_i$  and  $b_j$  we see that  $\nu(K_{10})$  is a modular form of weight 10 that vanishes with multiplicity 2 along  $\mathcal{A}_{1,1}$ . Therefore, by the cycle class of  $\mathcal{A}_{1,1}$  given in (1) it is a multiple of  $\chi_{10}$  and we normalize  $\chi_{10}$  by setting

$$\chi_{10} = \nu(K_{10}).$$

Consider  $K_{12} = K_3K_1^4 + K_3^4 + K_1^3K_3^3$ . It starts as follows

$$(a_0^2a_3^2 + a_0a_1a_2a_3 + a_0a_2^3 + a_1^3a_3)^4 b_3^4 + \dots$$

and one can check that it has order 0 along  $\mathcal{A}_{1,1}$ . This defines a modular form

$$\psi_{12} = \nu(K_{12}).$$

**Remark 5.2.** From the fact that  $\chi_{10}$  (resp.  $\psi_{12}$ ) vanishes with multiplicity 2 (resp. 0) along  $\mathcal{A}_{1,1}$  we see that the order of  $a_0$  and  $a_3$  along  $\mathcal{A}_{1,1}$  is 1.

Similarly,  $\nu(K_8)$  has order  $-8$  along  $\mathcal{A}_{1,1}$ , hence we find

$$\chi_{48} = \nu(K_8)\chi_{10}^4 \in S_{48}(\Gamma_2).$$

In this way we obtain modular forms  $\psi_1, \chi_{10}, \psi_{12}, \chi_{13}, \chi_{48}$ .

The relation  $K_{12} = K_8K_1^4 + K_3^4 + K_1^3K_3^3$  implies that

$$\chi_{10}^4\psi_{12} = \chi_{48}\psi_1^4 + \chi_{13}^4 + \chi_{13}^3\psi_1^3\chi_{10}. \quad (8)$$

Inside the ring  $\mathcal{R}_2(\mathbb{F}_2)$  we thus found generators

$$\psi_1, \chi_{10}, \psi_{12}, \chi_{13}, \chi_{48}$$

with a relation (8) of weight 52. These generate a subring  $R = \bigoplus_k R_k$  of  $\mathcal{R}_2(\mathbb{F}_2)$  with generating function for the dimensions

$$G = \frac{1 - t^{52}}{(1-t)(1-t^{10})(1-t^{12})(1-t^{13})(1-t^{48})}$$

and  $\dim R_k = k^3/1080 + O(k^2)$ . To see that  $R = \mathcal{R}_2(\mathbb{F}_2)$  one can use the degree of  $\text{Proj}(R)$

$$2 \deg(\lambda_1^3) = \frac{1}{1440} = \frac{52}{1 \cdot 10 \cdot 12 \cdot 13 \cdot 48},$$

(see [6]) or argue with the dimensions as follows. Let  $d(k) = \dim M_k(\Gamma_2)$  and  $r(k) = \dim R_k$ .

**Proposition 5.3.** We have  $d(k) = r(k)$  for  $k \geq 0$ .

**Proof.** We have  $d(k) \geq r(k)$  and we observed equality for  $k = 0, \dots, 13$  using the upper bound provided by (2), see (3) and (7). We assume by induction that we have  $d(k) = r(k)$  for  $k \leq m$ . Then by (2) we get an upper bound  $d(k) \leq r(k-10) + c(k)(c(k)+1)/2$  for  $k \leq m+10$  with  $c(k) = \dim M_k(\Gamma_1) = \lfloor k/12 \rfloor + 1$ . But we have  $r(k) - r(k-10) = c(k)(c(k)+1)/2$  for  $k \not\equiv 0, 1, 2 \pmod{12}$ . Indeed,

$$G - t^{10}G - \frac{1}{(1-t)(1-t^{12})^2} = -t^{12} \frac{t^{26} + t^{25} + t^{24} + t^{13} + t^{12} + 1}{t^{60} - t^{48} - t^{12} + 1}.$$

Hence,  $d(k) = r(k)$  for  $k \leq m + 10$  and  $k \not\equiv 0, 1, 2 \pmod{12}$ . But using  $d(k + 1) - d(k) \geq r(k + 1) - r(k)$ , to be proved in the next lemma, we get  $d(k) = r(k)$  for all  $k \leq m + 10$ . Induction finishes the proof. ■

**Lemma 5.4.** We have  $d(k + 1) - d(k) \geq r(k + 1) - r(k)$  for  $k \geq 0$ .

**Proof.** Because of (8) we have  $R_{k+1} = \psi_1 R_k \oplus N_{k+1}$  with  $N_{k+1}$  the subspace with basis the forms  $\chi_{10}^a \psi_{12}^b \chi_{13}^c \chi_{48}^d$  with  $10a + 12b + 13c + 48d = k + 1$  with  $a, b, c, d \geq 0$  and  $c \leq 3$ . We thus have  $r(k + 1) - r(k) = \dim N_{k+1}$ . We claim that  $N_{k+1} \cap \psi_1 M_k(\Gamma_2) = (0)$ , and this implies  $d(k + 1) - d(k) \geq \dim N_{k+1}$ . Suppose we have an  $f \in M_k(\Gamma_2)$  with  $f \notin R_k$  and  $\psi_1 f \in N_{k+1}$ . We can write  $\psi_1 f = P$  with  $P$  a polynomial in  $\chi_{10}, \psi_{12}, \chi_{13}, \chi_{48}$ , where no power of  $\chi_{13}$  is  $> 3$ . Then  $P = \nu(Q)$  with  $Q$  a polynomial in

$$K_{10}, K_8 K_1^4 + K_3^4 + K_3^3 K_1^3, K_3 K_{10}, K_8 K_{10}^4$$

that is divisible by  $K_1$  by assumption. In order that a non-zero  $Q$  is divisible by  $K_1$  we need at least one monomial of  $Q$  involving a power  $\geq 4$  of  $K_3 K_{10}$  since  $K_1, K_3, K_8$ , and  $K_{10}$  are algebraically independent. But this is excluded. ■

**Remark 5.5.** In characteristic different from 2 the invariant  $E$  of degree 15 gives rise to a generator, a modular form  $\chi_{35}$  of weight 35. But in our case, we get

$$K_{15} = K_3 K_{12} + K_{10} K_1^5$$

and  $\nu(K_{15})$  has order  $-2$  along  $\mathcal{A}_{1,1} \otimes \mathbb{F}_2$ , so if we set  $\chi_{25} = \chi_{10} \nu(K_{15}) \in S_{25}(\Gamma_2)$  we have

$$\chi_{25} = \chi_{13} \psi_{12} + \chi_{10}^2 \psi_1^5$$

and do not get a new generator.

**Remark 5.6.** The zero locus of  $\psi_1$  is the closure of the locus  $V_1$  of abelian surfaces of 2-rank  $\leq 1$ . The zero locus of the ideal  $(\psi_1, \chi_{13})$  consists of two components: the locus  $V_0$  of 2-rank 0, that is, the supersingular locus, and the closure of the intersection of  $\mathcal{A}_{1,1} \cap V_1$ , both with multiplicity 2. This fits the cycles classes:

$$13\lambda_1^2 = 3\lambda_1^2 + 10\lambda_1^2.$$

The class of  $V_0$  is  $3\lambda_2 = (3/2)\lambda_1^2$ , with  $\lambda_2$  the 2nd Chern class of  $\mathbb{E}$ , see [6, Thm. 2.4] and [4, Thm. 12.4].

### Funding

Part of the research of the first author was supported by the EPSRC Grant EP/N031369/1 and the Simons Collaboration in Arithmetic Geometry, Number Theory, and Computation via the Simons Foundation.

### References

- [1] Böcherer, S. and S. Nagaoka. "On mod  $p$  properties of Siegel modular forms." *Math. Ann.* 338 (2007): 421–33.
- [2] Cléry, F., C. Faber, and G. van der Geer. "Covariants of binary sextics and vector-valued Siegel modular forms of genus two." *Math. Ann.* 369 (2017): 1649–69.
- [3] Deligne, P. "Courbes elliptiques: Formulaire (d'après J. Tate)." *Modular Functions IV. Lecture Notes in Mathematics*, vol. 476, Berlin: Springer Verlag, (1975): 53–73.
- [4] Ekedahl, T. and G. van der Geer. "Cycle classes of the E-O stratification on the moduli of abelian varieties." *Algebra, Arithmetic, and Geometry*, Y. Tschinkel and Y. Zarhin (eds.), Progress in Mathematics 269. Basel: Birkhäuser, (2010).
- [5] Faltings, G. and C. L. Chai. "Degeneration of abelian varieties." *Ergebnisse der Math*, vol. 22. Springer, 1990.
- [6] van der Geer, G. "Cycles on the moduli space of abelian varieties." *Moduli of Curves and Abelian Varieties, Aspects Math.*, E33. Braunschweig: Vieweg, (1999).
- [7] van der Geer, G. "The ring of modular forms of degree two in characteristic three." arXiv:1912.05161.
- [8] Hartshorne, R. *Algebraic Geometry*. Graduate Texts in Mathematics, vol. 52. New York: Springer Verlag, 1977.
- [9] Igusa, J.-I. "Arithmetic variety of moduli for genus two." *Ann. of Math.* 72 (1960): 612–49.
- [10] Igusa, J.-I. "On the ring of modular forms of degree two over  $\mathbb{Z}$ ." *Am. J. Math.* 101 (1979): 149–83.
- [11] Ichikawa, T. "Siegel modular forms modulo  $p$ ." *Proceeding of the 4th Spring Conference Siegel Modular Forms and Abelian Varieties*, edited by Ibukiyama T., 105–16, 2007.
- [12] Liu, Q. "Courbes stables de genre 2 et leur schéma de modules." *Math. Ann.* 295 (1993): 201–22.
- [13] Liu, Q. "Modèles minimaux des courbes de genre deux." *J. Reine Angew. Math.* 453 (1994): 137–64.
- [14] Mumford, D. "Towards an enumerative geometry of the moduli space of curves." *Arithmetic and Geometry*, Papers dedicated to I. R. Shafarevich, Vol. II: Geometry. Progress in Math. 36, pp. 271–328.
- [15] Nagaoka, S. "Note on mod  $p$  Siegel modular forms." *Math. Z.* 235 (2000): 405–20.
- [16] Nagaoka, S. "Note on mod  $p$  Siegel modular forms, II." *Math. Z.* 251 (2005): 821–6.
- [17] Zagier, D. "Elliptic modular forms and their applications." *The 1-2-3 of Modular Forms, Universitext*, 1–103. Berlin: Springer, 2008.