



UvA-DARE (Digital Academic Repository)

Combinatorial RNAi against HIV-1

Liu, Y.P.

[Link to publication](#)

Citation for published version (APA):

Liu, Y. P. (2009). Combinatorial RNAi against HIV-1

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <http://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Chapter 4

Inhibition of HIV-1 by multiple siRNAs expressed from a single microRNA polycistron

Ying Poi Liu
Joost Haasnoot
Olivier ter Brake
Ben Berkhout
Pavlina Konstantinova

Laboratory of Experimental Virology,
Department of Medical Microbiology,
Center for Infection and Immunity Amsterdam (CINIMA),
Academic Medical Center, University of Amsterdam,
The Netherlands

Abstract

RNA interference (RNAi) is a powerful approach to inhibit human immunodeficiency virus type 1 (HIV-1) replication. However, HIV-1 can escape from RNAi-mediated antiviral therapy by selection of mutations in the targeted sequence. To prevent viral escape, multiple small interfering RNAs (siRNAs) against conserved viral sequences should be combined. Ideally, these RNA inhibitors should be expressed simultaneously from a single transgene transcript. In this study, we tested a multiplex microRNA (miRNA) expression strategy by inserting multiple effective anti-HIV siRNA sequences in the miRNA polycistron mir-17-92. Individual anti-HIV miRNAs that resemble the natural miRNA structures were optimized by varying the siRNA position in the hairpin stem to obtain maximal effectiveness against luciferase reporters and HIV-1. We show that an antiviral miRNA construct can have a greater intrinsic inhibitory activity than a conventional short hairpin (shRNA) construct. When combined in a polycistron setting, the silencing activity of an individual miRNA is strongly boosted. We demonstrate that HIV-1 replication can be efficiently inhibited by simultaneous expression of 4 antiviral siRNAs from the polycistronic miRNA transcript. These combined results indicate that a multiplex miRNA strategy may be a promising therapeutic approach to attack escape-prone viral pathogens.

Introduction

RNAi is an evolutionary conserved and sequence-specific gene silencing mechanism in eukaryotes (1,2). RNAi can be induced by double-stranded RNA that is processed by the RNase III-like enzyme Dicer into 21-25 bp siRNAs (3-5). The siRNA is incorporated into the RNA-induced silencing complex (RISC) in the cytoplasm and directs RISC to degrade an mRNA that is perfectly complementary to one (guide) strand of the siRNA (5).

Cellular miRNAs are the natural inducers of RNAi. Most miRNAs are synthesized as part of longer primary RNA transcripts (pri-miRNAs)(6-8). The pri-miRNAs are cleaved by the nuclear Drosha-DGCR8 complex to produce miRNA precursors (pre-miRNAs) of ~70 nt (9-12). Pre-miRNAs are transported by Exportin-5 to the cytoplasm, where they are cleaved by Dicer to produce the miRNA duplex of ~22 bp. The single-stranded mature miRNA targets RISC for mRNA cleavage (perfect complementarity) or translational repression (incomplete complementarity)(13,14). Several miRNAs are encoded in genomic clusters that are transcribed as polycistronic pri-miRNAs, allowing the production of multiple miRNAs from a single transcription unit (15,16). RNAi can be induced using RNA polymerase III promoter-driven shRNA expression vectors that direct siRNA expression. Another siRNA expression strategy uses a pre-miRNA backbone that is transcribed by polymerase II or III (17-20). An optimized pre-miRNA design includes the single stranded flanks of the pri-miRNA (21-23).

Currently, RNAi has been employed to inhibit the replication of a wide range of viruses including HIV-1, hepatitis C virus (HCV), hepatitis B virus (HBV), dengue virus, poliovirus, influenza virus A, coronavirus, herpesvirus and picornavirus (24,25). For HIV-1, potent inhibition has been reported with single shRNA and miRNA expression constructs (17,26-28). However, the therapeutic use of a single inhibitor is limited because of the rapid emergence of HIV-1 escape mutants (27,29,30). Minor sequence changes in the target sequence, sometimes even a single point mutation, are sufficient to overcome RNAi-mediated inhibition (30), thus demonstrating the exquisite sequence-specificity of RNAi.

Strategies to reduce the chance of viral escape include the simultaneous use of multiple shRNAs in a combinatorial RNAi approach, which increases the genetic barrier for viral escape (31,32). Similar strategies have previously been validated for antisense DNA and ribozymes (33-35). However, expression of these shRNAs necessitates multiple expression cassettes and the construction of rather complex vectors that will not easily provide equimolar shRNA expression levels. Furthermore, when the same promoter is reiterated in a lentiviral vector, recombination occurs with high frequency on the repeated sequences (36). Alternative anti-escape strategies include the use of a second-generation of siRNAs that target specific escape variants (37), the use of tandem siRNA transcripts (38), long hairpin RNAs (39) or the targeting of cel-

lular co-factors that are critically involved in viral replication (40-44). Another attractive approach is to express multiple antiviral siRNAs from a single polycistronic miRNA transcript, such as a natural genomic miRNA cluster that can be expressed from an RNA polymerase II promoter. This strategy is of particular interest for antiviral purposes because miRNA-like transcripts were shown to be more effective antivirals than regular shRNAs (17,45). Furthermore, using an RNA polymerase II promoter will allow lower and regulated expression, thereby reducing the risk of toxicity due to saturation of the RNAi machinery (46).

In this study, we designed a polycistronic transcript based on the mir-17-92 backbone to simultaneously express 4 anti-HIV siRNAs. To generate this transcript, we first constructed individual anti-HIV miRNAs that resemble the natural pri-miRNA structures. These hairpins were optimized for viral inhibition by varying the siRNA position in the hairpin stem. We show that the expression of individual miRNAs is greatly enhanced in multiplex hairpin transcripts that are properly processed into functional miRNAs. HIV-1 replication can be potently inhibited by simultaneous expression of 4 antiviral miRNAs. These combined results indicate that the multiplex miRNA strategy is a promising therapeutic approach against escape-prone viral pathogens.

Materials and methods

DNA constructs

The wild-type mir-17-19b polycistron was amplified from genomic DNA of 293T cells with primers OncoF1 and OncoR1 and TA-cloned in TOPO2.1 (Invitrogen). The mir-17-19b polycistron sequence was verified and is identical to the sequence in the NCBI database (NT_009952.14). The TOPO2.1/mir-17-19b construct was used as a template to generate the antiviral miRNA constructs. The construction consists of a 4-step fusion PCR as shown in the Supplementary Fig. 1. Briefly, the 5' flank of the pri-miRNA was amplified with a forward primer encoding a BamHI site and a reverse primer encoding the HIV-1 sequence at its 3' end (step 1). Similarly, the 3' flank of the pri-miRNA was amplified with a forward primer containing HIV-1 sequences and a reverse primer encoding BglII and XhoI sites (step 2). Two complementary oligonucleotides, creating the stem-loop structure of the antiviral miRNA, were annealed as described previously (step 3)(32). The partial sequence similarity between the fragments generated in step 1, 2 and 3 allowed their fusion by PCR with the outer forward F1 and reverse R2 primers (step 4), resulting in the antiviral miRNA. Supplementary Table 1 lists all oligonucleotides used in this study.

PCR amplification was performed in a 50 µl reaction containing 1 × PCR amplification buffer (Invitrogen), 1.5-2 mM MgCl₂ (optimized for each reaction), 25 pmol of each primer, 0.2 mM dNTPs and 2.5 units of AmpliTaq DNA polymerase (Perkin Elmer Applied Biosystem). The PCR program was as follows: 95°C for 5 min, 35 cycles of 2 min at 94°C, 1.5 min at 60°C, 2.5 min at 72°C and a final extension for 8 min at 72°C. The PCR products were separated on a 1% agarose gel stained with ethidium bromide and compared to a standard DNA size marker (Eurogentec). miRNA PCR products were excised from gel, purified with the QIAquick gel extraction kit (Qiagen),

digested with BamHI and XhoI and cloned in the corresponding sites of pcDNA6.2-GW/EmGFP-miR (Invitrogen). All miRNA constructs were sequenced with primers GFPseqF and mirR using the BigDye Terminator v1.1 Cycle Sequencing Kit (Perkin Elmer Applied Biosystem).

Multimerization of the individual pri-miRNA units was performed by digestion of a single miRNA hairpin construct with BamHI and XhoI and religation into the BglII/XhoI sites of pcDNA6-miRNA. By repeating this procedure we obtained constructs expressing different combinations of 1, 2, 3, 4 and 6 pri-miRNAs. The RNA structures formed by the transcripts were predicted with the Mfold program (47) at <http://frontend.bioinfo.rpi.edu/applications/mfold/> and found to be similar to the predicted conformation of the wild-type pri-miRNAs.

The firefly luciferase (FL) reporters containing HIV-1 target sequences pol47 (Luc-A^{pol47}), pol1 (Luc-B^{pol1}), gag5 (Luc-C^{gag5}), r/t5 (Luc-D^{r/t5}), ldr9 (Luc-E^{ldr9}) and the anti-HIV shRNAs have been described previously (32). The full-length HIV-1 molecular clone LAI (accession number AF33819.3) (48) was used to produce wild-type virus and to study its inhibition by the antiviral miRNAs and shRNAs.

Cells, DNA transfection and virus infection

Human embryonic kidney (HEK) 293T cells were grown as a monolayer in DMEM (Invitrogen) supplemented with 10% fetal calf serum (FCS) (Hybond), minimal essential medium nonessential amino acids, penicillin (100 U/ml) and streptomycin (100 µg/ml) at 37°C and 5% CO₂. Cells were trypsinized one day before transfection, resuspended in DMEM without antibiotics and seeded in 24-well plates at a density of 1.2 × 10⁵ cells per well. Cells were co-transfected with the indicated DNA constructs using Lipofectamine 2000 reagent (Invitrogen) according to the manufacturer's instructions. 1 ng of pRL plasmid (Promega) expressing renilla luciferase (RL) from the CMV promoter was added as an internal control for cell viability and transfection efficiency. All transfection experiments were controlled for equal DNA input by adding pBluescript SK- (Promega). Luciferase and renilla expression was measured with the dual-luciferase reporter assay system (Promega) according to the manufacturer's instructions. Virus production was determined by measuring the CA-p24 levels in the culture supernatant by ELISA (Abbott) (49). Subsequently, the cells were lysed to measure the renilla luciferase activities using the Renilla Luciferase Assay System (Promega) according to the manufacturer's protocol. Transfection experiments were corrected for between session variations as described previously (50).

The human T cell line SupT1 was cultured in 25 cm² flasks in RPMI 1640 medium supplemented with 10% FCS, penicillin (100 U/ml) and streptomycin (100 µg/ml) at 37°C and 5% CO₂. 200.000 SupT1 cells were infected with equal amounts virus (0.5 ng CA-p24) produced in 293T cells. When HIV-induced cytopathic effects were observed, cell and supernatant samples were stored at -80°C. Virus spread was followed by measuring the CA-p24 levels in the culture supernatant by ELISA.

siRNA detection by Northern blotting

One day before transfection 7.5 × 10⁵ 293T cells were plated in 6-well plates. Cells were transfected with 39 – 5000 ng of the shRNA construct and 2500 or 5000 ng of the miRNA-expression construct using Lipofectamine 2000 reagent (Invitrogen) according to the manufacturer's instructions. In all transfection experiments we added pBluescript SK- (Promega) to obtain identical DNA concentrations. Total cellular RNA was extracted two days post-transfection with the mirVana miRNA isolation kit (Ambion) according to the manufacturer's protocol. For Northern blot analysis, 15 µg

total RNA was heated for 5 min at 95°C before electrophoresis on a 15% denaturing polyacrylamide gel (precast Novex TBU gel, Invitrogen). To check for equal sample loading, the gel was stained with 2 µg/ml ethidium bromide in milliQ water for 20 min. Destaining was performed by rinsing the gel three times with milliQ water for 10 min. The ribosomal RNA (rRNA) bands were visualized under UV light. The RNA samples were electrotransferred to a positively charged nylon membrane (Boehringer Mannheim, GmbH, Mannheim, Germany) and crosslinked to the membrane using UV light at a wavelength of 254 nm (1200 µJ x 100). LNA oligonucleotide probes were 5' end labeled with the kinaseMax kit (Ambion) in the presence of 1 µl [γ -³²P]ATP (0.37 MBq/µl Amersham Biosciences). To remove unincorporated nucleotides, the probes were purified on Sephadex G-25 spin columns (Amersham Biosciences) according to the manufacturer's protocol. Hybridizations were performed at 42°C with labeled LNA oligonucleotides in 10 ml ULTRAhyb hybridization buffer (Ambion) according to the manufacturer's instructions. We used the oligonucleotide probes (LNA-positions underlined): 5' GTGAAGGGGCAGTAGTAAT 3' (pol47 probe), 5' ACAGGAGCAGATGATACAG 3' (pol1 probe), 5' GAAGAAAATGATGACAGCAT 3' (gag5 probe), 5' ATGGCAGGAAGAAGCGGAG 3' (r/t5 probe) and 5' AGATGGGTGCGAGAGCGTC 3' (ldr9 probe). The membranes were washed twice for 5 min at 42°C in 2 x SSC/0.1% SDS and twice for 15 min at 42°C in 0.1 x SSC/0.1% SDS. Signals were detected and quantified using a phosphorimager (Amersham Biosciences).

Lentiviral vector production and transduction

Lentiviral vector plasmids are derived from the construct pLenti6/V5-Dest (Invitrogen), which we renamed pLV. The miRNA cassettes, containing a GFP marker, were inserted into pLV using the Gateway-adapted BLOCK-iT Lentiviral Pol II miR RNAi Expression System (Invitrogen) according to the manufacturer's instructions. The sequences of all miRNA constructs were verified using the primers CMVf and V5r. The miRNA inhibitory potential was determined by co-transfection with appropriate luciferase reporters (results not shown). Lentiviral vector was produced in 293T cells (2.2×10^6) seeded in a 25 cm² flask. The next day, medium was replaced with 2.2 ml medium without antibiotics. Subsequently, pLV vectors expressing a miRNA (2.4 µg) were co-transfected with packaging plasmids pSYNGP (1.5 µg)(51), pRSV-rev (0.6 µg) and pVSVg (0.8 µg)(52) with 16 µl of Lipofectamine 2000 reagent and 1.5 ml Optimem (Gibco BRL). The second day, medium was replaced with fresh medium. On the third and fourth day, medium containing lentiviral vector was harvested and pooled. Cellular debris was removed by filtration through a FP30/0.45 CA-S filter (Schleicher & Schuell MicroScience) and 1 ml aliquots were stored at -80°C. Lentiviral stocks were titrated on SupT1 and 293T cells to determine the vector titer. SupT1 and 293T cells (1×10^5) were transduced with the pLV vector expressing an unrelated miRNA N (Invitrogen) or the antiviral miRNAs A^{pol47}, B^{pol1}, C^{gag5}, D^{r/t5}, E^{ldr9} and ACDE at a multiplicity of infection (MOI) of 0.15 as described previously (32). The pLV vector contains the blasticidin resistance gene, which allows the selection of transduced cells using 3,5 µg/ml blasticidin. The GFP marker encoded by the miRNA cassette was used to select GFP+ cells by FACS sorting approximately 10 days post-transduction.

Results

Design of antiviral miRNAs

The human mir-17-92 cluster on chromosome 13 encodes a 1 kb pri-miRNA polycistronic transcript with six pre-miRNAs that produces seven mature miRNAs (Fig. 1A, upper panel)(15). The pre-miR-17 hairpin encodes two miRNAs, one on the 5' side of the hairpin (miR-17-5p) and one on the 3' side (miR-17-3p). We amplified the sequences encoding the first five pri-miRNAs from the mir-17-92 polycistron and cloned it under the control of the cytomegalovirus (CMV) immediate early promoter (Fig. 1A, lower panel). We subcloned each individual pre-miRNA with at least 40-nt flanks on each side of the hairpin and systematically replaced the mature miRNA sequences with antiviral sequences as explained in detail in the Supplementary Fig. 1. The original miRNA names were replaced with letters A-E and we inserted 19- to 24-nt antiviral sequences against five HIV-1 genes (Fig. 1B, upper panel). The HIV-1 targets represent highly conserved sequences to which we successfully raised potent shRNA inhibitors (32). We set out to combine 2, 3, 4 or 5 antiviral miRNAs, which will eventually result in an antiviral pri-miRNA polycistron (Fig. 1B, lower panel).

We first determined if we could produce an optimal antiviral miRNA construct by varying the position and length of the anti-HIV sequence. For the initial experiments we chose miRNAs miR-20 and miR-19b (D^{wt} and E^{wt}) because they contain the smallest number of bulges and thus most resemble the original shRNA structure (Fig. 2). The original mature miRNA and the antiviral miRNA strand are boxed. We modified the passenger strand of the base paired stem to mimic structural features (mismatches, bulges and thermodynamic stability) of the natural pre-miRNA. We constructed a series of antiviral miRNAs with the effective r/t5 and ldr9 siRNAs in the D^{wt} backbone, which produces the antiviral guide strand from the 5' side of the hairpin duplex. The 19-nt antiviral siRNAs were positioned either at the 5' ($D^{r/t5-19}$ and $D^{ldr9-19}$) or 3' end ($D^{r/t5-19'}$ and $D^{ldr9-19'}$) of the original miRNA sequence (Fig. 2, upper panel). Additional constructs were made in which we extended the antiviral siRNAs at the 3' end to 23-nt, which is the actual size of the wild-type mature miRNA ($D^{r/t5-23}$ and $D^{ldr9-23}$) (Fig. 2, Table 1). We repeated this strategy for the r/t5 inhibitor in the E^{wt} hairpin, which produces the antiviral guide strand from the 3' side of the hairpin stem (Fig. 2, lower panel). We designed a similar set of constructs in which the 19-nt r/t5 siRNA was positioned at the 5' ($E^{r/t5-19}$) or 3' end ($E^{r/t5-19'}$) of the wild-type miRNA and an extended 23-nt version ($E^{r/t5-23}$).

Activity of the antiviral miRNAs

To determine the inhibitory activity of the miRNA constructs, we co-transfected 293T cells with the inhibitors and luciferase reporter constructs containing either the 50-nt r/t5 target (Luc- $D^{r/t5}$ and Luc- $E^{r/t5}$) or the 50-nt ldr9 target (Luc- D^{ldr9}). A plasmid encoding renilla luciferase (pRL) was included to correct for




Figure 1. Schematic of the mir-17-92 cluster and design of the antiviral miRNAs. (A) Genomic organization of the mir-17-92 polycistron (upper panel) and predicted secondary structure of the mir-17-19b transcript (A^{wt}-E^{wt}, lower panel). Mature miRNAs are indicated in red; CMV, cytomegalovirus immediate early promoter; pA, SV40 polyadenylation signal. (B) Scheme of the HIV-1 proviral genome and position of the target sequences (upper panel) used for the design of the antiviral miRNA polycistron (lower panel). Guide strands of the siRNA against HIV-1 are blue. Numbers between the pre-miRNAs represent the length of the flanks (in nucleotides). Original miRNA names were replaced with the letters A-E and target genes are indicated in superscript.

transfection variation and to monitor for cell viability that may be affected by off-target effects of the modified miRNAs and/or the antiviral siRNAs. Firefly and renilla luciferase expression was measured 48 hrs post-transfection. Firefly luciferase expression was normalized to the control renilla luciferase expression. Firefly luciferase expression in the presence of pBluescript (pBS) was set at 100% (Fig. 3A). A common pattern was observed in that most efficient inhibition was scored for the 19- and 23-nt siRNA positioned at the 5' end of the original miRNA sequence, which are the 19 and 23 variants of the D hairpin and the 19' and 23 variants of the E hairpin.

A similar trend was observed in HIV-1 inhibition studies (Fig. 3B). We co-transfected the HIV-1 molecular clone LAI and the miRNA constructs into 293T cells and virus production was measured as the CA-p24 level in the culture supernatant at 2 days post-transfection. We observed the strongest inhibition when the siRNA inhibitor was situated at the 5' end of the miRNA sequence. The optimized hairpins have a similar efficiency of inhibiting HIV-1 production as the shRNA constructs that were used as positive controls.




Figure 2. Structure of the antiviral miRNAs based on pre-miRNAs D^{wt} and E^{wt}. Upper panel: The r/t5 and ldr9 siRNAs were incorporated into the D^{wt} backbone, which produces the guide strand from the 5' side of the hairpin. The guide strand is marked in grey. We modified the passenger strand to mimic structural features (mismatches, bulges and thermodynamic stability) of the natural pre-miRNA. The 19-nt antiviral siRNAs were positioned either at the 5' (D^{r/t5-19} and D^{ldr9-19}) or 3' end (D^{r/t5-19'} and D^{ldr9-19'}) of the original miRNA and the length of the siRNAs was extended at the 3' end to 23-nt (D^{r/t5-23} and D^{ldr9-23}). Lower panel: The r/t5 siRNA was similarly incorporated into the E^{wt} pre-miRNA, which produces the guide strand from the 3' side of the hairpin. The structure of the original shRNAs sh^{r/t5} and sh^{ldr9} are presented. HIV-1 sequences are blue, mature wild-type miRNA sequences are red, pre-miRNA sequences are black, Watson-Crick base pairs are shown with dashes and G•U wobbles with dots.

Based on these initial findings, we designed additional miRNA constructs with 19-24 nt antiviral sequences at the 5' end of the miRNA: A^{pol47-24}, A^{pol47-19}, B^{pol1-22}, B^{pol1-19}, C^{gag5-23}, C^{gag5-19'}, E^{ldr9-23} and E^{ldr9-19'} (Table 1). Of the original D^{r/t5}, D^{ldr9} and E^{r/t5} constructs, we selected D^{r/t5-19} because it is the best inhibitor. As a consequence, the ldr9 inhibitor was introduced in the E hairpin. We tested the effectiveness of all new constructs against luciferase reporters and HIV-1 (Table 1). The observed inhibition is sequence-specific because non-matching miRNAs did not have any effect. Furthermore, the expression of the control renilla reporter was stable in all experiments. The miRNA constructs with 19-nt anti-HIV sequences showed a somewhat higher activity than the extended versions. We therefore selected the 19-nt inhibitors for the construction of antiviral miRNA polycistrons. For simplicity, we removed the 19 indications from the miRNA names (e.g. A^{pol47-19} becomes A^{pol47}).

Expression and processing of an antiviral miRNA

The original shRNA inhibitor $sh^{r/t5}$ seems slightly more active than the optimized $D^{r/t5}$ miRNA molecule (Fig. 3). This could be due to the use of different promoters (polymerase III U6 versus polymerase II CMV, respectively), but may also be due to differential RNA processing efficiencies. We therefore wanted to compare the siRNA level expressed from each construct. We titrated 39 – 2500 ng $sh^{r/t5}$ construct in 293T cells and compared that with 2500 ng $D^{r/t5}$ construct. Two days post-transfection, total cellular RNA was isolated from the transfected cells and analyzed on a Northern blot with an $r/t5$ probe




Figure 3. Inhibition of luciferase reporters and HIV-1 production by individual antiviral miRNAs. (A) Luciferase reporters (100 ng) were co-transfected in 293T cells with 20 ng antiviral miRNA construct and 1 ng pRL. The original shRNAs $sh^{r/t5}$ and sh^{ldr9} were used as positive controls and non-matching miRNAs as negative controls. Firefly and renilla luciferase expression was measured two days post-transfection. Normalized firefly luciferase expression in the presence of pBS was set to 100%. Mean values obtained in three independent experiments are shown (\pm SE). FL, firefly luciferase; RL, renilla luciferase. (B) 293T cells were co-transfected with 150 ng HIV-1 molecular clone LAI, 1 ng pRL and 20 ng of the indicated antiviral miRNA constructs. $sh^{r/t5}$ and sh^{ldr9} constructs were used as positive controls, D^{wt} and E^{wt} as negative controls. CA-p24 levels in the culture supernatant and renilla luciferase expression were measured two days post-transfection. Normalized CA-p24 expression in the presence of pBS was set at 100%. Error bars represent the standard deviation in four independent experiments.

that detects the guide (antisense) strand (Fig. 4). The original D^{wt} hairpin was used as a negative control. Quantification of the RNA bands showed that expression of the antiviral 19-nt siRNA by construct $D^{r/t5}$ is about 15-30 fold lower than the expression by construct $sh^{r/t5}$ (Fig. 4, see numbers below the blot). Although the siRNA expression level of $D^{r/t5}$ is significantly lower than that of $sh^{r/t5}$, the inhibitory effect measured in the luciferase and HIV-1 inhibition assays is comparable. This result suggests that the intrinsic inhibitory capacity of $D^{r/t5}$ is in fact much greater than that of $sh^{r/t5}$.




Figure 4. Expression and processing of siRNA from a miRNA versus an shRNA construct. We titrated 39 - 2500 ng $sh^{r/t5}$ construct in 293T cells and compared that with 2500 ng $D^{r/t5}$ construct. Northern blot analysis was performed on total cellular RNA. M, RNA size marker (nt) is shown on the left. Expression of siRNAs by 2500 ng $sh^{r/t5}$ was set at 100% and relative siRNA expression levels are indicated below the blot. Ethidium bromide staining of the 5S rRNA serves as sample loading control.

Multimerization of miRNAs increases the inhibitory activity

To address the effect of chaining of different hairpins for the silencing activity of the individual hairpins, we coupled the A^{pol47} inhibitor to wild-type pri-miRNAs. We constructed two variants of A^{pol47} with 2 or 6 hairpins in a single transcript: $A^{pol47}B^{wt}$ and $A^{pol47}A^{wt-E^{wt}}$ ($A^{wt-E^{wt}}$ represents the complete wild-type mir-17-19b polycistron). We tested the ability of these hairpins to inhibit the Luc- A^{pol47} reporter (Fig. 5A, left). Firefly luciferase expression was normalized to the renilla luciferase expression from the co-transfected pRL plasmid. We set the luciferase expression in the presence of pBS at 100%. The knockdown efficiency of the single A^{pol47} hairpin was ~40%, but chaining it with the B^{wt} hairpin or the $A^{wt-E^{wt}}$ cluster enhanced the silencing activity. As a negative control, we included the $A^{wt-E^{wt}}$ construct. As a positive control, we included the original shRNA. A similar pattern was observed for inhibition of HIV-1 production (Fig. 5A, right). Thus, expression of a miRNA inhibitor as part of a miRNA polycistronic transcript enhances the silencing activity.

Next, we constructed polycistronic hairpin constructs with different combinations of 2, 3 or 4 hairpins of the miRNA inhibitors A^{pol47} , B^{pol11} , C^{gag5} , $D^{r/t5}$ and E^{ldr9} : AB, AC, AD, ACC, ACD, ACCD, ACDE and ACDB^{wt}. We first determined the knockdown efficiency of each individual hairpin within the multiplex transcripts by co-transfection with the corresponding luciferase report-

Table 1. Structure and properties of the anti-HIV miRNAs

Name	Sequence (5'-3') and predicted structure	ΔG (kcal/mole)	Effectiveness		siRNA expression
			Luc	HIV	
A ^{wt}	GUCA ^{GA} AUAAGU ^{CA} AAGUCU ^A CA ^G UGCA ^G UAG ^U UG ^A _U CAGU ^{GG} UAUWACG ^{AUG} UUCACGU ^U AGC ^U AC ^U AC ^U _U	-34.3	-	-	-
A ^{po47-24}	GUCA ^{GA} AUAAGU ^{AU} UACUACU ^G CC ^C UUCAC ^G UUU ^U CG ^{AD} _U CAGU ^{GG} UAUWACG ^{AUG} AUGAUGGC ^U GG ^U AAGUG ^U AAA ^U GC ^U _U	-30.2	+	+	nd
A ^{po47-19}	GUCA ^{GA} AUAAGU ^{AU} UACUACU ^G CC ^C UUCAC ^G UAG ^U UG ^{AD} _U CAGU ^{GG} UAUWACG ^{AUG} AUGAUGGC ^U GG ^U AAGUG ^U ABC ^U AC ^U _U	-31.3	++	++	+
sh ^{po447}	UAAUGAGGACGGGGAAGU ^{UUC} _A UUUUAUACGCGCCCCUUCAC ^{AGA} _G	-35.8	+++	+++	+++
B ^{wt}	UUUU ^{CU} AAAG ^U GCA ^U UAG ^U GCA ^U UAG ^U UGAA ^{AG} _A UACGG ^{UUUC} CGU ^U GA ^U AUC ^U CGUC ^U AGC ^U CG ^U _U	-22.5	-	-	-
B ^{po11-22}	UUUU ^{CC} UUUA ^U CAU ^U CGU ^U CGUU ^U UGG ^U UGAA ^{AG} _U UACGG ^{UC} ACA ^U UUA ^U AC ^U CGA ^U GACA ^U AGC ^U CG ^U _U	-16.7	-	-	nd
B ^{po11-19}	UUUU ^{CC} UUUA ^U CAU ^U CGU ^U CGUU ^U UGG ^U UGAA ^{AG} _U UACGG ^{UC} ACA ^U UUA ^U AC ^U CGA ^U GACA ^U AUC ^U CG ^U _U	-20.3	-	-	-
sh ^{po11}	GACAUAGUAGACGAGGACA ^{UUC} _A UUUCUUUAUCACUUCUUCUUU ^{AGA} _G	-36.5	+++	+++	+++
C ^{wt}	GCAG ^U CC ^U UUUU ^U UAGUUUUUCAUAG ^U UUUCAC ^U UACA ^{AG} _A CGUC ^{CG} GG ^U GU ^U AGUCAAAACGUAC ^U AAACGU ^U UUG ^U _U	-39.1	-	-	-
C ^{gag5-23}	GCAG ^U CC ^U UUUU ^U UAGUUUUUCAUAG ^U UUCAC ^U UACA ^{AG} _A CGUC ^{CG} GG ^U GU ^U UAGUCAUUCUUCUUUACU ^U UUUCGU ^U AGUU ^U _U	-40.7	+	+	nd
C ^{gag5-19'}	GCAG ^U CC ^U UUUU ^U UAGUUUUUCAUAG ^U UUCAC ^U UACA ^{AG} _A CGUC ^{CG} GG ^U GU ^U UAGUCAUUCUUCUUUACU ^U UUUCGU ^U AGUU ^U _U	-40.3	++	+	+
sh ^{gag5}	GAGAAUUGUAGACCA ^{UUC} _A UUUCUUUUUACUACUGUCUUA ^{AGA} _G	-31.8	+++	+++	+++
D ^{wt}	GUAG ^C ACU ^A AAGUCU ^U UAGUAGU ^G CA ^G UAG ^U UG ^U _U CGUC ^U UGA ^{AA} UUCACGAGUUUUAUCUCU ^U AGU ^U AD ^U _A	-31.0	-	-	-
D ^{f15-23}	GUAG ^C ACC ^U CCUCUUCUUCUACCA ^U UGA ^U UG ^U _U CGUC ^U UGG ^U UGCAGAGGAGUUUCGUA ^U CUU ^U AD ^U _A	-42.1	++	++	nd
D ^{f15-19}	GUAG ^C ACC ^U CCUCUUCUUCUACCA ^U UAG ^U UG ^U _U CGUC ^U UGG ^U UGCAGAGGAGUUUCGUA ^U AUC ^U AD ^U _A	-39.0	+++	+++	+++
D ^{f15-19'}	GUAG ^C ACC ^U A ^C CCUCUUCUUCUACCA ^U UAG ^U UG ^U _U CGUC ^U UGA ^{AA} UGCAGAGGAGUUUCGUA ^U UAC ^U AD ^U _A	-38.2	+	+	nd
sh ^{f15}	AUUCGAGGAAAGCCGAG ^{UUC} _A UUUACCCUUCUUCUUCUUCU ^{AGA} _G	-39.5	+++	+++	+++
D ^{ldr9-23}	GUAG ^C ACU ^A CGUCUUCUACCCCAU ^U UCU ^U UG ^U _U CGUC ^U UGA ^{AA} UGCAGAGUUUUUUUAGA ^U AGA ^U AD ^U _A	-38.0	++	++	nd
D ^{ldr9-19}	GUAG ^C ACU ^A CGUCUUCUACCCCAU ^U UG ^U AG ^U UG ^U _U CGUC ^U UGA ^{AA} UGCAGAGUUUUUUUAGA ^U GA ^U UA ^U AD ^U _A	-41.0	++	++	nd
D ^{ldr9-19'}	GUAG ^C ACU ^A AGACUCUUCUACCCCAU ^U UG ^U UG ^U _U CGUC ^U UGA ^{AA} UUCGAGAGUUUUUUUUU AU ^U AD ^U _A	-38.7	-	+	nd
sh ^{ldr9}	AGAUGUUUUUUUAGAGCCUU ^{UUC} _A UUUCUACCCCAUUCUUCUUCU ^{AGA} _G	-34.7	+++	+++	+++
E ^{wt}	CACU ^U CUA ^U UUUUUAGUUUUUGCA ^U GG ^U UUUCA ^U CA ^U GC ^U _U GUGAU ^U GGUUU ^U CAGUCAAAACGU ^U CC ^U AAACGU ^U GU ^U CG ^U _U	-38.4	-	-	-
E ^{f15-23}	CACU ^U CUA ^U UUUUUAGUUUUUGAG ^U AA ^U AAGUU ^U CA ^U GC ^U _U GUGAU ^U GGUUU ^U CAGUCAAAACGU ^U CC ^U UUUCU ^U GU ^U CG ^U _U	-44.0	+	+	nd
E ^{f15-19}	CACU ^U CUA ^U UUUUUAGUUUUUGAG ^U AG ^U GAAGU ^U CA ^U GC ^U _U GUGAU ^U GGUUU ^U CAGUCAAAACGU ^U CC ^U CCUCU ^U GU ^U CG ^U _U	-44.6	+	+	nd
E ^{f15-19'}	CACU ^U CUA ^U UUUUUAGUUUUUGAG ^U AA ^U AA ^U U ^U CA ^U GC ^U _U GUGAU ^U GGUUU ^U CAGUCAAAACGU ^U CC ^U UUUCU ^U GU ^U CG ^U _U	-39.0	+	+	nd
E ^{ldr9-23}	CACU ^U CUA ^U UUUUUAGUUUUUGAG ^U UU ^U AGAGUUU ^U CA ^U GC ^U _U GUGAU ^U GGUUU ^U CAGUCAAAACGU ^U CC ^U UUUCU ^U GU ^U CG ^U _U	-44.0	-	-	nd
E ^{ldr9-19'}	CACU ^U CUA ^U UUUUUAGUUUUUGAG ^U UU ^U AGAGUUU ^U CA ^U GC ^U _U GUGAU ^U GGUUU ^U CAGUCAAAACGU ^U CC ^U UUUCU ^U GU ^U CG ^U _U	-44.4	+	+	+

HIV-1 sequences are blue; pre-miRNA sequences are black; mature miRNA sequences are red; guide strand is bold; -, no; +, 10-30%; ++, 30-70%; +++, 70-100%; nd, not determined.




Figure 5. Inhibition of luciferase reporters and HIV-1 production by antiviral polycistronic miRNAs. (A) Knockdown efficiencies of the antiviral miRNA A^{pol47} expressed from 1-, 2- and 6-hairpin transcripts: A^{pol47}, A^{pol47}B^{wt} and A^{pol47}A^{wt}-E^{wt}. Left panel: 293T cells were co-transfected with 100 ng Luc-A^{pol47}, 10 ng of different hairpin RNA constructs and 1 ng pRL. Two days post-transfection, firefly and renilla luciferase expression was measured. Normalized firefly luciferase expression with pBS was set at 100%. The A^{wt}-E^{wt} construct was used as negative control and sh^{pol47} as positive control. Right panel: 293T cells were co-transfected with 150 ng pLAI, 1 ng pRL and 5 ng of the hairpin constructs. Normalized CA-p24 level in the absence of inhibitor was set at 100%. FL, firefly luciferase; RL, renilla luciferase. (B) Luciferase knockdown efficiencies of 1, 2, 3 and 4 antiviral miRNA transcripts. 293T cells were co-transfected with 100 ng of the indicated luciferase reporter, 10 ng miRNA construct and 1 ng pRL. Two days post-transfection, firefly and renilla luciferase expression was measured. The A^{wt}-E^{wt} construct and a non-matching miRNA were used as negative controls. **CONTINUED ON NEXT PAGE ►**

- The original shRNA constructs sh^{pol47}, sh^{pol1}, sh^{gag5}, sh^{r/t5} and sh^{ldr9} were used as positive controls. (C) Inhibition of HIV-1 production by the miRNA constructs. 293T cells were co-transfected with 150 ng pLAI, 1 ng pRL and 5 ng 1, 2, 3 or 4 miRNA constructs. Two days post-transfection, CA-p24 levels in the culture supernatant and renilla luciferase expression were measured. Normalized CA-p24 level in the absence of inhibitor was set at 100%. Mean values obtained in four independent experiments are shown (\pm SE).

er into 293T cells (Fig. 5B, Supplementary Table 2). Constructs encoding the A^{wt}-E^{wt} cistron and individual wild-type or non-matching miRNAs were used as negative controls. For inhibitors A^{pol47}, C^{gag5} and E^{ldr9}, we observed a remarkable enhancement of silencing activity when chained to other hairpins. For instance, construct C^{gag5} is poorly active compared to construct AC or ACDE (Fig. 5B). Construct B^{pol1} does not exhibit any inhibitory activity, either alone or when combined with another hairpin as in AB. Since the B^{pol1} hairpin structure is rather unstable as predicted by the Mfold algorithm, this could be due to misfolding of the hairpin RNA. We therefore will not use hairpin B^{pol1} in the final polycistron construct. Hairpin D^{r/t5} does not benefit from linkage to other hairpins, but this is likely due to the high inhibitory activity of the individual D^{r/t5} inhibitor. Strong inhibition of Luc-D^{r/t5} was observed with D^{r/t5}, AD, ACD, ACCD and ACDE, indicating that generation of effective siRNAs from the multiplex hairpin transcripts does not depend on the miRNA position in the polycistron.

We further tested the ability of the antiviral miRNAs to inhibit HIV-1 by co-transfection with the HIV-1 molecular clone LAI (Fig. 5C, Supplementary Table 2). Consistent with the luciferase results, we observed a moderate HIV-1 inhibition by the single miRNA constructs, except for construct B^{pol1} (inactive) and D^{r/t5} (highly active) (Fig. 5C). Multimerization of the hairpins strongly enhanced the inhibition of HIV-1 production, which is due both to enhancement of the silencing activity of the individual hairpins and to the presence of multiple antiviral siRNAs. The A^{wt}-E^{wt} construct was used as a negative control and the shRNA constructs were used as positive controls.

Increased siRNA expression from polycistronic miRNA transcripts

We next performed Northern blot analysis of the antiviral siRNAs made by the different polycistron constructs (Fig. 6). The results are remarkably similar to the activity data in Fig. 5. For instance, siRNA expression of A^{pol47} is greatly increased by linkage to another hairpin as in AD (Fig. 6A). The Northern blot analysis of the B^{pol1} and the AB inhibitors provides an explanation for its inactivity as only the ~60-nt B^{pol1} precursor is observed (Fig. 6B, indicated by an arrow). To study whether the passenger siRNA strand of B^{pol1} is made, we performed a Northern blot analysis with the corresponding probe, but failed to detect any passenger siRNAs (results not shown). In contrast, properly processed siRNAs are produced from hairpin A^{pol47} in construct AB, indicating that the inactive B unit in the two-hairpin transcript does not negatively in-

fluence the active A unit (Fig. 6A and Fig. 6B). In addition, siRNA production from the A hairpin is boosted for the polycistronic transcripts compared to the single A^{pol47} hairpin transcript. For inhibitor C^{gag5}, we also observed a strong increase in siRNA production when the hairpin is combined with other hairpins in a polycistronic transcript (Fig. 6C). The D^{r/t5} inhibitor is expressed individually and does not benefit from chaining to other hairpins (Fig. 6D), which correlates nicely with the luciferase results (Fig. 5B). For inhibitor E^{ldr9}, we observed increased siRNA levels for ACDE compared to the single hairpin (Fig. 6E). The combined luciferase inhibition results and the Northern blot analyses demonstrate that the silencing activity of an individual hairpin RNA can be significantly enhanced when expressed in a polycistronic transcript.




Figure 6. Northern blot analyses of siRNAs derived from antiviral miRNA constructs. 293T cells were transfected with the indicated 1, 2, 3, and 4 antiviral miRNA constructs and siRNAs were detected in total cellular RNA using 19-nt complementary LNA oligonucleotide probes. Different miRNA transcripts were analysed: (A) A^{pol47}, (B) B^{pol11}, (C) C^{gag5}, (D) D^{r/t5} and (E) E^{ldr9}. As negative controls, pBS or miRNA constructs targeting another gene were used. The original shRNA constructs were used as positive controls, showing the mature siRNAs. Ethidium bromide staining of 5S rRNA serves as sample loading control. M, RNA size marker (nt) is shown on the left, the probe used, is shown on the right.

Table 2. Toxicity screen of antiviral miRNA constructs

Construct	Percentage of GFP + cells after transduction			
	Day 0	Day 4	Day 20	Day 36
pLV	0	0	0	0
N	2.8	3	1.5	4.5
A ^{pol47}	6	6	6	4.8
ACDE	2.4	3	1.5	2.9

HIV-1 replication is suppressed in cells stably expressing an antiviral miRNA polycistron

We next created stably transduced SupT1 cells with a lentiviral vector (pLV) expressing the individual miRNA constructs A^{pol47}, B^{pol1}, C^{gag5}, D^{r/t5}, E^{ldr9} and the polycistronic ACDE construct. pLV expressing the control miRNA N (Invitrogen) was used as negative control. To study the impact of the antiviral miRNAs on SupT1 cell viability, we set up a sensitive toxicity screen for cells




Figure 7. Inhibition of HIV-1 replication in SupT1 cells transduced with lentiviral vectors expressing antiviral miRNAs A^{pol47}, B^{pol1}, C^{gag5}, D^{r/t5}, E^{ldr9} and ACDE. Untransduced SupT1 cells and cells transduced with the unrelated miRNA N were used as negative controls. Cells expressing an extended triple shRNA construct (e-shRNA 3) serve as positive control. SupT1 cells stably expressing miRNAs A^{pol47}, B^{pol1}, C^{gag5}, D^{r/t5}, E^{ldr9} and ACDE were infected with HIV-1 and virus replication was monitored for 7 days by measuring CA-p24 in the culture supernatant. One representative experiment is shown, similar results were obtained in three independent experiments.

transduced with N, A^{pol47} and ACDE. We cultured the cells for 36 days after lentiviral transduction and followed the percentage of GFP+ (transduced) and GFP- (untransduced) cells by FACS (Table 2). We did not observe a decrease in the fraction of transduced cells, indicating a similar growth rate as untransduced cells.

The stably transduced SupT1 cells were selected by FACS sorting and subsequently infected with HIV-1 (0.5 ng of CA-p24). Virus replication was followed for 7 days by measuring the CA-p24 level in the culture supernatant. Fast virus replication and virus-induced cytopathic effects were observed in cells expressing miRNAs N, B^{pol1}, C^{gag5} and untransduced SupT1 cells (Fig. 7). HIV-1 replication was inhibited by the individual hairpins A^{pol47}, D^{r/t5} and E^{ldr9}. Virus replication was profoundly inhibited in SupT1 cells expressing the polycistronic miRNA construct ACDE and control cells that express an extended triple shRNA construct (e-shRNA 3)(manuscript in preparation).

Discussion

In this study, we designed a combinatorial RNAi approach against HIV-1 using the human mir-17-92 polycistron. We first constructed individual pre-miRNA transcripts against five conserved regions of HIV-1 under the control of a CMV promoter. We maintained the secondary structure of the original pre-miRNAs and included the single stranded flanks because these are important for proper miRNA processing and subsequent RISC loading (12). We used the CMV promoter to express the transcripts because most primary miRNAs are transcribed by RNA polymerase II (7), which also allows inducible or tissue specific miRNA expression (22,53,54). Previously, several reports have demonstrated effective gene knockdown in mammalian cells with siRNAs derived from miRNA precursors (17,18,22,53). However, none of these studies addressed the issue that a mature miRNA is typically 22-24 nt, whereas an siRNA is only 19-nt in length. Here, we demonstrate that positioning of 19-24 nt antiviral siRNA sequences at the 5' end of the pre-miRNA hairpin stem results in optimal HIV-1 inhibition.

Despite the optimization of the miRNA-like inhibitors, their activity is less than that of the original shRNA antivirals that were used to design the miRNAs. We therefore addressed whether the miRNA-like inhibitor is correctly processed and compared the amount of siRNA produced from the miRNA versus shRNA constructs. The siRNA level produced by the shRNA construct sh^{r/t5} is 15-30 fold higher than the D^{r/t5} miRNA, which is likely due to the use of different promoters (RNA polymerase III versus RNA polymerase II). Interestingly, the shRNA inhibitor did only show marginally higher inhibitory activity, suggesting that the intrinsic inhibitory activity of the miRNA-like inhibitor is in fact much greater than the shRNA variant. Consistent with these results, vectors encoding hairpin structures that closely resemble a natural pre-miRNA produced ~12 fold more mature siRNAs than vectors encoding simple

hairpin structures (21). The superior activity of natural miRNA-like inhibitors is likely attributed by the intrinsic properties of a miRNA, which is processed in the nucleus by Drosha, exported to the cytoplasm, processed further by Dicer and loaded into RISC. In the case of an shRNA, the Drosha step is bypassed, which could provide a less efficient entry into the RNAi pathway.

Moderate HIV-1 inhibitory activity was observed with constructs expressing a single miRNA hairpin. Interestingly, we demonstrate that co-expression of two or more hairpins in a single transcript greatly enhanced the silencing activity of each individual hairpin in the transcript. Northern blot analyses showed that the increased inhibitory activity correlates with higher siRNA expression levels. Multimerization of different miRNA hairpins is of particular interest for targeting of RNA viruses such as HIV-1 and HCV because of their extreme genetic diversity and potential for mutational escape. Two recent papers presented the potential of combinatorial RNAi using two miRNA-30 hairpins (55,56). In agreement with these studies, we showed an increase in RNAi activity upon multimerization of 2, 3, 4 or 6 hairpins in a single transcript. Another study used the miR-30 backbone to multiplex artificial miRNAs and reported decreased RNAi activity for the tandem plasmid, which may be due to miRNA processing problems (57). Thus, chaining of miRNAs needs a very careful transcript design. Most studies have used either the miR-30 or miR-155 backbone because these molecules have been studied extensively. Our study focuses on the natural mir-17-92 polycistron and we demonstrate that insertion of 4 antiviral siRNAs creates a transcript that is properly processed into functional antiviral miRNAs that effectively inhibit HIV-1 production and replication.

A major concern of the miRNA approach is the off-target effect on cellular transcripts with partial sequence complementarity. Such an off-target effect may require only a complementarity of 6-8 nt between the seed region of the miRNA and the target (58,59). Such a weak restraint results in numerous potential off-target genes for any miRNA. When multiple miRNAs are used, the number of potential off-targets will increase, increasing the chance of a negative effect on the treated cells. However, we observed no obvious cellular changes in a sensitive toxicity screen. Furthermore, the observed inhibition of firefly luciferase reporters clearly showed sequence-specificity as non-matching miRNAs did not have any effect and the expression of the control renilla reporter was not affected. Another study, in which artificial miRNAs were expressed in *Arabidopsis thaliana*, conferred viral resistance without cellular alterations, suggesting that off-target effects are not significant (60). Furthermore, recently evidence emerges that siRNA sequences inserted in a miRNA backbone do not compete for transport and incorporation into RISC, while competition was observed when the same siRNA sequences were presented as synthetic siRNAs or shRNAs (61). Nonetheless, off-targeting is a genuine concern for the development of any RNAi-based gene therapy against HIV-1 and the potential risk should be assessed properly in relevant *in vivo* models prior to an eventual clinical application (62).

We have shown in stably transduced T cell lines that multiple effective miRNAs inhibit HIV-1 replication much stronger than a single miRNA. These data, together with our previous shRNA studies (32,63), indicate that a combinatorial RNAi approach against HIV-1 results in an increased magnitude of inhibition and consequently a restriction of viral escape. Current strategies combine multiple polymerase III shRNA expression cassettes (63), which results in high expression levels. This may not always be desired in a gene therapy setting because of increased toxicity due to saturation of the RNAi machinery with siRNAs (46). The use of polymerase II promoters to express the miRNA polycistron will reduce this risk because of lower expression levels. In addition, polymerase II cassettes allow expression in a tissue-specific manner and inducible gene expression, which increases the flexibility for gene therapy and functional genomic applications (22,54).

For HIV-1 gene therapy applications the use of a hematopoietic or T cell specific promoter can increase the target cell specificity. An interesting candidate is the WAS promoter that is active in human hematopoietic precursor cells (CD34+) and T lymphocytes, B cells and dendritic cells (64). Another intriguing possibility is to use the HIV-1 LTR promoter to express the miRNA polycistron. Transcriptional activation of the HIV-1 LTR requires the viral Tat protein, which is produced only in HIV-1 infected cells, thereby allowing exquisite target cell specificity. This approach has previously been employed for shRNA expression (65). Thus, several approaches can be tested experimentally in order to optimize the antiviral miRNA polycistron strategy for HIV-1 inhibition.

In summary, one can effectively combat HIV-1 with multiple miRNA effector molecules transcribed from a single polycistronic transcript. We showed that expression of the miRNA polycistron results in the production of functional mature miRNAs that can efficiently and selectively inhibit HIV-1. Further optimization of this construct by increasing the target cell specificity and inducibility will be a further step towards a gene therapeutic approach against HIV-1.

Funding

HIV-1 RNA research in the Berkhoutlab is sponsored by ZonMw (VICI grant) and NWO-CW (TOP grant).

Acknowledgements

We thank Stephan Heynen for performing CA-p24 ELISA and Jens Gruber for useful discussions.

Supplementary Table 1. Oligonucleotides used in this study.


Oligo	Sequence (5'-3')	Target
OncoF1	CATgcccgcg CTGTAAAGAATCTTAAAG	
OncoR1	Cgctccgag GTATCTTGTCAATTAACAGTGG	mir17-19b
GFpseqF	GGCACTGGACGAGCTGTACAAG	
miR8	ACAAACTGGTGTATCTAGAG	1419-1429 pcDNA6.2
CMVf	CGCAATGGCGCTTAGGGCTG	1608-1628 pcDNA6.2
CMVr	ACCGAGGAGGGGTAGGGAT	nL.V
mir17F1	CAGTcgcagga tcc GAGGTGTTAATTC TAATTATC	pri-A ^{mi17}
mir17R1	GTGAGGGGCGAGTAGT ATACATATTTC GTCACTGG	A ^{mi17}
mir17F2	TGGTCTGTAGTATC AGCATTATG GTGACAGCTCCCT	A ^{mi17}
mir17R2	TAGctcgagtgcgcccagat CTCAACATCAACGAGCCCTGGCAGG	pri-A ^{mi17-24}
mir17F3	CAGAATAATG TATTA CTACTGCCCCCTCAGCTTTG CATATGTGGTAAAGTGAATGGT CGTAGTATCAGCATATATGGTG	A ^{mi17-24}
mir17R3	CACCATAATGTGTATAC TACCGACCAATTC ATTAC TTACGCACATATCGAAAGGTGAAGGGCAGTAGT ATACATATTTC GTG	
mir17F3A	CAGAATAATG TATTA CTACTGCCCCCTCAGCTAGT GATATGTGCATCTAGTGAATGGT CGTAGTATCCGCATATATGGTG	A ^{mi17-19}
mir17R3A	CACCATAATGGGATAC TACTACCGACCAATTC ACTAGATGCACATATC ACTAGTGAAGGGCAGTAGT ATACATATTTC GTG	
mir18F1	CATgctgcagga tcc CAAGGTGGAGGGCCGTGTGATGT	pri-B ^{mi18}
mir18R1	ATACAGGAGCAGATATACAGGAAC AAAAGCACTC	B ^{mi18}
mir18F2	AGTAGCC AATCTT ACACTGGCA TAAAGATATGT	B ^{mi18}
mir18R2	GACctcgagtgcgcccagat CTCTGCTGGCTTGAATTATG	pri-B ^{mi18}
mir18F3	TG TTCCGTATCACTGCTCCCTGCTGTA TCGTGGAAGTAGATTAGCATCGAAC AGTACG CAAT GTACACTGGCATAAAGAGTT	B ^{mi18-22}
mir18R3	AACTTCTTATCGCAGGTAA CACTTCTGCTGCTGATG CTATCTACTTCA CGATACAGGAGCAGATACAGGAACA	
mir18F3A	TGTTCCGTATCACTGCTCCCTGCTGTA TGTAAGTAGATTAGCA CTAGCAGTACG CGAT GTACACTGGCATAAAGAGTT	
mir18R3A	AACTTCTTATCGCAGGTAA CACTTCTGCTGCTGATG CTATCTACTTCA CTATACAGGAGCAGATACAGGAACA	
mir19aF1	CATgctgcagga tcc CAAGCCAAAGCATATAGG	pri-C ^{mi19}
mir19aR1	TGCTGTT CATCTCT CTAGTCA GAGGACTGC AAACAA	C ^{mi19}
mir19aF2	ATGCTGTCATC ATTTCTT CTAGT TGGTGGCTCTG ACTTTCC	C ^{mi19}
mir19aR2	GACctcgagtgcgcccagatct CGACACAATAAAGATACA	pri-C ^{mi19}
mir19aF3	AGTCCTCTG ACTAGAGAGATGAACAGCATAC AGAAGAATGTAGC ATGCTGTCACTA TTCTTCT CTAGT TGGTGGCTGC	C ^{mi19-23}
mir19aR3	GCAGGCCACCA CTAGA AGAAATGATGACAGC ATGCTCACTTCTT GTATGCTGTT CACTCTCC TAGTCA GAGGACT	C ^{mi19-23}
mir19aF3A	AGTCCTCTG ACTAGAGAGATGAACAGCATAC AGAAGAATGTAGC ATGCTGTCACTA TTCTTCT CTAGT TGGTGGCTGC	C ^{mi19-23}
mir19aR3A	GCAGGCCACCA CTAGA AGAAATGATGACAGC ATGCTCACTTCTT GTATGCTGTT CACTCTCC TAGTCA GAGGACT	
mir20F1	CTATgctgcagga tcc CAAGTAAAGAGCTGCTGCT	pri-D ^{mi20}
mir20R1	ATGGCAGGAGAG CTCGGAGCTGTCTACAGAACCTGTC	D ^{mi20-23&19}
mir20R1A	ATGGCAGGAGAGCTCGGAGCTGTCTACAGAACCTGTC	D ^{mi20-19}
mir20R1B	AGATGGT CCGAGAGCGTCTG CTACT CAGAAAGCTGTCA TCATCA	D ^{mi20-2&19}
mir20R1C	AGATGGT CCGAGAGCGTCT TAGTGT CTACAGAACCTGTCA TCATCA	D ^{mi20-19}
mir20F2	ATGGCAGGAGAG CTCGGAGCTGTCTACTGCTAGTCTAGTAA	D ^{mi20-2&19}
mir20F2A	ATGCAGGAGAG CTCGGAGTAA AGTACT CTAGTCTAGTAA	D ^{mi20-19}
mir20F2B	ATGGGTGT GAGAGCGAATG TACT CTAGTCTAGTAA	D ^{mi20-2&19}
mir20F2C	TAGTGGTGT GAGAGCGT TAAAGTACT CTAGTCTAGTAA	D ^{mi20-19}
mir20R2	GACctcgagtgcgcccagatct AAGCTGAGCTCTACAGCTA	D ^{mi20-19}
mir20F3	CTCTCTGTAGCACCTCCGCTCTCTCTG CCAT AGGATGTTAGTATTTT CATGGC GGAGGAAGCGG TGGTACT CTAGT	D ^{mi20-23}
mir20R3		
mir20F3A	CTCTCTGTAGCACCTCCGCTCTCTCTG CCAT GTAGT GTTTAGTAT CTAATGGCGGAGGAAGCGG TGGTACT CTAGT	
mir20R3A	CTCTCTGTAGCACCTCCGCTCTCTCCG CCAT TAGATA ACTAA CACTACATGGCAGGAGAAAGCGGAGGTG CTACAGAA	D ^{mi20-19}
mir20F3B	CTCTCTGTAGCACCTCTCTCCGCTCTCTCTG CCAT GTAGT GTTTAGTAT CTAATGGCGGAGGAGGAGTAAAGTACTGCTAGT	
mir20R3B	GCTAGCAGTACTT TACTCCGCT CTCCGCTCATGATA ACTAA CACTACATGGCAGGAGAAAGCGGAGGTAGT GTACTACAGAA	D ^{mi20-19}
mir20F3C	CTCTCTGTAGCACCTCTCTG CCAC CTCTCT GTTTAGTAT TAGAAGATGGGTGC AGAG CAAA GTACT CTAGT	D ^{mi20-23}
mir20R3C	GCTAGCAGTACTT TCGCT CTCACCCATCTT CTATA CTAAACAAGAGAGATGGGTGC AGAG CGCTGCT CTACAGAA	
mir20F3D	CTCTCTGTAGCACCTCTCTG CCAC CTCTCTG TAGTAT CTAGAATGGGTGC AGAG CAAA GTACT CTAGT	
mir20R3D	CTCTCTGTAGCACCTCTCTG CCAC CTCTCT AGATA CTAAACACTACAGATGGGTGC AGAG CGCTGCT CTACAGAA	D ^{mi20-19}
mir20F3E	CTCTCTGTAGCACCTTAGAGCGCTCTCG CA CCCATCTG TGTTAGTAT TAGTGGGTGC AGAG CGCTT AAAGTACT CTAGT	D ^{mi20-19}
mir20R3E	GCTAGCAGTACTT TAAAG CGCTCTCACACCAT TA AACTAAACA CA AGATGGGTGC AGAG CGCTT AGTGTACTACAGAA	
mir19bF1	CATgctgcagga tcc TGGGCTGTGCGCAATCA	pri-E ^{mi19}
mir19bR1	CTCCCTCTCTCTCCATCCAT CCATAGAA CAAGTGT CA GTAA	E ^{mi19-2&19}
mir19bR1A	CTGACCACTTTCTCCAT CCAT TAAGCATAGACAGTGTCTCAGT	E ^{mi19-2&19}
mir19bF2	CTCCGCTCTCTCTG CCAT CTGTGGTAGT	E ^{mi19-2&19}
mir19bF2A	CTCCGCTCTCTCTG CCAT CTGAGTGTGGTAGTGA AAAG GTCTG	E ^{mi19-19}
mir19bR2	TATctcgagtgcgcccagatct GCATTTCCACGCATCCCAAG	pri-E ^{mi19}
mir19bF3	GAC ACTGTTCT TAGTAGTGGTAGAGAGAGGAGAG CTAG CGCTGTG ATAT CTTCGCTGCTCCGCTTCTCTG CCAT CTG TAGT GA	E ^{mi19-23}
mir19bR3	CTACTACACAGATGGCAGGAGAGAGCGGAGCAGCGCAGAATATCACAGCGTAGATCCCTCTCC TACAT CCATAGAACAGTGTTC	E ^{mi19-23}
mir19bF3A	GAC ACTGTTCT TAGTAGTGGTAGAGAGAGGAG CT ATCCAGCTGTG ATAT CTTCGCTGCTCCGCTTCTCTG CCAT CTG TAGT GA	
mir19bR3A	CTACTACACAGATGGCAGGAGAGAGCGGAGCAGCGCAGAATATCACAGCGTAGATCCCTCTCC TACAT CCATAGAACAGTGTTC	E ^{mi19-19}
mir19bF3B	GAC ACTGTTCT TAGTAGTGGTAGAGAGAG CT AGGCTGTG ATAT CTTCGCTCCCTCTCTCTG CCAT CTG TAGT GA	
mir19bR3B	TCACTACACAGCTCAGATGGCAGGAGAGAGCGGAGCGAGAATATCACACAGCTGACCCATTTT CACAT CTAACCATAGAACAGTGTTC	E ^{mi19-19}

* HIV sequences are blue, miRNA sequences are red, pre-miRNA sequences are black upper case, pri-miRNA sequences are black bold, the effective anti-HIV miRNA sequences are bold, restriction sites are lower case, shaded in grey are the nucleotides that were changed to create G:U wobbles and mismatches.

Supplementary Table 2. Properties of the polycistronic antiviral miRNA constructs.

construct name	Inhibition					HIV	siRNA expression				
	Luc-A ^{pol47}	Luc-B ^{pol1}	Luc-C ^{8ags5}	Luc-D ^{r15}	Luc-E ^{ldr9}		pol47	pol1	gag5	r/t5	ldr9
A ^{mi17} B ^{w1}	+++	-	nd	nd	nd	+++	+++	nd	nd	nd	nd
AC	+++	-	+++	-	+++	+++	+++	-	+++	nd	nd
AD	+++	nd	-	+++	-	+++	+++	nd	-	+++	nd
ACC	+++	-	+++	-	+++	+++	+++	nd	+++	nd	nd
ACD	+++	nd	+++	+++	-	+++	+++	nd	+++	+++	nd
ACDE	+++	-	+++	+++	++	+++	+++	nd	+++	+++	+++
ACCD	+++	-	+++	+++	-	+++	+++	-	+++	+++	nd
ACDB ^{w1}	+++	-	+++	+++	-	+++	++	nd	+++	++	nd
A ^{mi17} A ^{w1} -E ^{w1}	+++	nd	nd	nd	nd	++	nd	nd	nd	nd	nd

-, no; +, 10-30%; ++, 30-70 %; +++, 70 -100 %; nd, not determined.



Supplementary figure 1. Construction of the anti-HIV miRNAs. The anti-HIV miRNAs were constructed by a 4-step fusion PCR. The 5' flank of the pri-miRNA was amplified with a forward primer F1 encoding a BamHI site and a reverse primer R1 encoding the HIV-1 sequence (blue) at its 3' end (step 1). Similarly, the 3' flank of the pri-miRNA was amplified with a forward primer F2 containing HIV-1 sequences (blue) and a reverse primer R2 encoding BglII and XhoI sites (step 2). Two complementary oligos F3 and R3, creating the stem-loop structure of the anti-HIV miRNA, were annealed (step 3). The partial sequence similarity between the PCR products from steps 1, 2 and 3 allowed their fusion by PCR with the outer forward F1 and reverse R2 primers (step 4), resulting in an anti-HIV miRNA.

References

1. Fire, A., Xu, S., Montgomery, M.K., Kostas, S.A., Driver, S.E. and Mello, C.C. (1998) Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature*, **391**, 806-811.
2. Meister, G. and Tuschl, T. (2004) Mechanisms of gene silencing by double-stranded RNA. *Nature*, **431**, 343-349.
3. Zamore, P.D., Tuschl, T., Sharp, P.A. and Bartel, D.P. (2000) RNAi: double-stranded RNA directs the ATP-dependent cleavage of mRNA at 21 to 23 nucleotide intervals. *Cell*, **101**, 25-33.
4. Hamilton, A.J. and Baulcombe, D.C. (1999) A species of small antisense RNA in posttranscriptional gene silencing in plants. *Science*, **286**, 950-952.
5. Hammond, S.M., Caudy, A.A. and Hannon, G.J. (2001) Post-transcriptional gene silencing by double-stranded RNA. *Nat.Rev.Genet.*, **2**, 110-119.
6. Lee, Y., Jeon, K., Lee, J.T., Kim, S. and Kim, V.N. (2002) MicroRNA maturation: stepwise processing and subcellular localization. *EMBO J.*, **21**, 4663-4670.
7. Lee, Y., Kim, M., Han, J., Yeom, K.H., Lee, S., Baek, S.H. and Kim, V.N. (2004) MicroRNA genes are transcribed by RNA polymerase II. *EMBO J.*, **23**, 4051-4060.
8. Cai, X., Hagedorn, C.H. and Cullen, B.R. (2004) Human microRNAs are processed from capped, polyadenylated transcripts that can also function as mRNAs. *RNA*, **10**, 1957-1966.
9. Lee, Y., Ahn, C., Han, J., Choi, H., Kim, J., Yim, J., Lee, J., Provost, P., Radmark, O., Kim, S. *et al.* (2003) The nuclear RNase III Drosha initiates microRNA processing. *Nature*, **425**, 415-419.
10. Denli, A.M., Tops, B.B., Plasterk, R.H., Ketting, R.F. and Hannon, G.J. (2004) Processing of primary microRNAs by the microprocessor complex. *Nature*, **432**, 231-235.
11. Han, J., Lee, Y., Yeom, K.H., Kim, Y.K., Jin, H. and Kim, V.N. (2004) The Drosha-DGCR8 complex in primary microRNA processing. *Genes Dev.*, **18**, 3016-3027.

12. Zeng, Y., Yi, R. and Cullen, B.R. (2005) Recognition and cleavage of primary microRNA precursors by the nuclear processing enzyme Drosha. *EMBO J.*, **24**, 138-148.
13. Cullen, B.R. (2004) Transcription and processing of human microRNA precursors. *Mol. Cell*, **16**, 861-865.
14. Kim, V.N. (2005) MicroRNA biogenesis: coordinated cropping and dicing. *Nat. Rev. Mol Cell Biol*, **6**, 376-385.
15. He, L., Thomson, J.M., Hemann, M.T., Hernando-Monge, E., Mu, D., Goodson, S., Powers, S., Cordon-Cardo, C., Lowe, S.W., Hannon, G.J. *et al.* (2005) A microRNA polycistron as a potential human oncogene. *Nature*, **435**, 828-833.
16. Suh, M.R., Lee, Y., Kim, J.Y., Kim, S.K., Moon, S.H., Lee, J.Y., Cha, K.Y., Chung, H.M., Yoon, H.S., Moon, S.Y. *et al.* (2004) Human embryonic stem cells express a unique set of microRNAs. *Dev. Biol*, **270**, 488-498.
17. Boden, D., Pusch, O., Silbermann, R., Lee, F., Tucker, L. and Ramratnam, B. (2004) Enhanced gene silencing of HIV-1 specific siRNA using microRNA designed hairpins. *Nucleic Acids Res*, **32**, 1154-1158.
18. Zeng, Y., Wagner, E.J. and Cullen, B.R. (2002) Both natural and designed micro RNAs can inhibit the expression of cognate mRNAs when expressed in human cells. *Mol. Cell*, **9**, 1327-1333.
19. Zeng, Y., Yi, R. and Cullen, B.R. (2003) MicroRNAs and small interfering RNAs can inhibit mRNA expression by similar mechanisms. *Proc. Natl. Acad. Sci. U.S.A.*, **100**, 9779-9784.
20. Zeng, Y., Cai, X. and Cullen, B.R. (2005) Use of RNA polymerase II to transcribe artificial microRNAs. *Methods Enzymol.*, **392**, 371-380.
21. Silva, J.M., Li, M.Z., Chang, K., Ge, W., Golding, M.C., Rickles, R.J., Siolas, D., Hu, G., Paddison, P.J., Schlabach, M.R. *et al.* (2005) Second-generation shRNA libraries covering the mouse and human genomes. *Nat. Genet.*, **37**, 1281-1288.
22. Stegmeier, F., Hu, G., Rickles, R.J., Hannon, G.J. and Elledge, S.J. (2005) A lentiviral microRNA-based system for single-copy polymerase II-regulated RNA interference in mammalian cells. *Proc Natl Acad Sci U S A*, **102**, 13212-13217.
23. Dickins, R.A., Hemann, M.T., Zilfou, J.T., Simpson, D.R., Ibarra, I., Hannon, G.J. and Lowe, S.W. (2005) Probing tumor phenotypes using stable and regulated synthetic microRNA precursors. *Nat. Genet.*, **37**, 1289-1295.
24. Haasnoot, P.C.J., Cupac, D. and Berkhout, B. (2003) Inhibition of virus replication by RNA interference. *J. Biomed. Sc.*, **10**, 607-616.
25. Haasnoot, P.C.J. and Berkhout, B. (2006) RNA interference: Its use as antiviral therapy. *Handbook of Experimental Pharmacology*. Springer-Verlag Berlin Heidelberg, Heidelberg, Vol. 173, pp. 117-150.
26. Anderson, J., Banerjee, A. and Akkina, R. (2003) Bispecific short hairpin siRNA constructs targeted to CD4, CXCR4, and CCR5 confer HIV-1 resistance. *Oligonucleotides.*, **13**, 303-312.
27. Das, A.T., Brummelkamp, T.R., Westerhout, E.M., Vink, M., Madiredjo, M., Bernards, R. and Berkhout, B. (2004) Human immunodeficiency virus type 1 escapes from RNA interference-mediated inhibition. *J. Virol.*, **78**, 2601-2605.
28. Lo, H.L., Chang, T., Yam, P., Marcovecchio, P.M., Li, S., Zaia, J.A. and Yee, J.K. (2007) Inhibition of HIV-1 replication with designed miRNAs expressed from RNA polymerase II promoters. *Gene Ther.*, **14**, 1503-1512.
29. Boden, D., Pusch, O., Lee, F., Tucker, L. and Ramratnam, B. (2003) Human immunodeficiency virus type 1 escape from RNA interference. *J. Virol.*, **77**, 11531-11535.
30. Westerhout, E.M., Ooms, M., Vink, M., Das, A.T. and Berkhout, B. (2005) HIV-1 can escape from RNA interference by evolving an alternative structure in its RNA genome. *Nucleic Acids Res.*, **33**, 796-804.
31. Berkhout, B. (2004) RNA interference as an antiviral approach: targeting HIV-1. *Curr. Opin. Mol. Ther.*, **6**, 141-145.
32. Ter Brake, O., Konstantinova, P., Ceylan, M. and Berkhout, B. (2006) Silencing of HIV-1 with RNA interference: a multiple shRNA approach. *Mol. Ther.*, **14**, 883-892.

33. Lisziewicz,J., Sun,D., Klotman,M., Agrawal,S., Zamecnik,P. and Gallo,R. (1992) Specific inhibition of human immunodeficiency virus type 1 replication by antisense oligonucleotides: an *in vitro* model for treatment. *Proc Natl Acad Sci U S A*, **89**, 11209-11213.
34. Bertrand,E., Pictet,R. and Grange,T. (1994) Can hammerhead ribozymes be efficient tools to inactivate gene function? *Nucleic Acids Res*, **22**, 293-300.
35. Ohkawa,J., Yuyama,N., Takebe,Y., Nishikawa,S. and Taira,K. (1993) Importance of independence in ribozyme reactions: kinetic behavior of trimmed and of simply connected multiple ribozymes with potential activity against human immunodeficiency virus. *Proc Natl Acad Sci U S A*, **90**, 11302-11306.
36. Ter Brake,O. and Berkhout,B. (2007) Lentiviral vectors that carry anti-HIV shRNAs: problems and solutions. *J Gene Med*, **9**, 743-750.
37. Ter Brake,O. and Berkhout,B. (2005) A novel approach for inhibition of HIV-1 by RNA interference: counteracting viral escape with a second generation of siRNAs. *Journal of RNAi and Gene Silencing*, **1(2)**, 56-65.
38. Liu,Y.P., Haasnoot,J. and Berkhout,B. (2007) Design of extended short hairpin RNAs for HIV-1 inhibition. *Nucleic Acids Res*, **35**, 5683-5693.
39. Konstantinova,P., de Vries,W., Haasnoot,J., Ter Brake,O., de Haan,P. and Berkhout,B. (2006) Inhibition of human immunodeficiency virus type 1 by RNA interference using long-hairpin RNA. *Gene Ther.*, **13**, 1403-1413.
40. Novina,C.D., Murray,M.F., Dykxhoorn,D.M., Beresford,P.J., Riess,J., Lee,S.K., Collman,R.G., Lieberman,J., Shankar,P. and Sharp,P.A. (2002) siRNA-directed inhibition of HIV-1 infection. *Nat.Med.*, **8**, 681-686.
41. Martinez,M.A., Gutierrez,A., Armand-Ugon,M., Blanco,J., Parera,M., Gomez,J., Clotet,B. and Este,J.A. (2002) Suppression of chemokine receptor expression by RNA interference allows for inhibition of HIV-1 replication. *AIDS*, **16**, 2385-2390.
42. Zhou,N., Fang,J., Mukhtar,M., Acheampong,E. and Pomerantz,R.J. (2004) Inhibition of HIV-1 fusion with small interfering RNAs targeting the chemokine coreceptor CXCR4. *Gene Ther.*, **11**, 1703-1712.
43. Jacque,J.M. and Stevenson,M. (2006) The inner-nuclear-envelope protein emerlin regulates HIV-1 infectivity. *Nature*, **441**, 641-645.
44. Nguyen,D.G., Wolff,K.C., Yin,H., Caldwell,J.S. and Kuhen,K.L. (2006) "UnPAKing" human immunodeficiency virus (HIV) replication: using small interfering RNA screening to identify novel cofactors and elucidate the role of group I PAKs in HIV Infection. *J Virol*, **80**, 130-137.
45. Qu,J., Ye,J. and Fang,R. (2007) Artificial microRNA-mediated virus resistance in plants. *J Virol*, **81**, 6690-6699.
46. Grimm,D., Streetz,K.L., Jopling,C.L., Storm,T.A., Pandey,K., Davis,C.R., Marion,P., Salazar,F. and Kay,M.A. (2006) Fatality in mice due to oversaturation of cellular microRNA/short hairpin RNA pathways. *Nature*, **441**, 537-541.
47. Zuker,M. (2003) Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic Acids Res.*, **31**, 3406-3415.
48. Peden,K., Emerman,M. and Montagnier,L. (1991) Changes in growth properties on passage in tissue culture of viruses derived from infectious molecular clones of HIV-1_{LAV}, HIV-1_{MAL} and HIV-1_{ELI}. *Virol.*, **185**, 661-672.
49. Jeeninga,R.E., Hoogenkamp,M., Armand-Ugon,M., de Baar,M., Verhoef,K. and Berkhout,B. (2000) Functional differences between the long terminal repeat transcriptional promoters of HIV-1 subtypes A through G. *J.Virol.*, **74**, 3740-3751.
50. Ruijter,J.M., Thygesen,H.H., Schoneveld,O.J., Das,A.T., Berkhout,B. and Lamers,W.H. (2006) Factor correction as a tool to eliminate between-session variation in replicate experiments: application to molecular biology and retrovirology. *Retrovirology*, **3**, 1-8.
51. Kotsopoulou,E., Kim, V.N., Kingsman,A.J., Kingsman,S.M. and Mitrophanous,K.A. (2000) A Rev-independent human immunodeficiency virus type 1 (HIV-1)-based vector that exploits a

- codon-optimized HIV-1 gag-pol gene. *J.Virol.*, **74**, 4839-4852.
52. Zufferey,R., Dull,T., Mandel,R.J., Bukovsky,A., Quiroz,D., Naldini,L. and Trono,D. (1998) Self-inactivating lentivirus vector for safe and efficient *in vivo* gene delivery. *J Virol*, **72**, 9873-9880.
 53. Chung,K.H., Hart,C.C., Al Bassam,S., Avery,A., Taylor,J., Patel,P.D., Vojtek,A.B. and Turner,D.L. (2006) Polycistronic RNA polymerase II expression vectors for RNA interference based on BIC/miR-155. *Nucleic Acids Res*, **34**, e53.
 54. Shin,K.J., Wall,E.A., Zavzavadjian,J.R., Santat,L.A., Liu,J., Hwang,J.I., Rebres,R., Roach,T., Seaman,W., Simon,M.I. *et al.* (2006) A single lentiviral vector platform for microRNA-based conditional RNA interference and coordinated transgene expression. *Proc Natl Acad Sci U S A*, **103**, 13759-13764.
 55. Sun,D., Melegari,M., Sridhar,S., Rogler,C.E. and Zhu,L. (2006) Multi-miRNA hairpin method that improves gene knockdown efficiency and provides linked multi-gene knockdown. *Biotechn.*, **41**, 59-63.
 56. Xia,X.G., Zhou,H. and Xu,Z. (2006) Multiple shRNAs expressed by an inducible pol II promoter can knock down the expression of multiple target genes. *Biotechn.*, **41**, 64-68.
 57. Zhou,H., Xia,X.G. and Xu,Z. (2005) An RNA polymerase II construct synthesizes short-hairpin RNA with a quantitative indicator and mediates highly efficient RNAi. *Nucleic Acids Res*, **33**, e62.
 58. Lewis,B.P., Burge,C.B. and Bartel,D.P. (2005) Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell*, **120**, 15-20.
 59. Birmingham,A., Anderson,E.M., Reynolds,A., Ilsley-Tyree,D., Leake,D., Fedorov,Y., Baskerville,S., Maksimova,E., Robinson,K., Karpilow,J. *et al.* (2006) 3' UTR seed matches, but not overall identity, are associated with RNAi off-targets. *Nat.Methods*, **3**, 199-204.
 60. Niu,Q.W., Lin,S.S., Reyes,J.L., Chen,K.C., Wu,H.W., Yeh,S.D. and Chua,N.H. (2006) Expression of artificial microRNAs in transgenic *Arabidopsis thaliana* confers virus resistance. *Nat. Biotechnol.*, **24**, 1420-1428.
 61. Castanotto,D., Sakurai,K., Lingeman,R., Li,H., Shively,L., Aagaard,L., Soifer,H., Gatignol,A., Riggs,A. and Rossi,J.J. (2007) Combinatorial delivery of small interfering RNAs reduces RNAi efficacy by selective incorporation into RISC. *Nucleic Acids Res*, **35**, 5154-5164.
 62. Haasnoot,J., Westerhout,E.M. and Berkhout,B. (2007) RNA interference against viruses: strike and counterstrike. *Nat.Biotechnol.*, **25**, 1435-1443.
 63. Ter Brake,O., 't Hooft,K., Liu,Y.P., Centlivre,M., von Eije K.J. and Berkhout,B. (2008) Lentiviral Vector Design for Multiple shRNA Expression and Durable HIV-1 Inhibition. *Mol Ther.*
 64. Charrier,S., Dupre,L., Scaramuzza,S., Jeanson-Leh,L., Blundell,M.P., Danos,O., Cattaneo,F., Aiuti,A., Eckenberg,R., Thrasher,A.J. *et al.* (2007) Lentiviral vectors targeting WASp expression to hematopoietic cells, efficiently transduce and correct cells from WAS patients. *Gene Ther.*, **14**, 415-428.
 65. Unwalla,H.J., Li,M.J., Kim,J.D., Li,H.T., Ehsani,A., Alluin,J. and Rossi,J.J. (2004) Negative feedback inhibition of HIV-1 by TAT-inducible expression of siRNA. *Nat.Biotechnol.*, **22**, 1573-1578.