



**UvA-DARE (Digital Academic Repository)**

**A supersymmetric model for lattice fermions**

Huijse, L.

[Link to publication](#)

*Citation for published version (APA):*

Huijse, L. (2010). A supersymmetric model for lattice fermions

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <http://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.


# Chapter 6

## Ground states of the 2D square lattice on the torus related to rhombus tilings

### 6.1 Introduction

In this chapter we give the proof for theorem 2, which can be formulated both in physical as well as mathematical terminology [35]. We prove the theorem in the mathematics context, namely we find the dimensions of the cohomology for the independence complex on the square lattice wrapped around a torus. In the physics context this translates to the statement that we found the total number of ground states for the supersymmetric model on the square lattice wrapped around a torus. For the interpretation and consequences of the theorem we refer to chapters 5, 7 and 8.

The solution is found by relating ground states, or equivalently elements of the cohomology, to tilings of the plane with two types of rhombi. As was mentioned before, this relation is inspired by the work of Jonsson [37, 39]. He first introduced the rhombi when he related the partition sum of hard squares with activity  $z = -1$  to these rhombus tilings. This is precisely the Witten index for our model on the square lattice, and also the Euler characteristic of  $H_Q$ . The Witten index is a lower bound to the number of ground states. The result presented here gives us, not just this bound, but the total number of ground states with their respective fermion number in terms of rhombus tilings. A rhombus tiling is obtained by tiling the plane with the rhombi depicted in figure 6.1, such that the entire plane is tiled and the rhombi do not overlap (they can have only a corner or a side in common). We call the tiles with area 4 diamonds and the ones with area 5 squares.



**Figure 6.1:** The diamonds on the left and squares on the right.

Although, the proof of the theorem is restricted to a certain class of periodicities on the square lattice, there is strong evidence that the theorem holds in general. This evidence comes from, on the one hand, the fact that Jonsson's result for the Witten index holds for general periodicities and on the other hand, numerical and analytic evidence for small systems (see chapter 7).

## 6.2 Outline of the proof

The proof presented in this chapter is quite involved and consists of several intermediate steps. Here we will give a brief outline of these steps. In section 2.2.3 we discussed the 'tic-tac-toe' lemma which plays a crucial role in the proof. The lemma relates  $H_Q$  to  $H_{Q_1}$  and  $H_{Q_2}$  when  $Q$  is written as  $Q_1 + Q_2$ . This is achieved by writing the lattice  $S$  as  $S_1 \cup S_2$  and letting  $Q_i$  act solely on  $S_i$ . The crucial step is to choose the right sublattices. It turns out that for the square lattice one should pick a set of disconnected points for  $S_1$  and a set of (disconnected) chains for  $S_2$  (for the details see definition 2).

First, in section 6.3, we will discuss the cohomology results for a single chain with various boundary conditions. These results are crucial in the first step of the 'tic-tac-toe' lemma, i.e. computing the cohomology of  $Q_2$ , since  $Q_2$  acts on a set of chains.

Second, in section 6.4, we consider the square lattice on the plane and on the cylinder, to illustrate the power of the 'tic-tac-toe' lemma for a relatively simple case. We choose the boundary conditions in such a way that  $H_{Q_2}$  is non-vanishing only for one value of  $f_1$  and  $f_2$ . Consequently,  $H_{12}$  and  $H_Q$  are trivially obtained from  $H_{Q_2}$ .

Finally, we wrap the square lattice around the torus. We then apply the same strategy as in section 6.4, and  $H_{Q_2}$  is easily obtained. Unfortunately, however, it has entries in several rows and columns of the double complex and computing  $H_{12}$  is highly non trivial.

As a first step (section 6.5.1), we compute  $H_{12}$  for a thin torus, such that the  $S_2$  sublattice consists of one chain only. For this case, we then show that  $H_Q = H_{12}$ , even though  $H_{12}$  has entries in multiple rows. The final step for this simple case is to relate the elements of  $H_Q$  to periodic sequences of tiles and identify the elements that give rise to the small number  $\Delta$  in (5.1).

In the last step (section 6.5.2), we finally present the proof of theorem 2. Here the sublattice  $S_2$  consists of an arbitrary number of chains. We proceed as in section 6.5.1 to obtain  $H_{12}$  and each step will be similar, but slightly more involved. Again we find that  $H_{12}$  does not have entries only in one row. In contrast to the thin torus case, however, we find that here  $H_Q$  is contained in but not equal to  $H_{12}$ . Using the 'tic-tac-toe' procedure, we reduce  $H_{12}$  to obtain  $H_Q$ . What we find is that all the elements of  $H_Q$  can be obtained by concatenating so-called building blocks and in the final step we map each building block to a sequence of tiles. It follows from this mapping, that the elements of  $H_Q$  map to all possible tilings, again with a small discrepancy  $\Delta$ , which is computed in the very last step.

Finally, in section 6.6 we present a counting formula for the number of tilings given the periodicities for which theorem 2 is proven to hold.

## 6.3 The cohomology of $Q$ on the chain

In the following sections we will often use the cohomology results for the supersymmetric model on the chain. These results can be found in [23, 22] (see also section 4.4), but will be restated here for completeness.

**Definition 1** *An open chain of length  $L$  is the graph  $G(V, E)$  with vertices  $V = \{v_j | j \in \mathbb{N}, j \leq L\}$  and edges  $E = \{(v_j, v_{j+1}) | j \in \mathbb{N}, j < L\}$ . A periodic chain of length  $L$  is a cycle defined by the graph  $G(V, E')$  where  $E' = E \cup \{(v_L, v_1)\}$ .*

In the following  $j \in \mathbb{N}$  ( $j$  can also be zero as long as the number of sites  $L$  is positive).

**Theorem 3** *The cohomology of  $Q$  on the periodic chain with  $L$  sites has:*

- 2 non-trivial cohomology classes with  $j$  fermions if  $L = 3j$ ,
- 1 non-trivial cohomology class with  $j$  fermions if  $L = 3j \pm 1$ .

*The cohomology of  $Q$  on the open chain with  $L$  sites has:*

- 1 non-trivial cohomology class with  $j$  fermions if  $L = 3j$  or  $L = 3j - 1$ ,
- 0 non-trivial cohomology classes if  $L = 3j + 1$ .

*Proof.* We prove this result for the periodic chain with  $L = 3j$  sites. We take  $S_2$  to be every third site and the remaining sites  $S_1$ . Remember that  $f_i$  is the number of fermions on sublattice  $S_i$ . Consider a single site on  $S_2$ . If both of the adjacent  $S_1$  sites are empty,  $H_{Q_2}$  is trivial:  $Q_2$  acting on the empty site does not vanish, while the filled site is  $Q_2$  acting on the empty site. So the empty site does not belong to the kernel of  $Q_2$ , whereas the filled site belongs to the image of  $Q_2$ . This leads to a vanishing  $H_{Q_2}$ , unless every site on  $S_2$  is forced to be empty by being adjacent to an occupied site. There are only two such configurations:

$$\begin{aligned} |\alpha\rangle &\equiv \cdots \bullet \square \circ \bullet \square \circ \bullet \square \circ \bullet \square \circ \bullet \square \circ \bullet \square \circ \bullet \square \circ \bullet \square \circ \bullet \square \circ \cdots \\ |\gamma\rangle &\equiv \cdots \circ \square \bullet \circ \square \bullet \circ \square \bullet \circ \square \bullet \circ \square \bullet \circ \square \bullet \circ \square \bullet \circ \square \bullet \circ \square \bullet \circ \square \bullet \circ \square \bullet \circ \cdots \end{aligned} \quad (6.1)$$

where the square represents an empty site on  $S_2$ . Both states  $|\alpha\rangle$  and  $|\gamma\rangle$  belong to  $H_{12}$ : they are closed because  $Q_1|\alpha\rangle = Q_1|\gamma\rangle = 0$ , and not exact because there are no elements of  $H_{Q_2}$  with  $f_1 = f - 1$  fermions, where  $f = L/3 = j$ . By the 'tic-tac-toe' lemma, there must be precisely two different cohomology classes in  $H_Q$ , and therefore exactly two ground states with  $j$  fermions.  $\square$

The proofs for the other cases are completely analogous. In the main proof we will need the representatives of the non-trivial cohomology classes of  $Q$  on the open chain. We will use the following notation: to denote a configuration with fermions on sites  $a, b, c$ , etc. we write  $|a, b, c \dots\rangle$ .

**Lemma 1** *A representative of the non-trivial cohomology classes of  $Q$  on the open chain with  $L = 3j$  or  $L = 3j - 1$  sites is*

$$|\phi\rangle \equiv |2, 5, 8 \dots 3j - 1\rangle, \quad (6.2)$$

*where on the dots the numbers always increase by three.*

*Proof.* From theorem 3 we know that the representative has  $j$  fermions. In the case that  $L = 3j$  it follows that  $|\phi\rangle$  is the only configuration with  $j$  fermions that belongs to the kernel of  $Q$ . Since the dimension of  $H_Q$  is one,  $|\phi\rangle$  must be a representative of the non-trivial cohomology class.

When  $L = 3j - 1$  there are two configurations that belong to the kernel of  $Q$ :  $|1, 4, 7 \dots 3j - 2\rangle$  and  $|2, 5, 8 \dots 3j - 1\rangle$ , however the dimension of  $H_Q$  is again just one. It follows that a linear combination of these two configurations will be in the image

of  $Q$ . In general, two states  $|s_1\rangle$  and  $|s_2\rangle$  are in the same cohomology class if one can write  $|s_1\rangle = |s_2\rangle + Q|s_3\rangle$  for some state  $|s_3\rangle$ . In that case both  $|s_1\rangle$  and  $|s_2\rangle$  are good representatives of the cohomology class. Since  $|\phi\rangle$  itself is not in the image of  $Q$  it is thus a good representative.  $\square$

## 6.4 The cohomology of $Q$ on the square lattice. Part I: Tilted rectangles and cylinders

Let us define  $\mathcal{R}(M, N)$  with  $M, N \geq 1$  as the subset of  $\mathbb{Z}^2$  given by the points  $(x, y)$  such that

$$y \leq x \leq y + M - 1 \quad \text{and} \quad -y + 1 \leq x \leq -y + N. \quad (6.3)$$

This defines a tilted rectangular part of the square lattice. We can also define  $\tilde{\mathcal{R}}(M, N)$  with  $M, N \geq 1$  as the subset of  $\mathbb{Z}^2$  given by the points  $(x, y)$  such that

$$y \leq x \leq y + M - 1 \quad \text{and} \quad -y \leq x \leq -y + N - 1. \quad (6.4)$$

Whereas  $\tilde{\mathcal{R}}(M, N)$  contains the point  $(0, 0)$ , it is excluded in  $\mathcal{R}(M, N)$ . The lattice  $\tilde{\mathcal{R}}(M, N)$  can be mapped to a lattice of the former type except when  $M$  and  $N$  are both odd. Finally, for  $M$  even the cylindrical version  $\mathcal{R}_c(M, N)$  can be obtained from  $\mathcal{R}(M + 1, N)$  by identifying the vertices  $(i, i)$  and  $(i + M/2, i - M/2)$ .

For the lattices  $\mathcal{R}(M, N)$ ,  $\tilde{\mathcal{R}}(M, N)$  and  $\mathcal{R}_c(M, N)$  the full cohomology problem has been solved using Morse theory [75]. These cases can also be solved using the ‘tic-tac-toe’ lemma. The crucial step is to choose the right sublattices. We take a set of disconnected sites for  $S_1$  and a set of (open or periodic) chains for  $S_2$  (see fig. 6.2).

**Definition 2** *More formally, for  $\mathcal{R}(M, N)$   $S_1$  is the set of points  $(x, y)$  that satisfy*


$$y \leq x \leq y + M - 1 \quad \text{and} \quad -y \leq x \leq -y + N - 1 \\ \text{and} \quad x = -y + 3s, \quad (6.5)$$

*with  $3 \leq 3s \leq N - 1$  and  $S_2$  is the set of points  $(x, y)$  that satisfy*

$$y \leq x \leq y + M - 1 \quad \text{and} \quad -y \leq x \leq -y + N - 1 \\ \text{and} \quad -y + 3p + 1 \leq x \leq -y + 3p + 2, \quad (6.6)$$


*with  $0 \leq 3p \leq N - 3$ . The sublattices can be defined similarly for  $\tilde{\mathcal{R}}(M, N)$ .*

To solve  $H_{Q_2}$  we start from the bottom-left chain. If a site on  $S_1$  directly above this chain is occupied, we are left with an isolated site on the bottom-left chain (see fig. 6.3), leading to a vanishing  $H_{Q_2}$  (see section 6.3). It follows that all sites directly above the bottom-left chain must be empty. Continuing this argument for subsequent chains one finds that all sites on  $S_1$  must be empty. However, in the case that  $N = 3l + 1$  we have a set of disconnected sites at the top-right that belong to  $S_2$ . From the previous argument we obtained that the sites of  $S_1$  directly below the top-right sites of  $S_2$  have to be empty. This implies that for  $N = 3l + 1$   $H_{Q_2}$  vanishes. When  $N \neq 3l + 1$  we find that all elements in



**Figure 6.2:** Sublattice  $S_1$  is indicated by circles and sublattice  $S_2$  is indicated by the fat lines. The bounding lines of  $\mathcal{R}(M, N)$  defined in (6.3) are drawn for various values of  $M$  and  $N$ . For the cylinder  $M$  should be even, but for the rectangle it can be odd as well. One easily checks that the length of the  $S_2$  chain is  $M$ . Note that for  $N = 3n + 1$  only half of the upper-right  $S_2$  chain is included in  $\mathcal{R}(M, N)$ .

$H_{Q_2}$  have all sites in  $S_1$  empty, thus computing  $H_{Q_1}(H_{Q_2})$  is a trivial step. The dimension of  $H_Q$  is related to the number of ground states, or equivalently, the number of non-trivial cohomology classes of  $Q$  on the chains that constitute  $S_2$ . Note that the length of these chains is  $M$  both for the tilted rectangles as well as for the cylinder. In the first case the chains have open boundary conditions, whereas in the latter the chains are periodic. Now, the number of non-trivial cohomology classes of  $Q$  for all these cases can be found in theorem 3.



**Figure 6.3:** A site directly above the bottom-left chain is occupied. This generates an isolated site on the bottom-left chain.

It follows that for the tilted rectangles,  $\mathcal{R}(M, N)$  and  $\tilde{\mathcal{R}}(M, N)$ , with  $N \neq 3l + 1$  we have

- i) no non-trivial cohomology classes for  $M = 3p + 1$
- ii) one non-trivial cohomology class for  $M \neq 3p + 1$ .

For the cylinder,  $\mathcal{R}_c(M, N)$ , with  $N \neq 3l + 1$  and  $M$  even we have

- i) one non-trivial cohomology class for  $M = 3p \pm 1$
- ii)  $2^K$  non-trivial cohomology classes for  $M = 3p$ , with  $K$  the nearest integer to  $N/3$ .

For  $N = 3l+1$  the non-trivial cohomology vanishes for both the rectangle and the cylinder.

## 6.5 The cohomology of $Q$ on the square lattice. Part II: The torus

We now define the doubly periodic lattices via two linearly independent vectors  $\vec{u} = (u_1, u_2)$  and  $\vec{v} = (v_1, v_2)$ . We wrap the square lattice around the torus by identifying all points  $(i, j)$  with  $(i + ku_1 + lv_1, j + ku_2 + lv_2)$ . The main result of [35] is the solution of the full cohomology of  $Q$  on the square lattice with doubly periodic boundary conditions defined by  $\vec{u} = (m, -m)$  and  $\vec{v} = (v_1, v_2)$  such that  $v_1 + v_2 = 3p$ . In particular, we obtain a direct relation between elements of  $H_Q$  and tiling configurations. This relation allows us to prove theorem 2, that was first conjectured by P. Fendley and is strongly inspired by the work of Jonsson [37, 39]. It is restated here for convenience.

*For the square lattice with periodicities  $\vec{v} = (v_1, v_2)$ ,  $v_1 + v_2 = 3p$  with  $p$  a positive integer and  $\vec{u} = (m, -m)$ , we find for the cohomology  $H_Q$*

$$N_n = \dim H_Q^{(n)} = t_n + \Delta_n \quad (6.7)$$

where  $N_n$  is the number of zero energy ground states with  $n$  fermions,  $t_n$  is the number of rhombus tilings with  $n$  tiles, and


$$\Delta_n = \begin{cases} \Delta \equiv -(-1)^{(\theta_m+1)p}\theta_d\theta_{d^*} & \text{if } n = [2m/3]p \\ 0 & \text{otherwise,} \end{cases} \quad (6.8)$$

with  $[a]$  the nearest integer to  $a$ . Finally,  $d = \gcd(u_1 - u_2, v_1 - v_2)$ ,  $d^* = \gcd(u_1 + u_2, v_1 + v_2)$  and

$$\theta_d \equiv \begin{cases} 2 & \text{if } d = 3k, \text{ with } k \text{ integer} \\ -1 & \text{otherwise.} \end{cases} \quad (6.9)$$

Computing the cohomology for these tori is far from trivial. First of all, computing  $H_{Q_2}$  does not imply that all sites on  $S_1$  are empty, instead there are many allowed configurations on  $S_1$ . Secondly, because of this, computing  $H_{Q_1}(H_{Q_2})$  becomes much more involved. Finally, we will see that, generally,  $H_Q$  will be contained in  $H_{12}$ , but not equal to  $H_{12}$ .

We will divide the proof into two parts. We start by proving the theorem for a specific torus, defined by  $\vec{u} = (m, -m)$  and  $\vec{v} = (1, 2)$ . This proof will already contain many steps that we use in the proof for the more general case, however, it will be deprived of certain complications. For instance, here we will find  $H_Q = H_{12}$ . As we said, this is not true in general, and in the second part of the proof a substantial part will be concerned with obtaining  $H_Q$  once we have found  $H_{12}$ .



**Figure 6.4:** The square lattice is wrapped around the torus by imposing periodicities  $\vec{v}$  and  $\vec{u}$ . Here  $\vec{v} = (1, 2)$  and  $\vec{u} = (m, -m)$ , consequently  $S_2$  consists of 1 chain. On the right we have drawn the cylinder, where periodicity in the  $\vec{u}$ -direction is still implied.

### 6.5.1 A special case: $S_2$ consisting of 1 chain

In this section we consider the case where  $\vec{v} = (1, 2)$  and  $\vec{u} = (m, -m)$ . It follows that  $S_2$  consists of exactly one periodic chain (see fig. 6.4).

The proof of theorem 2 for this case will consist of 4 steps:

1. We compute  $H_{Q_2}$ .
2. We compute  $H_{12} = H_{Q_1}(H_{Q_2})$  and show that its elements can be constructed from a finite number of building blocks, called motifs. A motif is characterized by a certain configuration on a finite number of subsequent  $S_1$  sites.
3. We show that  $H_Q = H_{12}$ .
4. We relate the elements of  $H_Q$  to tiling configurations by relating each motif to a small series of tiles.

#### Step 1

First we compute  $H_{Q_2}$  and we find the following:

**Lemma 2** *The cohomology of  $Q_2$  consists of all possible configurations on  $S_1$  except for configurations with a multiple of three  $S_1$  sites empty between two occupied  $S_1$  sites.*

*Proof.* The proof is relatively simple. First note that when a site on  $S_1$  is occupied, it blocks four subsequent sites on the  $S_2$  chain (see fig. 6.4). By occupying sites on  $S_1$  the periodic  $S_2$  chain is cut into smaller pieces of chain with open boundary conditions.



Consequently,  $H_{Q_2}$  vanishes when at least one of these smaller pieces has length  $3p + 1$ . This happens when the number of empty sites between two occupied sites on  $S_1$  is a multiple of three. We conclude that all configurations on  $S_1$  are allowed except for configurations with a multiple of three sites empty between two occupied sites.  $\square$

## Step 2

This step in the proof is the most involved. In the next section, where we prove theorem 2 in all generality, we will often refer back to the results obtained in this step. In this step we compute  $H_{Q_1}(H_{Q_2})$ , where  $H_{Q_2}$  was obtained in the previous step. Let us define  $f_1$  and  $f_2$  as the number of fermions on  $S_1$  and  $S_2$  respectively. Furthermore we shall adopt the following notation: an empty site on  $S_1$  is denoted by 0 and an occupied site is denoted by 1. A configuration on  $S_1$  can then be written as a series of 1's and 0's. In the following we shall consider all possible types of configurations on  $S_1$  that belong to  $H_{Q_2}$  and we shall investigate if they also belong to  $H_{12}$ .

If we consider a configuration on  $S_1$ , we note that there have to be at least two adjacent, empty  $S_1$  sites to allow for  $f_2 > 0$ . This is because two adjacent empty sites leave an open chain of two sites unblocked on  $S_2$  and this has an element in  $H_{Q_2}$  with  $f_2 = 1$ . A typical configuration will thus consist of alternating segments, where a segment is a sequence of  $S_1$  sites. The segments are characterized by the number of fermions on the part of the  $S_2$  chain corresponding to the segment, this is either zero or greater than zero. In a segment with  $f_2 > 0$  all  $S_1$  sites are empty and it contains at least two sites. On the other hand, a segment with  $f_2 = 0$  can have empty sites on  $S_1$ , but the empty sites cannot be adjacent. Finally, a segment with  $f_2 = 0$  will always start and end with an occupied  $S_1$  site. We will call this pair of occupied sites a pair of bounding sites. Note that a segment with  $f_2 = 0$  can consist of a single occupied site, in that case the bounding sites fall on top of each other and the pair of bounding sites is just this one site.

**Example 1** Consider the configuration "1101101010000100", there are two segments with  $f_2 = 0$ , formed by the first nine sites and the fourteenth site respectively. There are also two segments with  $f_2 > 0$  constituted by the rest of the sites. Finally, the first and the ninth site form a pair of bounding sites.

First, we consider the segments with  $f_2 > 0$ .

**Lemma 3**  $Q_1$  acting on a segment with  $f_2 > 0$  gives zero within  $H_{Q_2}$ .

*Proof.* Suppose this segment between a pair of bounding sites consists of  $l$  empty  $S_1$  sites. The corresponding  $S_2$  chain then has length  $L = 2l - 2$ . Since  $l = 3k \pm 1$ , we find  $L = 6k$  or  $L = 6k - 4$ . For these chain lengths the elements of the cohomology of  $Q_2$  have  $2k$  and  $2k - 1$  fermions respectively. We now distinguish two cases: a)  $Q_1$  acts on a site at the boundary of the segment, b)  $Q_1$  acts on a site away from the boundary.

a) In this case the length of the  $S_2$  chain in the new configuration is  $L' = L - 2$ . Thus  $L' = 6k - 2$  or  $L' = 6k - 6$ . On the new chain there are still  $2k$  or  $2k - 1$  fermions respectively. However, theorem 3 states that the cohomology for chain length  $6k - 2$  ( $6k - 6$ ) vanishes at all fermion numbers except  $f = 2k - 1$  ( $f = 2k - 2$ ). Thus the new

configuration does not belong to  $H_{Q_2}$  and it follows that this action of  $Q_1$  within  $H_{Q_2}$  gives zero.

b) In this case the action of  $Q_1$  cuts the  $S_2$  chain into two smaller chains of lengths  $L'_1$  and  $L'_2$ . Their total length is  $L'_1 + L'_2 = L - 4$ , since the occupied  $S_1$  site now blocks 4 sites on the  $S_2$  chain. For  $L = 6k$  we have  $L'_1 = 3k_1$  and  $L'_2 = 3k_2 + 2$  or  $L'_1 = 3k_1 + 1$  and  $L'_2 = 3k_2 + 1$ , where in both cases  $k_1 + k_2 = 2k - 2$ . For the latter case  $H_{Q_2}$  vanishes at all grades. In the first case  $H_{Q_2}$  is non-vanishing only for  $f = k_1 + k_2 + 1 = 2k - 1$ . However, the number of fermions on the  $S_2$  chains in the new configuration is  $f = 2k$  and thus it does not belong to  $H_{Q_2}$ . Similarly, one finds that for  $L = 6k - 4$ , the new configuration does not belong to  $H_{Q_2}$ . Again we obtain that this action of  $Q_1$  within  $H_{Q_2}$  gives zero. Finally, if the segment with  $f_2 > 0$  extends over the entire system, we are always in the case considered under b). However, the original chain length can now also be  $L = 6k - 2$  with  $2k - 1$  fermions on it. Under the action of  $Q_1$  we obtain a new chain of length  $L' = 6k - 6$ , which has a non-vanishing cohomology if and only if  $f = 2k - 2$ . So also in this case we find that the action of  $Q_1$  gives zero within  $H_{Q_2}$ .  $\square$

Second, we consider the segments with  $f_2 = 0$ .

**Lemma 4**  $H_{Q_1}(H_{Q_2})$  vanishes when the number of  $S_1$  sites between any pair of bounding sites in a segment with  $f_2 = 0$ , is  $3p + 1$  and it contains one element otherwise.

**Example 2** Consider a configuration with one empty site between a pair of bounding sites: "101", this is not an element of  $H_{Q_1}(H_{Q_2})$ , since  $Q_1$  on this configuration gives "111", which is also in  $H_{Q_2}$ . Now consider two sites between a pair of bounding sites. Then there are two configurations with one empty site: "1011" and "1101" and one configuration with all sites occupied "1111" (remember that the configuration "1001" does not have  $f_2 = 0$ ). It follows that  $Q_1$  acting on ("1101" - "1011") gives 2"1111", whereas  $Q_1$  acting on ("1101" + "1011") gives zero<sup>1</sup>. Consequently, we find that  $H_{Q_1}(H_{Q_2})$  consists of one element: the sum of the configurations with  $f_1 = 3$ .

*Proof.* We can solve  $H_{Q_1}(H_{Q_2})$  for an arbitrary number of sites between a pair of bounding sites, by realizing that this problem can be mapped to the normal chain. For the normal chain no two fermions can be adjacent, whereas here no two empty  $S_1$  sites can be adjacent. So we can map empty  $S_1$  sites to fermions on the chain and occupied  $S_1$  sites to empty sites in the normal chain. Finally,  $Q_1$  is mapped to  $Q^\dagger$  on the normal chain. For the chain  $H_{Q^\dagger}$  (which has the same dimension as  $H_Q$ ) vanishes when the length of the chain is  $3p + 1$  and it contains one element otherwise. So here we have that  $H_{Q_1}(H_{Q_2})$  vanishes when the number of sites between two occupied sites is  $3p + 1$  and it contains one element otherwise.  $\square$

For a segment with  $f_2 = 0$ , let us denote the representative of  $H_{Q_1}(H_{Q_2})$  by the pair of bounding sites with dots in between, for example we denote ("1101" + "1011") by "1 · 1". Even though, this is now a sum of configurations, we will still refer to this simply

<sup>1</sup>Note that the fermionic character of the particles is reflected in the sign here:  $Q_1$  acting on "1011" gives -"1111", whereas  $Q_1$  acting on "1101" gives +"1111". In the first case the particle is created on position 2 and has to hop over the particle at position 1, this gives a minus sign, in the second case the new particle is created at position 3 and thus has to hop over two particles, giving rise to no overall sign change. Also note that the states are not properly normalized, but this is not important for the argument.

as a configuration. It follows that, for a segment with  $f_2 = 0$ , two types of configurations are allowed. The two types can be distinguished by containing  $3s - 1$  dots or  $3s$  dots. Examples of the first type are: "1", "1 · · 1", "1 · · · · · 1", etc. Note that the configuration with  $s = 0$ , and thus with -1 dots between the pair of bounding sites, is "1". Examples of the second type are: "11", "1 · · · 1", "1 · · · · · 1", etc.

Combining lemma's 3 and 4, we find that  $H_{Q_1}(H_{Q_2})$  is spanned by all configurations that can be formed by concatenating the following motifs:

"000"

"1 ·<sub>3s-1</sub> 100"

"1 ·<sub>3s</sub> 100"

"1 ·<sub>3s-1</sub> 10000"

"1 ·<sub>3s</sub> 10000"

where ·<sub>3s</sub> means  $3s$  dots and, as before, "1 ·<sub>3s-1</sub> 1" with  $s = 0$  means "1".

Finally one can also have all zeroes for any length and all dots for any length. Note that if the number of  $S_1$  sites is a multiple of three, that both the configuration with all zeroes and the one with all dots account for two linearly independent elements of  $H_{12}$ . This is because the cohomology of  $Q$  acting the periodic chain with length a multiple of three has dimension two (see theorem 3).

**Example 3** *As an example, suppose we have  $\vec{v} = (1, 2)$  as always and  $\vec{u} = (10, -10)$ . This implies that  $S_1$  consists of 10 sites and with the defined motifs it follows that the following 12 elements belong to  $H_{12}$ : "1100000000", "1100000100", "1100100000", "1100100100", "1100110000", "1100 1 · · 100", "1 · · · · · 100", "1 · · · 100 000", "1 · · 100 100", "1000010000", "0000000000" and ".....". Note that the first nine motifs have periodicity 10 and thus account for ten elements of  $H_{12}$  each, whereas the motif "1000010000" has periodicity 5 and the last two motifs have periodicity 1. For each element one can easily compute the number of fermions and it follows that the first nine motifs have 7 fermions, the motif "1000010000" has 6 fermions and the last two motifs have again 7 fermions. So in total we have 92 elements in  $H_{12}$  with 7 fermions and 5 elements with 6 fermions.*

### Step 3

In the previous step we have obtained  $H_{12}$  for  $\vec{v} = (1, 2)$ . In this step we show that in this case this is equal to the cohomology of  $Q$ . We do this via the 'tic-tac-toe' procedure [33]. That is, we act on a configuration, say  $|\psi\rangle$ , with  $Q$ . The  $Q_2$  part will automatically give zero, but the  $Q_1$  part not necessarily, since we no longer restrict ourselves to the subspace  $H_{Q_2}$ . If it does give zero, we know that the configuration belongs to the kernel of  $Q$ . The configuration will thus belong to  $H_Q$  unless it also belongs to the image of  $Q$ . In that case, another configuration will map to this configuration at the end of the 'tic-tac-toe' procedure. So we continue with the configurations,  $|\psi_0\rangle$ , that do not belong to the kernel of  $Q_1$ . Since the image of  $|\psi_0\rangle$  does not belong to  $H_{Q_2}$  and it does belong to the kernel of  $Q_2$ , it must also belong to the image of  $Q_2$ . So we can write  $Q|\psi_0\rangle = Q_2|\phi\rangle$ , for some configuration  $|\phi\rangle$ . Now let us define a new state  $|\psi_1\rangle \equiv |\psi_0\rangle - |\phi\rangle$ . It then follows that  $Q|\psi_1\rangle = -Q_1|\phi\rangle$ . If this is zero, we have found that the state  $|\psi_1\rangle$  belongs to the kernel of  $Q$ . If it is non-zero we proceed as before: we try to find a configuration  $|\chi\rangle$ , such that  $Q_1|\phi\rangle = Q_2|\chi\rangle$  and define a new state  $|\psi_2\rangle \equiv |\psi_0\rangle - |\phi\rangle + |\chi\rangle$ , etc. This procedure ends,

either when we have found a state  $|\psi_n\rangle$  such that  $Q|\psi_n\rangle = 0$ , or when  $Q|\psi_n\rangle = |\tilde{\psi}\rangle$  with  $|\tilde{\psi}\rangle$  an element of  $H_{Q_1}(H_{Q_2})$ . In the latter case, we say  $|\psi_0\rangle$  maps to  $|\tilde{\psi}\rangle$  at the end of the 'tic-tac-toe' procedure and we conclude that neither  $|\psi_0\rangle$  nor  $|\tilde{\psi}\rangle$  belong to  $H_Q$ .

For the case we consider in this section, we will show that for each element  $|\psi_0\rangle$ , there is an element  $|\psi_n\rangle$  that belongs to the kernel of  $Q$ . So for each element in  $H_{12}$  we can find a corresponding element in  $H_Q$ , thus we obtain  $H_Q = H_{12}$ . In the next section, however, we will see that this is not true for general boundary conditions. We will then find that after several steps in the 'tic-tac-toe' procedure we map certain configurations in  $H_{12}$  to other configurations in  $H_{12}$ . It follows that the first do not belong to the kernel of  $Q$  and the latter belong to the image of  $Q$ . In that case  $H_Q$  is strictly smaller than  $H_{12}$ .


**Lemma 5**  $H_Q = H_{12}$  for  $\vec{v} = (1, 2)$  and  $\vec{u} = (m, -m)$ .

*Proof.* For the segments with  $f_2 = 0$ , we found that  $Q_1$  vanishes if we choose the states represented by the dots such that they are ground states of the normal chain with empty and occupied sites exchanged (see lemma 4). For the segments with  $f_2 > 0$ , we know from lemma 3 that the new configuration always belongs to the image of  $Q_2$ . That is,  $Q_1|\psi_0\rangle = Q_2|\phi\rangle$ , for some configuration  $|\phi\rangle$  if  $Q_1$  acts on a segment with  $f_2 > 0$ . So we can define a new configuration  $|\psi_1\rangle \equiv |\psi_0\rangle - |\phi\rangle$ , such that  $Q|\psi_1\rangle = -Q_1|\phi\rangle$ . Now  $Q_1$  either acts on a different segment with  $f_2 > 0$ , in which case the new configuration again belongs to the image of  $Q_2$ , or it acts on the same segment. In the latter case the new configuration is cancelled by the same configuration in which the two  $S_1$  sites are occupied in the reverse order due to the fermionic character of the particles. It thus follows that the 'tic-tac-toe' procedure always gives zero after as many steps as there are segments with  $f_2 > 0$ .  $\square$

#### Step 4


In this final step we will show that the dimension of  $H_Q$  (and the fermion number of each state) can be computed by counting all tiling configurations (and the number of tiles per configuration) with the four types of tiles depicted in figure 6.1. For the boundary conditions we consider here, the tilings reduce to a single layer sequence of only two types of tiles. Namely the two tiles that respect the boundary condition  $\vec{v} = (1, 2)$ . These tiles have two edges parallel to  $(1, 2)$  and then the diamond has the other two edges parallel to  $(1, -2)$  whereas the square has the other edges parallel to  $(2, -1)$  (see fig. 6.5(a)). Given the sublattices  $S_1$  and  $S_2$  there are three types of vertices: the ones that belong to  $S_1$ , the lower left sites of the  $S_2$ -chain and the upper right sites of the  $S_2$  chain. It follows that the diamond has one of three types of edges along the  $(1, -2)$  direction and a matching type of edge along the  $(1, 2)$  direction, the square can have one of three different types of edges along the  $(2, -1)$  direction and a matching type of edge along the  $(1, 2)$  direction. We conclude that we have 6 types of tiles, depicted in figure 6.5(b).

To establish theorem 2 we map each of the motifs obtained in step 2 to a unique sequence of tiles. The mapping for the four basis motifs, "100", "1100", "10000" and "110000", is shown in figure 6.6. Remember that each motif is modulo the addition of 3 zeroes and modulo the insertion of 3 dots. In terms of tilings, we find that each basis motif can be followed by an arbitrary repetition of the tiling corresponding to the 3 zeroes (see




**Figure 6.5:** Types of tiles we use to tile the square lattice with periodicities  $\vec{v} = (1, 2)$  and  $\vec{u} = (m, -m)$ . 6.5(a) Shows the diamond and square that respect  $\vec{v} = (1, 2)$ . 6.5(b) Shows the three types of diamonds and squares given the sublattices  $S_1$  and  $S_2$ .

fig. 6.7(a)). On the other hand, insertions of multiples of 3 dots correspond to inserting multiples of the tiling shown in figure 6.7(a) at the dotted line along the  $(1, 2)$  direction in the basis motifs. Some examples are shown in figure 6.7(b). Note that here we cannot easily write the motif of dots directly in the tiles (see fig. 6.7(b)), however, the mapping is still unambiguous.



**Figure 6.6:** The four basis motifs and the corresponding sequences of tiles.

Let us determine the number of fermions per motif. First of all, in a segment with  $f_2 > 0$ , the number of fermions is determined by the length of the corresponding  $S_2$  chain. It is easily verified, that for a segment with  $n$  empty  $S_1$  sites the corresponding chain has length  $2n - 2$ . Moreover, from theorem 3, we know that an element in the cohomology of  $Q$  on a chain with length  $L = 2n - 2$  contains  $[(2n - 2)/3]$  fermions, where  $[a]$  is the nearest integer to  $a$ . Similarly, we find that a segment with  $k$  dots contains  $[2k/3]$  fermions. Thus a segment with  $f_2 = 0$ , consisting of  $k$  dots and a the pair of bounding sites, contains  $[2k/3] + 2$  fermions. From these formulae we find for the four basis motifs "100", "1100", "10000" and "110000", that they contain 2, 3, 3 and 4 fermions respectively. Furthermore, an insertion of 3 zeroes, corresponds to increasing  $n$  by 3, and thus increasing the number of fermions,  $[(2n - 2)/3]$ , by 2. Equivalently, inserting 3 dots corresponds increasing  $k$  by 3, and thus again increasing the number of fermions,  $[2k/3]$ , by 2. If we compare this



**Figure 6.7:** On the left we show the sequences of tiles that correspond to the motifs with 3 zeroes (top) and 3 dots (bottom). The addition of 3 zeroes to a basis motif corresponds to attaching the sequence of tiles corresponding to the 3 zeroes to the sequence of tiles corresponding to the basis motif. An insertion of 3 dots in a basis motif corresponds to inserting the sequence of tiles corresponding to the 3 dots at the dotted line along the  $(1, 2)$  direction in the sequence of tiles corresponding to the basis motif. The insertions of 3 dots in each of the four basis motifs and the corresponding tiling are shown on the right as examples. From these examples it is clear that we cannot write the 3 dots directly in the corresponding sequence of tiles. However, the mapping is still unambiguous.

to the number of tiles in the tilings that correspond to these motifs, we find that they exactly agree. Furthermore, the number of sites in a motif is given by three times the number of  $S_1$  sites in a motif, since there are 2  $S_2$  sites for every  $S_1$  site. On the other hand, for the tiles we find that the area of the diamond is 4 and the area of the square is 5. It is now easily verified that the number of fermions per site for the motifs is the same as the number of tiles per area for the corresponding tiling. Thus we find that, not only is the number of elements in the cohomology of  $Q$  directly related to the number of tilings with the two tiles of figure 6.5(a), but also the number of fermions for each element corresponds to the number of tiles in the corresponding tiling.

One can verify that with these sequences of tiles, and the rules for concatenating them, one can obtain every possible tiling. Each tile can be preceded by a certain type of square and diamond and it can be followed by another type of square and diamond. In total this gives four possibilities for the surrounding neighbors. It can be checked that for each tile all four possibilities can be constructed with the given sequences of tiles and the rules for concatenating them.

Finally, the configurations with all zeroes or all dots account for the extra term in (5.1) in theorem 2 repeated here for convenience:

$$\Delta_i \equiv \begin{cases} -(-1)^{(\theta_m+1)p}\theta_d\theta_{d^*} & \text{if } i = [2m/3]p \\ 0 & \text{otherwise.} \end{cases} \quad (6.10)$$

Remember that

$$\theta_d \equiv \begin{cases} 2 & \text{if } d = 3k, \text{ with } k \text{ integer} \\ -1 & \text{otherwise} \end{cases} \quad (6.11)$$

and with  $\vec{v} = (1, 2)$  and  $\vec{u} = (m, -m)$  we have  $p = 1$ ,  $d = \gcd(u_1 - u_2, v_1 - v_2) = \gcd(2m, -1) = 1$  and  $d^* = \gcd(u_1 + u_2, v_1 + v_2) = \gcd(0, 3) = 3$ . It follows that the extra term is  $-2$  for  $m = 3n$  and  $+2$  otherwise.

Let us consider the configuration with all zeroes, which clearly has periodicity 1. If the number of zeroes is a multiple of three, i.e.  $m = 3n$ , the configuration accounts for 2 ground states, otherwise it accounts for 1 ground state. The number of fermions in this configuration is  $i = [2m/3]$ , i.e. the nearest integer to  $2m/3$ . From the mapping (fig. 6.7(a)) it is clear that the configuration corresponds to a tiling with periodicity 3 if  $m = 3p$ . If  $m \neq 3p$ , however, there is no corresponding tiling. Exactly the same holds for the configuration with all dots. It follows that for  $m = 3p$  the tilings overcount the number of ground states by 2 and for  $m \neq 3p$  the tilings fail to count 2 ground states.


Note that the choice of sublattices  $S_1$  and  $S_2$  has increased the number of tilings unrelated by a lattice symmetry by a factor of three (see fig. 6.5). Indeed when computing the number of ground states with the motifs given in step 2 it turns out that one discovers each tiling three times (given that the tiling is not completely uniform, that is all diamonds or all squares).

**Example 4** *Let us consider the case of example 3 again. So we have  $\vec{v} = (1, 2)$  and  $\vec{u} = (10, -10)$ . One possibility is to cover the lattice with 6 squares. This tiling has a unit cell of size 5 and thus this tiling accounts for 5 ground states. The number of tiles is 6 and thus the ground states will have 6 fermions. We can also cover the lattice with 2 squares and 5 diamonds. The 2 squares can be placed between the diamonds in three independent ways. Each of these three tilings has a unit cell of size 30 and consists of 7 tiles, so they account for 90 ground states with 7 fermions.*

*We compare this with the 12 configurations found in example 3. The motif "1000010000" has periodicity 5 and accommodates 6 fermions, so this corresponds to the uniform tiling with all squares. The configurations with all zeroes and all dots account for two ground states with 7 fermions and have no corresponding tiling. Finally, there are 9 configurations with periodicity 10 and 7 fermions, which account for 90 ground states. Using the mapping given in figure 6.6, we find that these configurations can be split into three groups of three, each group corresponding to one of the tilings with 2 squares and 5 diamonds. For example the motifs "1100000100", "1100110000" and "1100 1 · ·100" correspond to the tiling where the two squares are adjacent. They can be distinguished by considering for example the first of the two squares. In each motif it will be of a different type, where the three types are given in figure 6.5(b) (see fig. 6.8).*

### 6.5.2 The general case: $S_2$ consisting of $p$ chains

In the previous section we had  $\vec{v} = (1, 2)$ . In this section we relax this condition to  $\vec{v} = (v_1, v_2)$  with  $v_1 + v_2 = 3p$  with  $p$  a positive integer. It follows that we get  $p$   $S_2$  chains with their accompanying  $S_1$  sites stacked on top of each other. For this situation we will prove theorem 2. The proof consists of 5 steps:



**Figure 6.8:** The square lattice with periodicities  $\vec{v} = (1, 2)$  and  $\vec{u} = (10, -10)$  can be tiled with 2 squares and 5 diamonds. One of these tilings, with the two squares adjacent is shown on the left. The choice of sublattices splits this tiling into three tilings. These three tilings and their corresponding motifs are shown on the right.

1. We compute  $H_{Q_2}$ .
2. We compute  $H_{12} = H_{Q_1}(H_{Q_2})$ .
3. We compute  $H_Q$  starting from  $H_{12}$  via the 'tic-tac-toe' procedure.
4. We relate the elements of  $H_Q$  to tiling configurations by relating each motif to a small series of tiles.
5. We compute  $\Delta_i$ .

### Step 1


As in the previous section we shall start by computing the cohomology of  $Q_2$ . We will define two types of configurations that do not belong to  $H_{Q_2}$  and then find that  $H_{Q_2}$  consists of all configurations except these two types.

**Lemma 6** *A configuration that contains an occupied site  $(k, l)$  on the  $S_1$  lattice, such that the sites  $(k + 1, l + 2)$  and  $(k + 2, l + 1)$  and/or the sites  $(k - 1, l - 2)$  and  $(k - 2, l - 1)$  are empty, does not belong to  $H_{Q_2}$ .*

*Proof.* It is easily verified (see fig. 6.9) that in this configuration the  $S_2$  sublattice contains the isolated site(s)  $(k + 1, l + 1)$  and/or  $(k - 1, l - 1)$ . This site can be either occupied or empty, which leads to a vanishing  $H_{Q_2}$ .  $\square$

Note that in the previous section this situation never occurred because for each occupied site  $(k, l)$ , the sites  $(k + 1, l + 2)$  and  $(k - 1, l - 2)$  were automatically occupied due to the boundary condition set by  $\vec{v} = (1, 2)$ . The second type of configuration that does not belong to  $H_{Q_2}$  follows from a generalization of lemma 2. Remember that occupying  $S_1$  sites causes the  $S_2$  chains to break into smaller open chains. The length of these open






**Figure 6.9:** A configuration is shown where an occupied  $S_1$  site is surrounded by empty sites. This site isolates a site on the  $S_2$  chains directly below and above the site.

chains now depends on the number of empty  $S_1$  sites directly below and above the  $S_2$  chain. For an example see figure 6.10.

**Lemma 7** *If, for a certain configuration, the sum of the number of empty  $S_1$  sites directly below and above an open  $S_2$  chain is  $3s$ , the configuration does not belong to  $H_{Q_2}$ .*

*Proof.* It is easily verified that the open  $S_2$  chain corresponding to the  $3s$  empty  $S_1$  sites has length  $3(s - 1) + 1$ . This leads to a vanishing  $H_{Q_2}$ .  $\square$



**Figure 6.10:** Part of a configuration is shown. The number of empty  $S_1$  sites directly below and above the  $S_2$  chain is 12. The  $S_2$  sublattice thus contains an isolated chain of length 10. Consequently, this configuration does not belong to  $H_{Q_2}$ .

A configuration does not belong to  $H_{Q_2}$  if it contains one or more isolated open chains on the sublattice  $S_2$  with length  $3p + 1$ . It is easy to see that all such configurations fall into the class of configurations described in lemma 6, or lemma 7, or both. It follows that all configurations that do not fall into either of these classes belong to  $H_{Q_2}$ .

## Step 2

As in the previous section, we will now compute  $H_{12} = H_{Q_1}(H_{Q_2})$ .

**Definition 3** *Define a row of  $S_1$  sites as the set of  $S_1$  sites directly above one  $S_2$  chain.*

Note that the configurations in  $H_{Q_2}$  again contain segments where  $f_2$ , the number of fermions on the  $S_2$  sublattice, is zero and segments where it is non-zero.

**Lemma 8** *Lemma 4 for  $H_{12}$  holds for each row of  $S_1$  sites.*

That is, in the segments where  $f_2 = 0$ , the cohomology of  $Q_1$  vanishes when the number of  $S_1$  sites between any pair of bounding sites is  $3p+1$  and it contains one element otherwise. The proof can be found in the previous section. It follows that, in the segments where  $f_2 = 0$ , two types of configurations on a row of  $S_1$  sites are allowed. Using the notation of the previous section, the two types can be distinguished by containing  $3s-1$  dots or  $3s$  dots.

**Lemma 9** *The configurations in  $H_{12}$  have spatially separated columnar segments where  $f_2 = 0$  and segments where  $f_2 > 0$ . The width of a column in a segment where  $f_2 = 0$  can vary between  $3s+1$  and  $3s+2$   $S_1$  sites, whereas the width of a column in a segment where  $f_2 > 0$  can vary between  $3p-1$  and  $3p+1$   $S_1$  sites. In the latter case, two consecutive rows never both have width  $3p$  and the difference in their widths is at most 1 (or -1).*

*Proof.* This follows from combining lemma's 6, 7 and 8. □

An example is shown in figure 6.11. From lemma 9 it follows that we only have to consider columns of width varying between 1 and 2 in the segments where  $f_2 = 0$  separated by columns of width varying between 2 and 4 in the segments where  $f_2 > 0$ . All other configurations can be obtained from these configurations by inserting multiples of 3 dots in the segments where  $f_2 = 0$  over the entire height of the columns, and, similarly, by inserting multiples of 3 zeroes in the segments where  $f_2 > 0$  over the entire height of the columns.


We now turn to the segments where  $f_2 > 0$ . Remember that in the previous section this step was easy because all  $S_1$  sites in the segment where  $f_2 > 0$  were blocked by fermions on the  $S_2$  chain. Here, however, that is not the case. The first thing we note in this case is the following.

**Lemma 10** *The  $S_1$  sites within a column marking a segment where  $f_2 > 0$  have to be empty if they are away from the boundaries with adjacent columns marking a segment where  $f_2 = 0$ .*


*Proof.* This follows directly from lemma 6. □

From this lemma it follows that we only have to consider the  $S_1$  sites on the boundary between a segment where  $f_2 > 0$  and a segment where  $f_2 = 0$ . In fact, we will argue that we only have to consider the boundary where the segment with  $f_2 > 0$  is to the right of a segment with  $f_2 = 0$  (and not the boundary on the other side).

First, however, we introduce a new notation where a configuration is fully characterized by the boundaries between the two types of segments ( $f_2 = 0$  and  $f_2 > 0$ ). From lemma 6 it follows that these boundaries are an arbitrary sequence of steps of  $+(2, 1)$  and  $+(1, 2)$ . However, in the new notation we shall tilt the lattice by  $-45^\circ$ , such that the rows of  $S_1$  sites are horizontal. If we then draw the boundary as a collection of vertical lines between two  $S_1$  sites that are to the left and to the right of the boundary, we find that



**Figure 6.11:** Part of a configuration is shown.




**Figure 6.12:** Part of a configuration is shown with a mapping to the new notation.

the boundaries have a zigzagged shape. The segments where  $f_2 > 0$  will be white and the segments where  $f_2 = 0$  will be grey. For an example see figure 6.12.

Suppose for a moment that we would have a completely disconnected graph, that is, just a collection of disconnected vertices. Then each site can be both empty and occupied. It is clear that each configuration with a least one empty site does not belong to the kernel of  $Q$ , whereas each configuration with at least one occupied site belongs to the image of  $Q$ . It follows that  $H_Q$  vanishes at all grades. Here we do not have a disconnected graph, however, it turns out that the division in grey and white regions is similar to disconnecting the graph.

We define a special notation for a site that can be both empty and occupied. If this site is to the right of a grey region we shall denote this site with a dot, whereas when it is to the left of a grey region the site will be shaded. That is, suppose there are two configurations that both belong to  $H_{Q_2}$  and obey lemma 9, such that these two configurations differ by one site only. Then we can summarize these two configurations in one picture by denoting this particular site by a dot if it is to the right of a grey region or by shading the site if it is to the left of a grey region. For an example see figure 6.13. Moreover, we can summarize  $2^n$  configurations in one picture if the picture contains  $n$  sites with dots or shaded sites. We make a distinction between sites to the left and to the right of the grey region, because we will argue that the configurations with a site with a dot do not belong to  $H_{12}$ . Clearly this is a choice, we could also have chosen to argue that the configurations with a shaded site do not belong to  $H_{12}$ .


Let us consider a boundary that separates a grey segment on the left from a white segment on the right. There are only a few such configurations that may have a site on the boundary with a dot.



**Figure 6.13:** The big brackets indicate how the configurations at the top can be summarized using the notation introduced in the text. The two left-most configurations at the top differ by one site in the *right*-most boundary of the grey region, they can therefore be summarized by the left-most picture at the bottom by denoting this site with a dot. The middle two configurations at the top differ by one site in the *left*-most boundary of the grey region, they can therefore be summarized by the middle picture at the bottom by shading this site. Finally, all four configurations at the top can be summarized by the right-most picture at the bottom.

**Lemma 11** *There are 8 possible configurations with a site with a dot in a boundary that separates a grey segment on the left from a white segment on the right. The configurations are depicted in figure 6.14(c).*

*Proof.* The first restriction follows from lemma 6. That is, both when the site is empty as well as when it is occupied, the configuration should satisfy the lemma. This restriction



**Figure 6.14:** In three steps we find that there are 8 possible configurations with a site with a dot in a boundary that separates a grey segment on the left from a white segment on the right: (a) It follows from lemma 6 that a site with a dot must have occupied sites to the upper left and lower left and empty sites to the upper right and lower right. (b) There are four possibilities for the left-most boundary of the grey segment, following from the two shaded sites being empty or occupied. (c) There are two possibilities for the right-most boundary of the white segment.

is depicted in figure 6.14(a). Then the second restriction follows from lemma 8, that is, the width of a grey column varies between 1 and 2 modulo 3. It follows that next to the site with the dot there can only be one grey site (modulo 3). Combining this again with lemma 6, we find four possibilities for the left boundary of the grey segment. The four possibilities can be summarized in one picture with the notation defined above, see figure 6.14(b), the shaded sites can be both empty and occupied. Finally, it follows from lemma 7 that there are only two possible right most boundaries for the white segment, each modulo columns of width 3, see figure 6.14(c).  $\square$

From lemma 11 it follows that if there is more than one site with a dot in the same boundary, they are sufficiently far away to be independent. That is, each of these sites can be both empty and occupied independent of the configuration of the other dotted sites. Also note that if we select one of the 8 configurations with a dot from figure 6.14(c), the rest of the system can take on any configuration independent of the configuration of the dotted site. Note that this resembles a disconnected graph.

We are now ready to solve  $H_{Q_1}(H_{Q_2})$ .

**Lemma 12** *All configurations that contain a boundary between a grey segment to the left and a white segment to the right, such that this boundary contains one or more sites with a dot, do not belong to  $H_{Q_1}(H_{Q_2})$ .*

*Proof.* A site with a dot can be either empty or occupied. Suppose the site is empty and we act with  $Q_1$  on the configuration. If  $Q_1$  can act non-trivially only on the site under consideration we are done, since the configuration in which the site with the dot is empty does not belong to the kernel of  $Q_1$  and the configuration in which it is occupied belongs to the image of  $Q_1$ . This proves the lemma for this case.

If, however,  $Q_1$  can act non-trivially also on other sites, there are four scenarios: a) The other site is in the same boundary. b) The other site is in the left boundary of the grey region under consideration. c) The other site is in the right boundary of the white region under consideration. d) The other site is further away from the region under consideration than the first three cases.

For scenario a), we know that the other site is also a site with a dot. It follows that the configuration with both dotted sites empty does not belong to the kernel of  $Q_1$ . The sum of the configurations with one of the dotted sites empty belongs to the image of  $Q_1$ . The difference of the configurations with one of the dotted sites empty does not belong to

the kernel of  $Q_1$ , because it maps to the configuration with both dotted sites occupied. Clearly, the latter configuration belongs to the image of  $Q_1$ . So for this scenario the lemma is proven.

For scenario b) we distinguish two cases. First, the other site and the dotted site can be occupied simultaneously. In this case we can prove the lemma via the same argument as we did for scenario a). Second, the other site and the dotted site *cannot* be occupied simultaneously. This only happens when the other site is in the same row as the dotted site. In this case the sum of the configurations with one of them occupied is in the image of  $Q_1$ , but the difference belongs to the kernel of  $Q_1$  and does not belong to the image of  $Q_1$ . The latter is thus an element of  $H_{Q_1}(H_{Q_2})$ . However, we have the freedom to decide to keep only the configuration in which the other site is occupied and the dotted site is empty as a representative of this element. At this point it becomes clear why we only consider configurations with a site with a dot, and not configurations with a shaded site. For scenario c) we can again distinguish two cases. In the first case, the configuration of the other site and the dotted site are independent and the lemma is proven as for scenario a). In the second case, the other site and the dotted site *cannot* be occupied simultaneously. There are again three configurations under consideration. The configuration with both sites empty does not belong to  $\ker Q_1$ , the sum of the configurations with one of the two sites occupied belongs to  $\text{Im } Q_1$  and, finally, the difference again is an element of  $H_{Q_1}(H_{Q_2})$ . And as under b), we choose to represent this element with the configuration where the dotted site is empty and the other site occupied.

Finally, for scenario d) it is clear that the configuration of the other site and the dotted site are always independent and the lemma is proven as for scenario a).

In the four scenarios we considered, there was just one other site on which  $Q_1$  acts non-trivially. If there are more sites on which  $Q_1$  acts non-trivially, the lemma clearly holds when these sites can again be empty or occupied independent of the dotted site. However, if they are not all independent, the proof is more lengthy, but analogous to the proofs of the second case in scenarios b) and c).

□


**Lemma 13** *All the configurations that belong to  $H_{Q_1}(H_{Q_2})$  are a sequence of alternating grey and white columns subject to the conditions in lemma 9, such that the left-most boundary of all the white columns does not contain any sites with a dot.*

*Proof.* This is a direct consequence of lemma 12.

□

As an example we consider the case where  $\vec{v} = (6, 6)$  and  $\vec{u} = (m, -m)$ , that is, we stack four rows of  $m$   $S_1$  sites separated by four  $S_2$  chains. All configurations in  $H_{12}$  can be obtained by concatenating the configurations depicted in figure 6.15, with eventual insertions of grey and/or white columns of width 3, such that the boundary conditions are satisfied<sup>2</sup>. That is, each row should in the end have width  $m$  or, equivalently, the right-most boundary should fit with the left-most boundary. Finally, there is a configuration with all zeroes (one entirely white segment) and a configuration with all dots (one entirely grey segment).

<sup>2</sup>These configurations are obtained as follows. First consider all possible white segments satisfying the boundary condition in the  $\vec{v}$ -direction, then construct all possibilities for the grey segments to the left of these white segments.



**Figure 6.15:** Building blocks of the configurations spanning  $H_{Q_1}(H_{Q_2})$  for  $\vec{v} = (6, 6)$  and  $\vec{u} = (m, -m)$ , modulo insertions of grey and white columns of width 3.

### Step 3

In the previous step we have determined  $H_{12}$ . According to the 'tic-tac-toe' lemma, the cohomology of  $Q$  is equal to or contained in  $H_{12}$ :  $H_Q \subseteq H_{12}$ . In the previous section we found that for  $\vec{v} = (1, 2)$ , we have  $H_Q = H_{12}$ . For general  $\vec{v}$ , however, this is not the case. That is, within  $H_{12}$ , there are configurations that are not in the kernel of  $Q$  and there are configurations that are in the image of  $Q$ . To find out which configurations do not belong to  $H_Q$ , we follow the 'tic-tac-toe' procedure [33] as described in step 3 of section 6.5.1.

In the previous section, we found via the 'tic-tac-toe' procedure that we could find for each element  $|\psi_0\rangle$ , that belongs to  $H_{12}$ , but not to  $\ker Q$ , an element  $|\psi_n\rangle$  that does belong to  $\ker Q$ . In this section, however, we will find that for some elements  $|\psi_0\rangle$ , the 'tic-tac-toe' procedure leads to a corresponding element  $|\tilde{\psi}\rangle$ , that also belongs to  $H_{12}$ . We then say that  $|\psi_0\rangle$  maps to  $|\tilde{\psi}\rangle$  at the end of the 'tic-tac-toe' procedure and we conclude that neither  $|\psi_0\rangle$  nor  $|\tilde{\psi}\rangle$  belong to  $H_Q$ .

We now prove some rules for the 'tic-tac-toe' procedure specific to the configurations we obtained in the previous step.

**Lemma 14** *Let  $Q$  act on an empty  $S_1$  site  $(k, l)$ , such that for the preceding  $S_1$  sites on that row we have:  $(k-1, l+1)$  and  $(k-3s-2, l+3s+2)$  are occupied and the intermediate sites are dotted. Then the new configuration with  $(k, l)$  occupied, is also the image of  $Q_1$  acting on the configuration with  $(k, l)$  occupied and one less fermion in the preceding sites  $(k-1, l+1)$  to  $(k-3s-1, l+3s+1)$ .*

*Proof.* For general  $s$  we can denote the original configuration as " $1 \cdot_{3s} 10$ ", the new configuration is then " $1 \cdot_{3s} 11$ ". From lemma 4 we know that, if the number of  $S_1$  sites between a bounding pair is  $3s+1$ ,  $H_{Q_1}$  vanishes. Consequently, each configuration that is in the kernel of  $Q_1$  is also in the image of  $Q_1$ . Now, since the configuration " $1 \cdot_{3s} 11$ " is in the kernel of  $Q_1$  and the number of  $S_1$  sites between the bounding pair is  $3s+1$ , it must also be in the image of  $Q_1$ . Thus, there is a configuration with one less fermion between the

bounding pair that maps to this configuration under the action of  $Q_1$ .  $\square$

**Example 5** For  $s = 0$  this is easily understood: the original configuration will have "110" on the  $S_1$  sites  $(k-2, l+2)$  through  $(k, l)$ . Acting on this with  $Q$  gives "111", however, this can also be obtained by acting with  $Q$  on "101".

**Lemma 15** Acting with  $Q$  on a white segment away from the boundary, gives zero.

*Proof.* The proof is analogous to the proof of lemma 3. The length of the  $S_2$  chains in the white region is  $L = 3k$  or  $L = 3k - 1$  each containing  $k$  fermions. If  $Q_1$  acts on a site above this chain and away from the boundary, it will cut the  $S_2$  chain into 3 pieces. One of length 1 and two of lengths  $L'_1$  and  $L'_2$ , such that  $L'_1 + L'_2 = L - 3$ . We will now argue that the new configuration with the smaller  $S_2$  chains, always belongs to  $\text{Im } Q_2$ . This implies that we can always continue to the next step in the 'tic-tac-toe' procedure. If the chain of length 1 contains a fermion, the new configuration clearly belongs to  $\text{Im } Q_2$ . If it is empty there are  $k$  fermions on the other two chains. For  $L = 3k$  their combined length is  $L'_1 + L'_2 = 3(k-1)$ , so  $L'_1 = 3k_1$  and  $L'_2 = 3k_2$  or  $L'_1 = 3k_1 + 1$  and  $L'_2 = 3k_2 - 1$ , where in both cases  $k_1 + k_2 = k - 1$ . For the second case the cohomology vanishes for all fermion numbers because of the length  $L'_1$ . For the first case the cohomology is non-vanishing only if  $f = k_1 + k_2 = k - 1$ , however, there are  $k$  fermions. So for both cases the new configuration belongs to  $\text{Im } Q_2$  (since it belongs to  $\ker Q_2$  and not to  $H_{Q_2}$ ). For  $L = 3k - 1$  we find  $L'_1 = 3k_1$  and  $L'_2 = 3k_2 - 1$  or  $L'_1 = 3k_1 + 1$  and  $L'_2 = 3k_2 - 2$ , where in both cases  $k_1 + k_2 = k - 1$ . The rest of the argument is the same as before.

From the above it follows that we can always continue with the next step in the 'tic-tac-toe' procedure. Now suppose that in this next step we act with  $Q_1$  on the same row as in the first step. Since there are no fermions between these two  $S_1$  sites, this configuration will cancel against the configuration where the two  $S_1$  sites are occupied in the reverse order. It follows that we only have to consider acting with  $Q_1$  on each row just once.

It is now easily verified that, since there are as many  $S_2$  chains as there are  $S_1$  rows, we can always continue the 'tic-tac-toe' procedure until we get zero.  $\square$


In this lemma we restricted ourselves to  $Q$  acting on  $S_1$  sites away from the boundary. We will see in the following that if we allow  $Q$  to act on sites at the boundary, the 'tic-tac-toe' procedure can map one configuration in  $H_{12}$  to another configuration in  $H_{12}$ . The crucial point is that, when we act with  $Q_1$  on a site at the boundary, the length of at least one of the  $S_2$  chains below and above this site is reduced by 1. If the original length was  $3k$ , the new length is  $3k - 1$  and both have non-vanishing cohomology for  $f = k$ . In that case we cannot use this  $S_2$  chain to write the new configuration as  $Q_2$  of some other configuration. It follows that to continue the 'tic-tac-toe' procedure, we have to use the other  $S_2$  chain. However, if this chain was already used in a previous step, the 'tic-tac-toe' procedure could end. Before we continue with an example that illustrates this point, we will argue that it is enough to consider  $Q$  acting only on sites at the boundary. This follows from lemma 9; if the 'tic-tac-toe' procedure ends because we have obtained a configuration that does not belong to  $\text{Im } Q_2$  (nor to  $\text{Im } Q_1$ ), this configuration must belong to  $H_{12}$ . From lemma 9 we know that configurations in  $H_{12}$  have spatially separated columnar grey and white segments that do not branch. It follows that we can only map one configuration in  $H_{12}$  to another by either creating a new grey column in a white column, or by (locally)



increasing the width of a grey column. Since the first possibility is excluded by lemma 15, we conclude that we can restrict  $Q$  to act only on sites at the boundary. As in step 2 we will restrict ourselves to the left-most boundary to avoid overcounting.

Let us consider an example of a configuration that does belong to  $H_{12}$ , but does not belong to  $H_Q$ , i.e. it maps to another configuration in  $H_{12}$  at the end of the 'tic-tac-toe' procedure.

**Example 6** Consider the configuration shown on the left in figure 6.16. We label the three  $S_2$  chains (not shown explicitly) between the four  $S_1$  rows; chain 1, chain 2 and chain 3 ( $c_1$ ,  $c_2$  and  $c_3$ ) from top to bottom. Similarly, we label the  $S_1$  rows; row 1 to row 4 ( $r_1$  to  $r_4$ ) from top to bottom. The  $S_2$  chains have lengths  $L_{c_1} = 6$ ,  $L_{c_2} = 5$  and  $L_{c_3} = 3$  and thus contain 2, 2 and 1 particle respectively. Now consider the left-most, empty  $S_1$  sites in the middle two rows. Occupying the left-most, empty  $S_1$  site on row 2 reduces the length of  $c_1$  from 6 to 5. There will still be 2 particles on  $c_1$  and since the chain of length 5 has non-vanishing cohomology at grade 2, the configuration on this chain will in general not belong to  $\text{Im } Q_2$ . Occupying the left-most, empty  $S_1$  site on row 3 reduces the length of  $c_3$  from 3 to 2. Again the configuration on this chain will not belong to  $\text{Im } Q_2$ , since the chain of length 2 has non-vanishing cohomology at grade 1. It follows that if we occupy either of these  $S_1$  sites in the 'tic-tac-toe' procedure, we have to use  $c_2$  to write the new configuration as  $Q_2$  of some other configuration. By definition this is always possible in the first step of the procedure. However, also by definition, we can do this only once since  $Q^2 = 0$ . It follows that, after two steps in the 'tic-tac-toe' procedure, we obtain a new configuration (see fig. 6.16 on the right) that has 2, 1 and 1 particles on the  $S_2$  chains from top to bottom and belongs to  $H_{12}$ . Consequently, both the original as well as the final configuration do not belong to the cohomology of  $Q$ , although they do belong to  $H_{12}$ .




**Figure 6.16:** On the left we depict the configuration that does belong to  $H_{12}$ , but not to  $H_Q$ , since it does not belong to the kernel of  $Q$ . Instead it maps to the configuration depicted on the right under the 'tic-tac-toe' procedure. This configuration also belongs to  $H_{12}$ , but not to  $H_Q$ , since it belongs to  $\text{Im } Q$ . The configuration on the left has 4 particles on sublattice  $S_1$  and 5 particles on sublattice  $S_2$ , divided as 2, 2, 1 over the  $S_2$  chains from top to bottom. The configuration on the right has one more particle in total; it has 6 particles on  $S_1$  and it has 4 particles on  $S_2$ , divided as 2, 1, 1 over the  $S_2$  chains from top to bottom.

As we anticipated, the crucial point in this example is that the length of an  $S_2$  chain is reduced from  $3k$  to  $3k - 1$ , since this limits the options to continue the 'tic-tac-toe' procedure. In fact, in the 'tic-tac-toe' procedure, we can only reach a configuration that is not in the image of  $Q_2$  if the length of an  $S_2$  chain is reduced from  $3k$  to  $3k - 1$ . To find the configurations in  $H_{12}$  that map to another configuration in  $H_{12}$  under the action

of  $Q$  in the most efficient way<sup>3</sup>, we will start the 'tic-tac-toe' procedure by occupying an  $S_1$  site, such that this happens. It follows that we can then only use the other  $S_2$  chain to continue the procedure. We will then, again for efficiency, continue the procedure by again occupying an  $S_1$  site such that there is just one  $S_2$  chain that we can use to continue the procedure. This means that we will act with  $Q_1$  on consecutive rows, either moving upwards or downwards along the boundary.

In the previous step we constructed all possible configurations with a site *with a dot* in the left-most boundary of a white segment. Here we will construct all possible configurations with a site in the left-most boundary of a white segment, such that occupying this site reduces the length of an  $S_2$  chain from  $3k$  to  $3k - 1$ . We shall call such sites 'critical reducer sites'. We start with the white segment and obtain the configurations depicted in figure 6.17. For these configurations occupying the left-most site of the middle row reduces the length of at least one of the adjacent  $S_2$  chains from  $3k$  to  $3k - 1$ . For the two configurations on the left, occupying this site reduces the length of both  $S_2$  chains to  $3k - 1$ . It follows that the new configuration must belong to  $\text{Im } Q_1$  (see lemma 14), otherwise it was a site with a dot in the previous step. So we do not have to consider these two configurations. This same reasoning tells us that the grey region to the left of the middle row should have width 1 modulo 3, otherwise the new configuration would belong to  $\text{Im } Q_1$ . This leads to the 12 possibilities in figure 6.18.




**Figure 6.17:** The possible boundaries of a white segment are shown, such that the left-most boundary contains a critical reducer site. That is, occupying this site reduces the length of at least one of the adjacent  $S_2$  chains from  $3k$  to  $3k - 1$ .


Note that, indeed, occupying the critical reducer site at the boundary, leads to reducing the length of one of the  $S_2$  chains from  $3k$  to  $3k - 1$ , for all these configurations. For efficiency we continue the 'tic-tac-toe' procedure either upwards or downwards, such that at every step in the procedure there is just one  $S_2$  chain that we can use to continue the procedure. The direction we should follow, is indicated by the arrow in figure 6.19. Note that we dropped the two configurations for which the grey segment had width 2, because of lemma 14.

It is now clear that if we stack a configuration for which the 'tic-tac-toe' procedure goes downwards on top of a configuration for which it goes upwards, the 'tic-tac-toe' procedure ends. In particular, it maps the old configuration to a new configuration that is also in  $H_{12}$ . We can increase the number of steps necessary in the 'tic-tac-toe' procedure by stacking

<sup>3</sup>By 'the most efficient way' we mean the shortest sequence of occupying  $S_1$  sites in the 'tic-tac-toe' procedure that maps one configuration in  $H_{12}$  to another. This is the most efficient way, because as soon as this happens, we know that both configurations do not belong to  $H_Q$ , independent of all the other terms created under the action of  $Q$ .




**Figure 6.18:** The 12 possible configurations such that the left-most site of the middle row is a critical reducer and occupying this site does not lead to a configuration that is in  $\text{Im } Q_1$ .




**Figure 6.19:** If the critical reducer site (the left-most site of the middle row) is occupied in the first step of the 'tic-tac-toe' procedure, the procedure should be continued in one direction only, as explained in the text. This direction is indicated by the arrow.

rows for which the grey segment has width 1 modulo 3 and the width of the white segment alternates between 3 and 4 modulo 3 (see fig. 6.20). Examples of the stacked configurations and the configurations they map to are shown in figure 6.21. Here the sites with connected dots can be either all empty or all occupied. The configuration with all the sites empty maps to the configuration with all the sites occupied under the 'tic-tac-toe' procedure. However, if the configurations on the left in figure 6.19 are not combined with one of the configurations on the right in figure 6.19, the 'tic-tac-toe' procedure will end with a state that is in the kernel of  $Q$  (as long as we only let the sites on the left-most boundary participate).



**Figure 6.20:** Stacking these configurations with the configurations of figure 6.19, increases the number of steps in the 'tic-tac-toe' procedure.

At this point we have identified a certain set of configurations that does belong to  $H_{12}$ , but



**Figure 6.21:** Some examples are shown of configurations that do belong to  $H_{12}$ , but not belong to  $H_Q$ . Here the sites with connected dots can be either all empty or all occupied.

does not belong to  $H_Q$ . However, we have to make a final step before we can identify all configurations in  $H_Q$  with tiling configurations. This is due to the fact that certain parts of configurations seem to belong to  $H_Q$ , but they do not respect the boundary conditions. Note that, at this point, we have reduced all possible motifs to the following set:

"100"

"1100"

"10000"

"110000"

which can be separated by single insertions of the motifs:

"1000"

"11000"

all modulo insertions of three dots and three zeroes along an entire column. Each of the four basis motifs, comes with two directions, determined by whether the boundaries between the grey and white segments follows the direction  $(-1, -2)$  or  $(-2, -1)$ .

**Definition 4** We assign a letter to each of the four basis motifs:

$A_i \equiv "100"$

$B_i \equiv "1100"$

$C_i \equiv "10000"$

$D_i \equiv "110000"$


where the subscript  $i$  is 1 or 2 when the direction of the motif is  $(-1, -2)$  or  $(-2, -1)$  respectively.

Note that the direction of a motif is not defined if neither the motif directly above it nor the motif directly below it is the same. We will start, however, by considering cases in which this does not happen. At the end of step 4 we will encounter a case where this point needs some attention.

We now want to study whether a vertical sequence of a certain motif can be followed by a sequence of a different motif, eventually, with an insertion of one of the motifs with 3 zeroes: "1000" or "11000".

**Example 7** As an example let us start with a sequence of motif  $A_1$ . This sequence could be followed by motifs  $A_2$ ,  $B_1$  and  $C_2$ . However, it cannot be followed by motif  $B_2$ , because

it would not belong to  $H_{12}$ . Nor can it be followed by motifs  $C_1$  or  $D_i$ , because it would not belong to  $H_Q$  (see fig. 6.22).



**Figure 6.22:** At the top, we show, from left to right, motif  $A_1$  followed by the motifs  $A_2$ ,  $B_1$  and  $C_2$ . On the bottom-left, we see that a configuration in which  $A_1$  is followed by  $B_2$  contains a site with a dot in the left-most boundary of the white segment. The other two configurations on the bottom, show that configurations in which  $A_1$  followed by  $C_1$  or  $D_i$  do not belong to  $H_Q$ . Here we used the notation of figure 6.21.

Similarly we find the following:

- motif  $B_1$  can only be followed by motif  $C_2$ .
- motif  $C_2$  can only be followed by motif  $B_1$ .
- motif  $A_i$  can be followed by motifs  $A_j$ ,  $B_1$  and  $C_2$ .
- motif  $D_i$  can be followed by motifs  $D_j$ ,  $B_1$  and  $C_2$ .
- motif  $B_2$  can only follow after motif  $C_1$ .
- motif  $C_1$  can only follow after motif  $B_2$ .


Finally, we know from lemma's 7 and 8 that grey and white columns cannot branch or have end points, since their width always oscillates between 1 and 2 modulo 3 or 2, 3 and 4 modulo 3 for the grey and white segments respectively. Consequently, grey and white columns may wind around the torus several times, but they will always close to form a loop.

Let us combine this observation with the rules we found for stacking motifs. Consider, for example, motif  $A_1$ , which can be followed by motifs  $A_2$ ,  $B_1$  and  $C_2$ . However, motifs  $B_1$  and  $C_2$  can only be followed by  $C_2$  and  $B_1$  respectively. Consequently, if motif  $A_1$  is followed by either of these two motifs, we can never fulfill the boundary conditions, because the column cannot be closed to form a loop. Thus configurations in which motif  $A_1$  is followed by motifs  $B_1$  and  $C_2$  do not belong to  $H_Q$ . In this same spirit we obtain the following lemma.

**Lemma 16** *For configurations that belong to  $H_Q$  the following holds:*

- motif  $B_1$  can only be followed by motif  $C_2$  and vice versa.
- motif  $A_1$  can only be followed by motif  $A_2$  and vice versa.
- motif  $D_1$  can only be followed by motif  $D_2$  and vice versa.
- motif  $B_2$  can only be followed by motif  $C_1$  and vice versa.


For motifs  $A_i$  and  $D_i$  the width of the white segment does not change, thus the motif with direction 1 can follow directly below or above this same motif with direction 2. For the motifs  $B_i$  and  $C_i$ , however, there is an intermediate motif of the type "1000" or "11000". Which of the two can be determined via the 'tic-tac-toe' procedure. If we read the motifs of the rows from top to bottom, we find that a sequence of  $B_1$  motifs will be followed by "1000", to be followed by a sequence of  $C_2$  motifs. Then the  $C_2$  motifs will be followed by "11000", which is then to be followed by another sequence of  $B_1$  motifs. On the other hand, a sequence of  $B_2$  motifs will be followed by "11000", followed directly by a sequence of  $C_1$  motifs. Finally, the  $C_1$  motifs will be followed by "1000", followed directly by another sequence of  $B_2$  motifs (see fig. 6.23). It is readily checked that any other choice gives a configuration that does not belong to  $H_{12}$ , but not to  $H_Q$ .




**Figure 6.23:** Two configurations are shown in which motif  $B_i$  is followed by motif  $C_j$  (where  $i \neq j$ ) and vice versa, with the correct intermissions of the motifs "1000" and "11000" (indicated by the arrows).

#### Step 4

We are now ready to make the identification with the tiles. For the four basis motifs  $A_1$  through  $D_1$  the identification is shown in figure 6.24 and  $A_2$  through  $D_2$  are identified with a tiling in figure 6.25. Note that to distinguish motif  $X_1$  from  $X_2$ , where  $X = A, B, C$  or  $D$ , one has to consider also the motif on the row above or below this motif. These motifs can be followed by an arbitrary threefold of zeroes. Let us define the motif  $E \equiv "000"$ . For this motif we can also distinguish a direction, because its boundary will follow the left-most boundary of the white segment it is attached to. From figures 6.24 and 6.25 it is clear that the motifs  $X_i$  can be followed by motif  $E_i$ , where the  $i$  should be the same. For



**Figure 6.24:** The motifs  $X_1$  and the corresponding tilings. Note that this mapping was already found in section 6.5.1 (see fig. 6.6).



**Figure 6.25:** The motifs  $X_2$  and the corresponding tilings.




**Figure 6.26:** On the left the motifs  $E_1$  (on the left) and  $E_2$  (on the right) and the corresponding tilings. On the right the tilings corresponding to insertions of 3 dots into the motifs  $X_1$  (on the left) and  $X_2$  (on the right).

the motifs  $B_1$  and  $C_2$  there is an exception: when the motif above these motifs is "11000" and "1000" respectively, they are followed by  $E_2$  and  $E_1$  respectively.


There can also be insertions of multiples of three dots in the four basis motifs. How this translates into tilings is shown in figure 6.27. More precisely, note that each basis motif  $X_1$  contains a dotted line, connecting two  $S_1$  sites along the direction 1 (black) and equivalently, all basis motifs  $X_2$  contain a dotted line with direction 2 (blue). An insertion of three dots in a basis motif corresponds to an insertion of two tiles, shown in figure 6.26, at this dotted line. Note that here we cannot easily write the motif of dots directly in the tiles, however, the mapping is still unambiguous.

Finally, we have the motifs "1000" and "11000". Which tilings these motifs correspond to depends on whether they occur between the motifs  $B_1$  and  $C_2$  or the motifs  $B_2$  and  $C_1$ . In fact, in the first case, "1000" will correspond to the same sequence of tiles as  $B_1$

and "11000" to the same tiling as  $C_2$ . Similarly, in the latter case, "1000" and  $B_2$ , on the one hand, and "11000" and  $C_1$ , on the other hand, correspond to the same tilings. For an example see figure 6.28.



**Figure 6.27:** On the top (bottom), insertions of 3 dots into the motifs  $X_1$  ( $X_2$ ) and the corresponding tilings are shown.



**Figure 6.28:** On the left (right), we show a configuration in which motifs  $C_2$  and  $B_1$  ( $C_1$  and  $B_2$ ) alternate with the corresponding tilings. Note that the motifs "1000" and "11000" correspond to different tilings on the left than on the right.

With these identifications there is one ambiguity, but it is easily dealt with. If there is a column in which the motifs alternate indefinitely between "1100" and "11000", we cannot determine whether the motif "1100" is of type  $B_1$  or  $B_2$ . The same happens when the motifs "10000" and "1000" alternate indefinitely: the motif "10000" could be of type  $C_1$  or  $C_2$ . Note that if we choose to identify the first with  $B_1$  and the second with  $C_2$ , the



corresponding tilings would be indistinguishable. This also happens when we choose  $B_2$  and  $C_1$ . So we conclude that we should either choose  $B_1$  and  $C_1$  or  $B_2$  and  $C_2$ . The ambiguity is thus lifted by simply deciding that we will always choose, say,  $B_1$  and  $C_1$ . Finally, we note that again the number of fermions in a certain configuration is the same as the number of tiles in the corresponding tiling. For the four basis motifs and the motifs with 3 zeroes or 3 dots, this follows from the arguments in section 6.5.1. For the motifs "1000" and "11000", we should look at figure 6.28. If the motif "1000" sits between motifs  $B_1$  and  $C_2$ , the number of fermions in these three rows is  $3 * 3$  and the number of sites is 3 times the number of  $S_1$  sites:  $3 * (2 * 4 + 5)$ . So 9 fermions on 39 sites. Compare this with the corresponding tiling: it contains 2 times 3 tiles of area 4 and once 3 tiles of area 5. So 9 tiles with total area 39. Similarly, if the motif "11000" sits between motifs  $B_1$  and  $C_2$ , the number of fermions in these three rows is  $3 * 3$  and the number of sites is  $3 * (2 * 5 + 4)$ . The corresponding tiling contains 2 times 3 tiles of area 5 and once 3 tiles of area 4. For the corners between motifs  $B_2$  and  $C_1$  the comparison is slightly more subtle. Following the same arguments as above, we find that in this case the number of fermions in the motifs "1000" and "11000" do not agree with the number tiles in the corresponding tiling. However, the discrepancy is minus one in one case and plus one in the other, and since the boundary conditions dictate that the number of "1000"-motifs equals the number of "11000"-motifs, the discrepancies exactly cancel.

## Step 5

The final step concerns the small correction  $\Delta$  in equation (5.1). With the four basis motifs, horizontal insertions of multiples of three dots and three zeroes and vertical insertions of the motifs "1000" and "11000", we can represent all elements in  $H_Q$ . With the mappings given in the previous step, we find a corresponding tiling for each of these elements. On the other hand, each possible tiling can be constructed with the small sequences of tiles given in the previous step. Thus we find that for each possible tiling there is a corresponding element in  $H_Q$ . Furthermore, we found that the number of fermions and the number of tiles agree. So we find  $N_i = t_i$ , that is, the number of elements in  $H_Q$  with  $i$  fermions equals the number of tilings of the square lattice with  $i$  tiles. However, there is a small discrepancy in this one-to-one relation for the configurations with all zeroes or all dots. For  $\vec{u} = (m, -m)$  and  $v_1 + v_2 = 3p$ , it is readily verified that these configurations contain  $i = [2m/3]p$  fermions. In the following we will first compute the number of elements of  $H_Q$  that these configurations account for. We shall call this  $N^{(a)}$ , where  $a$  stands for anomalous. We will then compute  $t^{(a)}$ , the number of tilings consisting only of the tiles that correspond to either all zeroes or all dots (see fig. 6.26). Combining these results we obtain  $\Delta \equiv N^{(a)} - t^{(a)}$ . Finally, since we found a one-to-one correspondence between tilings and elements of  $H_Q$  for  $i \neq [2m/3]p$ , theorem 2 will then be established with  $\Delta_i$  as in equation (5.2).

As we discussed in section 6.5.1 for  $\vec{v} = (1, 2)$ , the configurations with all dots and all zeroes actually correspond to multiple elements of the cohomology if there is a multiple of 3  $S_1$  sites per row, that is if  $\vec{u} = (3n, -3n)$ . This is a direct consequence of theorem 3, which says that a periodic chain with length  $3j$  has two ground states. In fact, these configurations account for  $2^p$  elements each, where  $p = (v_1 + v_2)/3$  is the total number of  $S_1$  rows or, equivalently, of  $S_2$  chains. On the other hand, for  $\vec{u} = (m, -m)$  with  $m \neq 3n$

they each represent one element of the cohomology. So we find  $N^{(a)} = 2^{p+1}$  for  $m = 3n$  and  $N^{(a)} = 2$  otherwise.

Now, let us look at the corresponding tilings. For  $\vec{u} = (m, -m)$  with  $m \neq 3n$  there is no corresponding tiling, thus there is a discrepancy of 2 between the number of tilings and the number of elements in the cohomology. That is  $\Delta \equiv N^{(a)} - t^{(a)} = 2$  for  $m \neq 3n$ . For  $\vec{u} = (3n, -3n)$  there are tilings corresponding to the configurations with all zeroes or all dots. Along the  $\vec{u}$  direction these tilings have periodicity 3. The periodicity in the other direction is more involved. Given the boundary condition  $\vec{v} = (2r + s, r + 2s)$  the tiling makes  $r$  steps in the  $(2, 1)$  direction and  $s$  steps in the  $(1, 2)$  direction, in arbitrary order. However, because of the periodicity of 3 in the  $\vec{u}$  direction one can also end at  $(2r + s + 3l, r + 2s - 3l)$ , that is,  $r + 3l$  steps in the  $(2, 1)$  direction and  $s - 3l$  steps in the  $(1, 2)$  direction, again in arbitrary order. Thus we find

$$t^{(a)} = 2 * 3 \sum_{l \geq -r/3} \binom{r+s}{r+3l}.$$

If we define  $r = 3k + c$ , where  $c \in 0, 1, 2$ , we can write  $t$  as:

$$\begin{aligned} t^{(a)} &= 6 \sum_{l=0} \binom{r+s}{c+3l} \\ &= 6 \sum_{l=0} \left[ \binom{r+s-2}{c+3l-2} + 2 \binom{r+s-2}{c+3l-1} + \binom{r+s-2}{c+3l} \right] \\ &= 6 \sum_{l=0} \left[ \binom{r+s-2}{l} + \binom{r+s-2}{c+3l-1} \right] \\ &= 6 * 2^{r+s-2} + 6 \sum_{l=0} \left[ \binom{r+s-2}{c+3l-1} \right]. \end{aligned}$$

Repeating these steps  $d$  times, such that  $2d \leq r + s$ , we find

$$\begin{aligned} t^{(a)} &= 6 \sum_{l=1}^d 2^{r+s-2l} + 6 \sum_{l=0} \binom{r+s-2d}{c+3l-d} \\ &= \sum_{l=0}^{2d-1} 2^{r+s-l} + 6 \sum_{l=0} \binom{r+s-2d}{c+3l-d} \\ &= \begin{cases} 2^{r+s+1} - 2 + 6 \sum_{l=0} \binom{0}{c+3l-d} & \text{if } r+s = 2d \\ 2^{r+s+1} - 4 + 6 \sum_{l=0} \binom{1}{c+3l-d} & \text{if } r+s = 2d+1. \end{cases} \end{aligned}$$

For the last term we find

$$\begin{aligned} 6 \sum_{l=0} \binom{0}{c+3l-d} &= \begin{cases} 6 & \text{if } d = 3b + c \\ 0 & \text{otherwise.} \end{cases} \\ 6 \sum_{l=0} \binom{1}{c+3l-d} &= \begin{cases} 0 & \text{if } d = 3b + c + 1 \\ 6 & \text{otherwise.} \end{cases} \end{aligned}$$

We now compare the expression for  $t^{(a)}$  with the expression for the number of elements in the cohomology represented by the configurations with all zeroes and all dots,  $N^{(a)}$ . For

$\vec{v} = (2r + s, r + 2s)$  this is  $N^{(a)} = 2 * 2^{(v_1+v_2)/3} = 2^{r+s+1}$ . So we finally find

$$\Delta = N^{(a)} - t^{(a)} = \begin{cases} -4 & \text{if } r + s = 2d \text{ and } r - s = 6b \\ 2 & \text{if } r + s = 2d \text{ and } r - s = 6b \pm 2 \\ 4 & \text{if } r + s = 2d + 1 \text{ and } r - s = 6b + 3 \\ -2 & \text{if } r + s = 2d + 1 \text{ and } r - s = 6b \pm 1. \end{cases}$$

Combining this with the result  $\Delta = 2$  for  $\vec{u} = (m, -m)$  with  $m \neq 3n$ , this can be cast in the compact form of equation (5.2).

## 6.6 Counting formula for tilings

In the previous sections we proved a theorem that expresses the number of ground states in terms of tiling configurations. In this section we derive a counting formula for the number of tilings of the plane with periodicities given by  $\vec{u} = (m, -m)$  and  $\vec{v} = (2\alpha + \beta, \alpha + 2\beta)$ . An important property of the periodic tilings is that a given tiling can be specified completely by giving a collection of corners of tiles in the tiling that form a closed loop along a non-contractible direction of the torus when connected by edges of length  $\sqrt{5}$ . There is one constraint, namely that in the collection of edges between the corners at least one should be along the directions  $(2, 1)$  or  $(1, 2)$  and at least one should be along the directions  $(2, -1)$  or  $(1, -2)$ .

Let us consider an example. Suppose we have the tiling shown in figure 6.29 with periodicities  $\vec{u} = (6, 0)$  and  $\vec{v} = (0, 6)$ . It is not difficult to see that the set of corners  $\{(0, 0), (1, 2), (3, 1), (5, 2)\}$  completely determines the tiling. However, the set  $\{(0, 0), (1, 2), (3, 3), (4, 5)\}$  does not fully determine the tiling. Indeed in this case none of the edges between the corners is along the directions  $(2, -1)$  or  $(1, -2)$ .

This property is the main reason why the number of tilings and thus the number of ground states of the doubly periodic square lattice does not grow exponentially with the 2D volume, but at most with the linear dimensions of the system.

In the following we will use this property to obtain a counting formula for the number of tilings of the plane with periodicities given by  $\vec{u} = (m, -m)$  and  $\vec{v} = (2\alpha + \beta, \alpha + 2\beta)$ . We will first consider the closed loop along the non-contractible direction of the torus given by  $\vec{u} = (m, -m)$ . Notice that a tiling with periodicity  $\vec{u}$  will only exist if we can write

$$(m, -m) = r_1(1, -2) + s_1(2, -1) + r_2(2, 1) + s_2(1, 2),$$


where the  $r_i$  and  $s_i$  are integers. For  $m$  positive we will have  $r_1, s_1 \geq 0$  and  $r_2, s_2$  will have the same sign. If we solve these equations for  $r_2$  and  $s_2$  we find

$$\begin{cases} r_2 = m - (4r_1 + 5s_1)/3 \\ s_2 = -m + (5r_1 + 4s_1)/3. \end{cases}$$


From this it follows that there are two regions in the parameter space of  $r_1, s_1$  where  $r_2, s_2$  have the same sign. These regions are bound by the lines

$$\begin{cases} r_2 = 0 & \rightarrow s_1 = (3m - 4r_1)/5 \\ s_2 = 0 & \rightarrow s_1 = (3m - 5r_1)/4. \end{cases}$$

The regions are shown in figure 6.30.



**Figure 6.29:** We show a tiling which obeys the periodicities  $\vec{u} = (6, 0)$  and  $\vec{v} = (0, 6)$ .



**Figure 6.30:** The lines  $s_1 = (3m - 4r_1)/5$  and  $s_1 = (3m - 5r_1)/4$  are shown and the region where  $r_2, s_2$  have the same sign are indicated.

Now suppose we have  $r_2 = \lambda$  and  $s_2 = \mu$ . Let us determine the constraints on  $\lambda$  and  $\mu$  set by  $m$ . First of all, we have

$$\begin{cases} r_1 = (m + 4\lambda + 5\mu)/3 \\ s_1 = (m - 5\lambda - 4\mu)/3 \end{cases}$$

and since  $m$  is positive we have  $r_1, s_1 \geq 0$  from which it follows that

$$\begin{cases} 5\lambda + 4\mu \leq m & \text{for } \lambda, \mu \geq 0 \\ 4|\lambda| + 5|\mu| \leq m & \text{for } \lambda, \mu \leq 0. \end{cases}$$

Furthermore, we find that, since  $r_1$  and  $s_1$  are integers, we must have

$$\begin{cases} m + 4\lambda + 5\mu = 3q_1 \\ m - 5\lambda - 4\mu = 3q_2, \end{cases}$$

with  $q_1$  and  $q_2$  positive integers. From this we find, with  $q \in \mathbb{Z}$ ,

$$\lambda - \mu = \begin{cases} 3q & \text{for } m = 3p \\ 3q - 1 & \text{for } m = 3p + 1 \\ 3q + 1 & \text{for } m = 3p + 2. \end{cases}$$

Together with the condition that  $\lambda$  and  $\mu$  have the same sign, these are the conditions imposed by the periodicity  $\vec{u}$ . We now also impose the periodicity in the other direction,  $\vec{v} = (2\alpha + \beta, \alpha + 2\beta)$ . We only need consider the case where  $\alpha$  and  $\beta$  are both positive, since the other cases can be reduced to this case either by reflecting in  $\vec{u}$  or translating by  $\vec{u}$  or both. It is clear that the periodicity  $\vec{v}$  imposes the following condition: either  $\lambda = \mu = 0$  or

$$\lambda = \alpha/k \quad \text{and} \quad \mu = \beta/k,$$

with  $k \in \mathbb{Z}$ .

With all these condition in place, we can finally write down a counting formula. As we have emphasized at the start a tiling can be fully specified by a collection of corners that form a closed loop around an incontractible direction of the torus. Now suppose that we know that this loop contains  $\kappa$  edges in the  $(1, -2)$ -direction,  $\nu$  edges in the  $(2, -1)$ -direction,  $\lambda$  edges in the  $(2, 1)$ -direction and  $\mu$  edges in the  $(1, 2)$ -direction. Since the order of the edges in the directions  $(2, 1)$  and  $(1, 2)$  can be chosen independently from the order of the edges in the directions  $(2, -1)$  and  $(1, -2)$ , it follows that the number of tilings specified by this collection of edges is

$$\frac{5(\kappa\lambda + \mu\nu) + 4(\kappa\mu + \lambda\nu)}{(\kappa + \nu)(\lambda + \mu)} \binom{\kappa + \nu}{\nu} \binom{\lambda + \mu}{\mu},$$

where the first factor is the total number of sites divided by the total number of tiles in the tiling.

Putting everything together we find that the number of tilings  $R_{\vec{u}, \vec{v}}$  for  $\vec{u} = (m, -m)$  and  $\vec{v} = (2\alpha + \beta, \alpha + 2\beta)$  with  $m$  positive and  $\alpha, \beta \in \mathbb{Z}_{\geq 0}$  is given by

$$R_{\vec{u}, \vec{v}} = A_0 + A_k, \tag{6.12}$$

where

$$A_0 = \begin{cases} \frac{9}{2} \binom{\alpha+\beta}{\beta} \binom{2m/3}{m/3} & \text{for } m = 3p \\ 0 & \text{otherwise,} \end{cases} \tag{6.13}$$

and

$$A_k = \sum_k' \frac{5(\kappa\alpha + \nu\beta) + 4(\kappa\beta + \nu\alpha)}{(\kappa + \nu)(\alpha + \beta)} \binom{(\alpha + \beta)/|k|}{\beta/|k|} \binom{\kappa + \nu}{\nu},$$

with

$$\kappa = \frac{1}{3}\left(m + \frac{4\alpha}{k} + \frac{5\beta}{k}\right) \quad \text{and} \quad \nu = \frac{1}{3}\left(m - \frac{5\alpha}{k} - \frac{4\beta}{k}\right).$$

The prime denotes that the sum is restricted to run over integer values of  $k$  such that

$$\begin{cases} (5\alpha + 4\beta)/m \leq k & \text{if } k > 0 \\ (4\alpha + 5\beta)/m \leq |k| & \text{if } k < 0, \end{cases}$$

furthermore we impose  $\binom{a}{b} = 0$  if  $a$  and/or  $b$  is non-integer. Finally, we should have

$$\begin{cases} \frac{\alpha-\beta}{k} = 3q & \text{for } m = 3p \\ \frac{\alpha-\beta}{k} = 3q - 1 & \text{for } m = 3p + 1 \\ \frac{\alpha-\beta}{k} = 3q + 1 & \text{for } m = 3p + 2 \end{cases}$$

with  $q \in \mathbb{Z}$ .

Note that the counting formula derived here is consistent with the result for the growth behavior of the number of tilings (5.4) obtained in [38]. It is easily verified that for  $m \bmod 3 = 0$  and  $m$  and  $\alpha = \beta$  large  $A_0$  is the leading term in (6.12). Finally, writing  $\alpha = \beta = \lambda q$  and  $m = 3\mu q$  with  $q$  large and using Stirling's approximation, one quickly recovers (5.4) from (6.13).