



UvA-DARE (Digital Academic Repository)

Computability of Diagrammatic Theories for Normative Positions

Pascucci, M.; Sileno, G.

DOI

[10.3233/FAIA210333](https://doi.org/10.3233/FAIA210333)

Publication date

2021

Document Version

Final published version

Published in

Legal Knowledge and Information Systems

License

CC BY-NC

[Link to publication](#)

Citation for published version (APA):

Pascucci, M., & Sileno, G. (2021). Computability of Diagrammatic Theories for Normative Positions. In E. Schweighofer (Ed.), *Legal Knowledge and Information Systems: JURIX 2021: The 34th Annual Conference, Vilnius, Lithuania, 8-10 December 2021* (pp. 171-180). (Frontiers in Artificial Intelligence and Applications; Vol. 346). IOS Press. <https://doi.org/10.3233/FAIA210333>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Computability of Diagrammatic Theories for Normative Positions

Matteo PASCUCCI^a, Giovanni SILENO^{b,1}

^a *Slovak Academy of Sciences, Bratislava, Slovakia*

^b *University of Amsterdam, Amsterdam, The Netherlands*

Abstract. Normative positions are sometimes illustrated in diagrams, in particular in didactic contexts. Traditional examples are the Aristotelian polygons of opposition for deontic modalities (squares, triangles, hexagons, etc.), and the Hohfeldian squares for obligative and potestative concepts. Relying on previous work, we show that Hohfeld's framework can be used as a basis for developing several Aristotelian polygons and more complex diagrams. Then, we illustrate how logical theories of increasing strength can be built based on these diagrams, and how those theories enable us to determine in a computably efficient way whether a set of normative positions can be derived from another set of normative positions.

Keywords. Computable Normative Theories and Diagrams and Hohfeldian relationships and Normative Positions and Polygons of Opposition

Introduction

Diagrams are acknowledged to be effective instruments for didactic purposes: they are commonly used to illustrate in an intuitive and accessible way the most various conceptual models, procedures, systems. Examples abound in engineering and theoretical sciences, but archetypes of diagrams include also the *Aristotelian square of opposition* [1], used in philosophical, linguistic, literary, and semiotic studies; and, within legal studies, the *two squares of fundamental normative relationships* due to Hohfeld [2,3]. Reference to regular shapes may be not by chance; cognitive studies show that symmetries facilitate perception of structure, memorization and recall [4]. Indeed, the motivating intuition behind this research is that diagrams may be useful to create user-friendly interfaces for the analysis of legal/contractual constructs. Rather than inspecting hundreds of sentences in the text of a contract, a subject may more easily figure out her normative relations (duties, rights, etc.) with the other parties by navigating or exploring a diagram-construed model. One may also ask whether these diagrams have similarly interesting computational properties; if this is the case, unveiling diagrammatic theories may benefit on multiple levels. For this reason, the present work focuses on the computational treatment of *Aristotelian diagrams* to represent normative positions, building upon previous interpretations of Hohfeld's squares relating these two types of diagrams [5,6,7,8,9].

¹Corresponding Author: g.sileno@uva.nl. Matteo Pascucci was supported by Štefan Schwarz Fund (project "A fine-grained analysis of Hohfeldian concepts") and VEGA Grant No. 2/0117/19; he thanks Jonas Raab for discussion. Giovanni Sileno was supported by NWO for the DL4LD project (628.009.001) and the HUMAINER AI project (KIVI.2019.006). This article is the result of a joint research work of the two authors.




Figure 1. The two Hohfeldian squares: (left) the deontic square, (also obligative or of the first-order), and (right) the potestative square (or of the second-order).

The paper proceeds as follows. Section 1 introduces the notion of a normative position, the two Hohfeldian squares, and the notion of an Aristotelian diagram. Section 2 provides a formal language to encode normative positions and, reorganizing and extending previous results in [8], analyses the representation of Hohfeld’s deontic square and of three versions of Hohfeld’s potestative square (change-centered, force-centered and outcome-centered) in terms of Aristotelian diagrams. Section 3 shows how to develop logical theories over a diagram relying on the syntactic notion of an *inference tree*. These will be called *diagrammatic theories*. The analysis of a complex diagram which combines the three potestative squares of opposition is used as an example. Section 4 focuses on the computational dimension of diagrams, many aspects of which have recently gained attention in the literature (see, e.g., [10]). In particular, it shows that diagrammatic theories enable one to compute in polynomial time whether a finite set of normative positions can be derived from another.

1. Normative positions on diagrams

A *normative position* is described by a statement involving a normative concept, one or more normative parties related to that concept and a certain type of behaviour of one of those parties. For instance, consider the following:

- it is obligatory for the company to pay an annual fee;
- borrowers have a duty towards lenders to bring back the relevant goods;
- by accepting a sale offer, a buyer creates a duty upon the seller to deliver.

Two families of normative positions have been extensively analysed by Hohfeld [2,3]. He proposed to graphically visualize their relations via two diagrams, a *deontic* and a *potestative* square, which are reproduced in Fig. 1.

Many formal accounts of Hohfeld’s analysis have been proposed (see, e.g., Lindahl [11], Makinson [12], and, more recently, Markovich [13] or Sileno and Pascucci [14]). Furthermore, some works [5,6,7] have shown that relationships on Hohfeldian squares can be used to construct Aristotelian polygons of opposition. The fundamental advantage of the latter over Hohfeld’s squares is that they *unambiguously express logical relations* of a certain kind between normative statements. Aristotelian polygons can be combined to form more complex figures, whence we will generally speak of *Aristotelian diagrams*. At the basis of an Aristotelian diagram are four types of logical relations [10].




Figure 2. Aristotelian square for basic deontic modalities, with subaltern (green), contrary (orange), sub-contrary (blue) and contradictory (red) bindings, and named vertices (A, E, I, O). Sub-alternity is *directed*: from A to I, and from E to O, i.e., from orange towards blue lines.

Definition 1 (Aristotelian Relation). *An Aristotelian relation between two sentences ϕ and ψ is either subalternation, contrariety, sub-contrariety or contradiction, where:*

- ϕ is a subalternate of ψ iff the truth of ψ implies the truth of ϕ ;
- ϕ and ψ are contrary iff at most one between ϕ and ψ can be true;
- ϕ and ψ are sub-contrary iff at most one between ϕ and ψ can be false;
- ϕ and ψ are contradictory iff ϕ is true precisely when ψ is false.

Definition 2 (Aristotelian Diagram). *Given a finite set of sentences Γ , an Aristotelian diagram over Γ is a graph whose vertices are labelled by elements of Γ . Each vertex v is labelled by a distinct sentence. An edge e of the graph connecting two vertices v_1 and v_2 is associated with an Aristotelian relation R_e between the sentences labelling v_1 and v_2 .*

In the normative domain, the simplest example of Aristotelian diagram is the *square of opposition for basic deontic modalities* (Fig. 2).

2. Formalization

To enable a more rigorous analysis of normative positions, we here use a language \mathcal{L} of first-order logic to encode sentences, following [11]. Moreover, refining and extending recent work [8], we show how to construct squares of oppositions starting from Hohfeld’s squares and, subsequently, more complex Aristotelian diagrams.

2.1. Language

Language \mathcal{L} has variables x, y etc. for normative parties and α, β , etc. for action types. It has constants p, q , etc. for normative parties and constants A, B , etc. for action types. Symbol $\overline{}$ (overline) denotes action complementation: \overline{A} is the complement of A (the type of any action not instantiating A). We assume that $\overline{\overline{A}} = A$. Hohfeldian (and other) relations are encoded as n -ary predicates.² Finally, \mathcal{L} has standard propositional connectives ($\neg, \wedge, \vee, \rightarrow, \equiv$) and quantifiers (\forall, \exists). We omit quantification over variables for normative parties, interpreting an expression $\phi(x, y, \dots)$ as implicitly having the form $\forall x \forall y \dots \phi(x, y, \dots)$. Thus, while $\text{Claim}(x, y, A)$ means “for all x , for all y : x has a claim that A be performed by y ”, $\text{Claim}(p, q, A)$ means “ p has a claim that A be performed by q ”.

²Sometimes the argument of a relation is a statement involving another relation. However, no quantification on such statements is employed; therefore, \mathcal{L} remains a first-order language.

2.2. First-order Hohfeldian relations

The formal renderings of the fundamental deontic relations identified in Hohfeld's framework, for two normative parties p and q and an action type A , are the following: $\text{Claim}(p, q, A)$, $\text{Liberty}(p, q, A)$, $\text{Duty}(p, q, A)$ and $\text{NoClaim}(p, q, A)$. We can map all relationships to a single primitive, e.g. Claim :

$$\begin{aligned}\text{NoClaim}(x, y, A) &\equiv \neg\text{Claim}(x, y, A) \\ \text{Duty}(y, x, A) &\equiv \text{Claim}(x, y, A) \\ \text{Liberty}(y, x, A) &\equiv \neg\text{Claim}(x, y, \bar{A})\end{aligned}$$

This choice leads to the following set of labels DR with respect to a given action type A :

$$\text{DR} = \{\text{Claim}(p, q, A), \text{Claim}(p, q, \bar{A}), \neg\text{Claim}(p, q, \bar{A}), \neg\text{Claim}(p, q, A)\}$$

The set DR naturally gives rise to a deontic square of opposition. The only additional principle needed is the following, used to characterize *subalternate statements*: $\text{Claim}(x, y, A) \rightarrow \neg\text{Claim}(x, y, \bar{A})$, which can be seen as corresponding to the *Obligatory*(ϕ) \rightarrow *Permitted*(ϕ) axiom used in deontic logics.

2.3. Second-order Hohfeldian relations

Potestative relations concern *actions* that trigger changes of first-order or even second-order relations, such as, for instance, an action B creating a duty for a party q with respect to a party p to perform an action A . A possible way of writing that p has such a power would be by means of a predicate expression *Ability*(p, B, R) (cf. the predicate *has_ability* investigated in [14]), where R is a Hohfeldian relation issued at B 's performance by p ; for instance, *Ability*($p, B, \text{Claim}(p, q, A)$). Different characterizations of actions exist in human language, mapping to different levels of abstraction [15], e.g. the *behavioural* or *procedural* characterization, relating to the action task, or the *productive* characterization, relating to its outcome. In the following, we similarly provide different definitions of power constructed at different abstraction levels (force, outcome, change).

2.3.1. Force-centered power

At behavioural level, power relations can be seen in analogy to force fields determining attraction, repulsion, and absence of those (independence) at the occurrence of interventions ([7], [16, Ch.4]). To express this, we need to separate the *stimulus* component (a particular type of action, such as a verbal command) and the consequent target *manifestation* (e.g. a type of action that is due or expected on the basis of the stimulus, cf. the concept of *pliance*). If the action-manifestation is denoted by the action type symbol A , then, the action-stimulus can be conveniently represented via the symbol "A" to emphasize the shared connection between signal and expected performance.

If stimulus and manifestation converge, i.e. A is always performed in correspondence to its stimulus, we have a *positive force-centered power*:

$$\overset{\rightarrow}{\leftarrow} \text{Power}(x, y, A) \equiv \text{Ability}(x, \text{"A"}, \text{Claim}(x, y, A))$$

If stimulus and manifestation diverge, i.e. A is never performed in correspondence to its stimulus, we have a *negative force-centered power*:³

$$\overset{\leftarrow\rightarrow}{\text{Power}}(x, y, A) \equiv \text{Ability}(x, "A", \text{Claim}(x, y, \bar{A}))$$

From these concepts we can define a new set of potestative relations $\text{PR}^{\leftarrow\rightarrow}$ as labels for a force-centered potestative square of opposition, obtained by taking into account all possible combinations of positive- vs. negative-force power and Boolean negation:

$$\text{PR}^{\leftarrow\rightarrow} = \{\overset{\rightarrow\leftarrow}{\text{Power}}(p, q, A), \overset{\leftarrow\rightarrow}{\text{Power}}(p, q, A), \overset{\leftarrow\rightarrow}{\neg\text{Power}}(p, q, A), \overset{\rightarrow\leftarrow}{\neg\text{Power}}(p, q, A)\}$$

The subalternity is here captured by the logical principle: $\overset{\rightarrow\leftarrow}{\text{Power}}(x, y, A) \rightarrow \overset{\leftarrow\rightarrow}{\neg\text{Power}}(x, y, A)$ which is acceptable because otherwise the same stimulus "A" would generate two conflicting first-order relations.

2.3.2. Outcome-centered power

At the outcome or productive abstraction level, we may abstract the triggering action B , and focus only the output, e.g. $R = \text{Claim}(p, q, A)$. *Positive outcome-centered power*, expressed in the form $\text{Power}(p, q, A)$, means that p has the power of issuing a duty upon q to A . It can be defined via an existential quantification on the set of action types:

$$\text{Power}(x, y, A) \equiv \exists\beta : \text{Ability}(x, \beta, \text{Claim}(x, y, A))$$

We can similarly define a *negative outcome-centered power* (to release a duty to A) as:

$$\overline{\text{Power}}(x, y, A) \equiv \exists\beta : \text{Ability}(x, \beta, \neg\text{Claim}(x, y, A))$$

As before, we can form a set of potestative relations PR as labels for an outcome-centered potestative square of opposition:

$$\text{PR} = \{\text{Power}(p, q, A), \overline{\text{Power}}(p, q, A), \neg\overline{\text{Power}}(p, q, A), \neg\text{Power}(p, q, A)\}$$

where subalternity is captured by: $\text{Power}(p, q, A) \rightarrow \neg\overline{\text{Power}}(p, q, A)$. This principle can be explained as such: to create a duty, this duty needs not to be holding: $\text{Power}(x, y, A) \rightarrow \neg\text{Claim}(x, y, A)$. Dually, to release a duty, this needs to exist: $\overline{\text{Power}}(x, y, A) \rightarrow \text{Claim}(x, y, A)$; its contrapositive provides the subalternity relation.

2.3.3. Change-centered power

Given a target relation R , e.g. a due performance $\text{Claim}(p, q, A)$, one can define power also as the ability of p to *affect* q in any sense with respect to this relation R . This proposal, originally made by O'Reilly [6], can be reframed in our framework as the ability of changing q 's position in the (e.g. deontic) square of opposition of which R is part. Using the proposed notation we have:

$$\begin{aligned} \text{Power}_{\text{O'Reilly}}(x, y, B, A) \equiv & \text{Ability}(x, B, \text{Claim}(x, y, A)) \vee \text{Ability}(x, B, \text{Claim}(x, y, \bar{A})) \\ & \vee \text{Ability}(x, B, \neg\text{Claim}(x, y, A)) \vee \text{Ability}(x, B, \neg\text{Claim}(x, y, \bar{A})) \end{aligned}$$

³This position is neglected in the analytical literature but it is critically important in institutional-construction terms: it posits the denial to recognize another normative system which attempts to positively enact a certain power, see e.g. the Dutch Declaration of Independence, the Act of Abjuration (1581) towards Spain. [7]




Figure 3. Map of potestative relations defined in terms of triggering action (force-centered square of opposition, the left one), in terms of outcome (middle square), in terms of change or affecting outcomes (change-centered square of opposition, the right one). For visual clarity, labels of vertices are simplified so as to consist only of a (possibly negated) predicate without its arguments. Occurrences of negation before a predicate are standard, whereas occurrences after a predicate denote action complementation; for instance, we denote the power to issue a prohibition, i.e., $\text{Power}(p, q, \bar{A})$, as Power^- . Notice that the leftmost square is vertically mirrored and the rightmost square underwent a 90 degree clockwise rotation. Colours are as usual.

A *positive change-centered* power corresponds to the ability of affecting the target relation by any triggering action:

$$\text{Power}^+(x, y, A) \equiv \exists \beta : \text{Power}_{\text{OReilly}}(x, y, \beta, A)$$

A *negative change-centered* power corresponds to the ability of the agent to perform an action without affecting the target relation:

$$\text{Power}^-(x, y, A) \equiv \exists \beta : \neg \text{Power}_{\text{OReilly}}(x, y, \beta, A)$$

Labels for an Aristotelian square are, this time:

$$\text{PR}^\pm = \{ \text{Power}^+(p, q, A), \text{Power}^-(p, q, A), \neg \text{Power}^-(p, q, A), \neg \text{Power}^+(p, q, A) \}$$

Subalternity is here encoded by: $\neg \text{Power}^-(x, y, A) \rightarrow \text{Power}^+(x, y, A)$.

2.3.4. Relationships amongst powers

The previous formulas can be applied to discover different relationships between the distinct forms of powers. First, the convergence or divergence with due performance in force-centered powers map directly or dually to outcome-centered powers:

$$\overset{\rightarrow \leftarrow}{\text{Power}}(x, y, A) \rightarrow \text{Power}(x, y, A) \quad \overset{\leftarrow \rightarrow}{\text{Power}}(x, y, A) \rightarrow \text{Power}(x, y, \bar{A})$$

Following the contrapositive, the absence of positive-outcome power to produce a duty (meaning that there is no triggering action to obtain this) maps *a fortiori* to the absence of a positive-force power for doing so:

$$\neg\text{Power}(x, y, A) \xrightarrow{\leftrightarrow} \neg\text{Power}(x, y, A) \quad \neg\text{Power}(x, y, \bar{A}) \xrightarrow{\leftrightarrow} \neg\text{Power}(x, y, A)$$

Second, positive-change power holds if any outcome-center power holds:

$$\text{Power}^+(x, y, A) \leftrightarrow \text{Power}(x, y, A) \vee \text{Power}(x, y, \bar{A}) \\ \vee \overline{\text{Power}(p, q, A)} \vee \overline{\text{Power}(p, q, \bar{A})}$$

Assuming again that there is always some available action, we have, dually, that:

$$\text{Power}^-(x, y, A) \leftrightarrow \neg\text{Power}(x, y, A) \wedge \neg\text{Power}(x, y, \bar{A}) \\ \wedge \neg\overline{\text{Power}(p, q, A)} \wedge \neg\overline{\text{Power}(p, q, \bar{A})}$$

Introducing those relationships, we obtain the Aristotelian diagram in Fig. 3.

3. Diagrammatic theories

It is possible to define logical theories of different strength based on an Aristotelian diagram. These can be called *diagrammatic theories*. A diagrammatic theory over a diagram \mathcal{D} encodes at least all logical relations among formulas used as labels in \mathcal{D} . A diagrammatic theory will be here defined in terms of the notion of an *inference tree*.

Definition 3 (Inference Tree). *An inference tree T is an irreflexive and intransitive tree (N, \rightsquigarrow) with a single root ($N \neq \emptyset$ is a finite set of nodes and, for any $n, m \in N$, $n \rightsquigarrow m$ means that m is an immediate successor of n) where each $n \in N$ is associated with a finite set of formulas $\Gamma \neq \emptyset$ and has a rank. The rank of the root is $\mathbf{0}$; furthermore, if $\text{rank}(n) = \mathbf{i}$, n is associated with a set Θ , and $n \rightsquigarrow m$, then $\text{rank}(m) = \mathbf{i} + \mathbf{1}$ and m is associated with a set $\Sigma \supseteq \Theta$. Nodes with no successors are said to be leaves of T . A maximal \rightsquigarrow -chain of nodes $\sigma = (n_1, \dots, n_k)$ is a branch of T .*

Formulas in sets associated with nodes of a tree T can be uniformly substituted. Furthermore, an *equivalence relation* $Eq \subset \mathcal{L} \times \mathcal{L}$ can be established in order to replace, in any set Γ associated with a node n , a formula ϕ with a formula ψ , provided that $Eq(\phi, \psi)$.

Definition 4 (Set Immediate Inference). *If Δ is a set of formulas (associated with a node) ranked with \mathbf{i} and Δ' a set of formulas (associated with a node) ranked with $\mathbf{i} + \mathbf{1}$ in a branch σ of a tree T , then we say that Δ' can be immediately inferred from Δ within σ .*

Definition 5 (Diagrammatic Theory). *A diagrammatic theory \mathbb{DT} based on a diagram \mathcal{D} is a set of inference trees satisfying the following properties, for each formulas ϕ and ψ that label some vertex of \mathcal{D} :*

- if ψ is a subalternant of ϕ in \mathcal{D} , then some branch σ of a tree T in \mathbb{DT} encodes an inference of the form $\Delta \cup \{\phi\} \rightsquigarrow \Delta \cup \{\phi, \psi\} \cup \Gamma$;
- if ψ and ϕ are contraries in \mathcal{D} , then some branch σ of a tree T in \mathbb{DT} encodes an inference of the form $\Delta \cup \{\phi\} \rightsquigarrow \Delta \cup \{\phi, \neg\psi\} \cup \Gamma$;
- if ψ and ϕ are sub-contraries in \mathcal{D} , then some branch σ of a tree T in \mathbb{DT} encodes an inference of the form $\Delta \cup \{\neg\phi\} \rightsquigarrow \Delta \cup \{\neg\phi, \psi\} \cup \Gamma$;
- if ψ and ϕ are contradictories in \mathcal{D} , then some branch σ of a tree T in \mathbb{DT} encodes an inference of the form $\Delta \cup \{\phi\} \rightsquigarrow \Delta \cup \{\phi, \neg\psi\} \cup \Gamma$ and some branch σ' of a tree T' in \mathbb{DT} encodes an inference of the form $\Delta' \cup \{\psi\} \rightsquigarrow \Delta' \cup \{\neg\phi, \psi\} \cup \Gamma'$.

According to Definitions 4 and 5, the relation of immediate inference in a diagrammatic theory based on a diagram \mathcal{D} encodes at least all Aristotelian relations between formulas labelling vertices of \mathcal{D} . However, the use of the equivalence relation Eq allows one to reduce the number of Aristotelian relations to be encoded, as we will clarify below. Here we just assume $Eq(\phi, \neg\neg\phi)$, for every $\phi \in \mathcal{L}$.

T' is a *sub-tree* of T iff the root of T' is a node $n \in T$ and the other nodes of T' are all those that (i) occur in branches of T to which n belongs, and (ii) have a higher rank than n in those branches. The cardinality of a set of formulas Γ will be denoted as $|\Gamma|$. The notion of inference within a branch is obtained by combining the transitive closure of the notion of immediate inference and the subset relation, as indicated below.

Definition 6 (Set Inference). *A set of formulas Γ can be inferred from a set of formulas Δ within a branch σ of a tree T iff there is $\Gamma' \supseteq \Gamma$ s.t.:*

- both Δ and Γ' belong to σ ;
- the rank of Γ' in σ is not lower than the rank of Δ in σ .

According to Definition 6, if $\Gamma \subseteq \Delta$, then Γ can always be inferred from Δ within a branch of a tree. In order to check whether Γ can be inferred from Δ in a branch σ of a tree T , one has to compare pairs of sets (checking whether they are identical, one is a subset of the other, etc.). Derivation is a particular kind of inference, as per the following definition.

Definition 7 (Set Derivability - Trees). *A set of formulas Γ is derivable from a set of formulas Δ within a tree T iff there is a sub-tree T' of T whose root is a node n labelled by Δ and Γ can be inferred from Δ within all branches of T' .*

The derivability of Γ from Δ within a diagrammatic theory \mathbb{DT} is defined in terms of a finite sequence of derivations within trees of \mathbb{DT} , as below.

Definition 8 (Set Derivability - Diagrammatic Theories). *A set of formulas Γ can be derived from a set of formulas Δ within a diagrammatic theory \mathbb{DT} iff there are trees T_1, \dots, T_{n-1} in \mathbb{DT} and sets of formulas $\Delta_1, \dots, \Delta_n$ s.t.:*

- $\Delta = \Delta_1$ and $\Gamma = \Delta_n$;
- for $1 \leq j < n$, Δ_{j+1} can be derived from Δ_j within tree T_j .

Example Below is an example of an inference tree that can be used in a diagrammatic theory built over Fig. 3. It captures inferences from a set of formulas Δ including the label of the E -corner in the change-centered (rightmost) square. Each line represents a node of the tree (which, in this, case has no branches) and starts with the node's rank:

$$\begin{aligned}
 0 : \Delta_0 &= \Delta \cup \{\neg\text{Power}^+(p, q, A)\} \rightsquigarrow \\
 1 : \Delta_1 &= \Delta_0 \cup \{\text{Power}^-(p, q, A)\} \rightsquigarrow \\
 2 : \Delta_2 &= \Delta_1 \cup \{\neg\text{Power}(p, q, A) \wedge \neg\overline{\text{Power}}(p, q, A) \wedge \neg\overline{\text{Power}}(p, q, \bar{A}) \wedge \neg\text{Power}(p, q, \bar{A})\} \rightsquigarrow \\
 3 : \Delta_3 &= \Delta_2 \cup \{\neg\text{Power}(p, q, A), \neg\text{Power}(p, q, A)\} \rightsquigarrow \\
 4 : \Delta_4 &= \Delta_3 \cup \{\neg\overleftarrow{\text{Power}}(p, q, A)\}
 \end{aligned}$$

Any diagrammatic theory including this tree allows one (due to Definitions 4, 6 and 7) to derive any set $\Gamma \subseteq \Delta_i$, for $0 \leq i \leq 4$, from the starting set of formulas Δ_0 .

4. Decidability and complexity

Given two finite sets $\Gamma, \Delta \subset \mathcal{L}$, we will say that the problem of checking whether Γ can be derived from Δ within a diagrammatic theory \mathbb{DT} is the *derivability problem for finite sets* in \mathbb{DT} . We now illustrate that such a problem can be effectively computed. First, we need to define an auxiliary notion.

Definition 9 (Tree traversal). *The traversal of a tree T with reference to a formula ϕ and a set Δ is a procedure which can be described as follows (we assume that Δ occupies the root of T):*

- *Following the order of ranks, for any set of formulas Γ with rank \mathbf{i} in T , we compare ϕ with all formulas in Γ and keep track of whether ϕ occurs in Γ or not.*
- *The procedure terminates when either (positive outcome) there is a rank \mathbf{j} s.t. all sets of formulas with rank \mathbf{j} include ϕ or (negative outcome) all sets of formulas with all ranks available in T have been checked.*

Notice that, in case of a positive outcome of the traversal, due to Definitions 6 and 7, $\Delta \cup \{\phi\}$ is derivable from Δ within T . If the number of nodes in T is l , and $\max\{|\Theta \setminus \Sigma| : \text{rank}(\Theta) = \mathbf{1} + \text{rank}(\Sigma)\} = k$ (hereafter, *the maximum successor difference*), then a traversal of T with reference to ϕ and Δ requires up to $k * l$ moves.

Definition 10 (Theory traversal). *The traversal of a diagrammatic theory \mathbb{DT} with reference to a formula ϕ and a set of formulas Δ is the traversal of all trees T in \mathbb{DT} with reference to ϕ and Δ . The outcome is positive iff it is positive for some T in \mathbb{DT} .*

If the number of trees in \mathbb{DT} is h , the maximum number of nodes in a tree of \mathbb{DT} is l , and the maximum successor difference in a tree of \mathbb{DT} is k , then a traversal of \mathbb{DT} with reference to ϕ and Δ requires up to $(k * l) * h$ moves.

Theorem 1 (Decidability). *The derivability problem for finite sets within a diagrammatic theory is decidable.*

Proof. Consider two finite sets of formulas Δ and Γ , such that $\max(|\Delta|, |\Gamma|) = n$. Let \mathbb{DT} be the diagrammatic theory at issue, consisting of h inference trees. To see whether Γ is derivable from Δ , we first need to compare the sets Γ and Δ , an operation with time complexity $O(n)$. There are four relevant Cases: (1) $\Gamma = \Delta$; (2) $\Gamma \subset \Delta$; (3) $\Gamma \supset \Delta$; (4) $\Gamma \not\subset \Delta$, $\Gamma \not\supset \Delta$ and $\Gamma \neq \Delta$. In Cases 1 and 2, by Definition 8, we can immediately conclude that Γ is derivable from Δ within \mathbb{DT} . In Cases 3 and 4 one considers the set $\Gamma \setminus \Delta$. Let $|\Gamma \setminus \Delta| = m$. We know that the elements of $\Gamma \setminus \Delta$ can be enumerated ($m < n$ by construction). We take the first formula ϕ_1 in $\Gamma \setminus \Delta$ and perform a *traversal* of \mathbb{DT} with reference to ϕ_1 and Δ . If the traversal produces a negative outcome, then the whole procedure terminates and Γ is not derivable from Δ . Otherwise, there is a tree T s.t., due to Definitions 6, 7 and 9, $\Delta \cup \{\phi\}$ is derivable from Δ within T . We take $\Delta_1 = \Delta \cup \{\phi_1\}$ and then perform a *traversal* of \mathbb{DT} with reference to ϕ_2 and Δ_1 . The procedure is reiterated: at each step a traversal of \mathbb{DT} is performed with reference to a new formula $\phi_j \in \Gamma \setminus \Delta$ and the set of formulas Δ_{j-1} obtained at the previous step. In accordance with Definition 8, Γ is derivable from Δ iff each traversal of \mathbb{DT} performed ends with a positive outcome. □

Theorem 2 (Complexity). *The algorithm to solve the derivability problem for finite sets in a diagrammatic theory takes polynomial time.*

Proof. In all Cases (1-4) mentioned in the proof of Theorem 1 the procedure terminates in at most $((k * l) * h) * m$ moves. \square

5. Final remarks

Recent years have witnessed a renewed interest in diagrams and logical geometry: our work provides a further contribution towards their application in the analysis of normative problems. We introduced a syntactic method of reasoning over Aristotelian diagrams that encode relations between normative positions. The method can be automated and allows one to derive a finite set of normative positions from another in polynomial time. In our opinion, reasoning methods associated with diagrams are very promising and, due to the cognitive efficacy of visual aids, they are potentially accessible to a broader audience. Future investigations concern: (i) a theoretical and empirical comparison with other methods for (normative) automated reasoning, (ii) an integration of temporal/causal components within the definition of normative positions, and (iii) the consolidation of a structured taxonomy of Aristotelian diagrams for normative reasoning.

References

- [1] T. Parsons. The traditional square of opposition. In E.N. Zalta, editor, *The Stanford Encyclopedia of Philosophy*. Metaphysics Research Lab, Stanford University, Fall 2021 edition, 2021.
- [2] W.N. Hohfeld. Some fundamental legal conceptions as applied in judicial reasoning. *Yale Law Journal*, 23(1):16–59, 1913.
- [3] W.N. Hohfeld. Fundamental legal conceptions as applied in judicial reasoning. *Yale Law Journal*, 26(8):710–770, 1917.
- [4] A.B. Markman and D. Gentner. The effects of alignability on memory. *Psychological Science*, pages 363–367, 1997.
- [5] L.W. Sumner. *The Moral Foundation of Rights*. Clarendon Press, 1987.
- [6] D.T. O’Reilly. Using the square of opposition to illustrate the deontic and alethic relations. *University of Toronto Law Journal*, 45(3):279–310, 1995.
- [7] G. Sileno, A. Boer, and T. van Engers. On the interactional meaning of fundamental legal concepts. In R. Hoekstra, editor, *Proceedings of JURIX 2014*, pages 39–48, 2014.
- [8] M. Pascucci and G. Sileno. The search for symmetry in Hohfeldian modalities. In A. Basu, G. Stapleton, S. Linker, C. Legg, E. Manalo, and P. Viana, editors, *Diagrammatic Representation and Inference. Proceedings of Diagrams 2021*, pages 87–102. Springer, 2021.
- [9] J. A. de Oliveira Lima, C. Griffo, J. P. A. Almeida, G. Guizzardi, and M. I. Aranha. Casting the light of the theory of opposition onto hohfeld’s fundamental legal concepts. *Legal Theory*, 27(1):2–35, 2021.
- [10] L. Demey. Computing the maximal Boolean complexity of families of Aristotelian diagrams. *Journal of Logic and Computation*, 28:1323–1339, 2018.
- [11] L. Lindahl. *Position and Change: A Study in Law and Logic*. Synthese Library. Springer, 1977.
- [12] D. Makinson. On the formal representation of rights relations. *Journal of Philosophical Logic*, 15:403–425, 1986.
- [13] R. Markovich. Understanding Hohfeld and formalizing legal rights: the Hohfeldian conceptions and their conditional consequences. *Studia Logica*, 108(1):129–158, 2020.
- [14] G. Sileno and M. Pascucci. Disentangling deontic positions and abilities: a modal analysis. In F. Calimeri, S. Perri, and E. Zumpano, editors, *Proceedings of CILC 2020*, volume 2710, pages 36–50, 2020.
- [15] J.F. Sowa. *Knowledge Representation: Logical, Philosophical, and Computational Foundations*, volume 13. MIT Press, 2000.
- [16] G. Sileno. *Aligning Law and Action*. PhD thesis, University of Amsterdam, 2016.