



UvA-DARE (Digital Academic Repository)

Blocks of monodromy groups in complex dynamics

Jones, R.; Peters, H.

DOI

[10.1007/s10711-010-9499-2](https://doi.org/10.1007/s10711-010-9499-2)

Publication date

2011

Document Version

Final published version

Published in

Geometriae dedicata

[Link to publication](#)

Citation for published version (APA):

Jones, R., & Peters, H. (2011). Blocks of monodromy groups in complex dynamics. *Geometriae dedicata*, 150(1), 137-150. <https://doi.org/10.1007/s10711-010-9499-2>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Blocks of monodromy groups in complex dynamics

Rafe Jones · Han Peters

Received: 26 January 2009 / Accepted: 12 April 2010 / Published online: 27 May 2010
© The Author(s) 2010. This article is published with open access at Springerlink.com

Abstract Motivated by a problem in complex dynamics, we examine the block structure of the natural action of iterated monodromy groups on the tree of preimages of a generic point. We show that in many cases, including when the polynomial has prime power degree, there are no large blocks other than those arising naturally from the tree structure. However, using a method of construction based on real graphs of polynomials, we exhibit a non-trivial example of a degree 6 polynomial failing to have this property. This example settles a problem raised in a recent paper of the second author regarding constant weighted sums of polynomials in the complex plane. We also show that degree 6 is exceptional in another regard, as it is the lowest degree for which the monodromy group of a polynomial is not determined by the combinatorics of the post-critical set. These results give new applications of iterated monodromy groups to complex dynamics.

Keywords Iterated monodromy groups · Complex dynamics · Polynomial iteration · Post-critically finite polynomials · Conservative polynomials · Constant weighted sum of iterates

Mathematics Subject Classification (2010) 37F10 · 20B99

Rafe Jones' work was partially supported by NSF grant DMS-0852826 and Han Peters' work was partially supported by NSF grant DMS-0757856.

R. Jones (✉)
Department of Mathematics and CS, College of the Holy Cross, Worcester, MA, USA
e-mail: rjones@holycross.edu

H. Peters
Department of Mathematics, University of Amsterdam, Amsterdam, Netherlands
e-mail: h.peters@uva.nl

1 Introduction

Let G be a subgroup of the permutation group of a finite set S . A subset $E \subset S$ is called a G -block if for every permutation $\sigma \in G$ we have that $\sigma(E) \cap E$ is equal to either E or \emptyset . Motivated by a problem in a recent paper [9] of the second author we study G -blocks for monodromy groups of polynomial iterates. These groups are natural finite quotients of the iterated monodromy group of the given polynomial. The latter have attracted attention mainly for their group-theoretic interest (see e.g. [1, 2, 4, 6]), and to a lesser extent for the light they shed on the dynamics of the given polynomial (e.g. [5, Chapter 6]). The results of the present paper give a new application of iterated monodromy groups to complex dynamics, as well as results on their structure. We remark that little remains known in general about iterated monodromy groups; see [7] for some general discussion and open questions.

The monodromy group $MG_n(f)$ of the n th iterate $f^{\circ n}$ of a complex polynomial f acts by definition on the set R_n of n th preimages of a generic point. Indeed, this action extends to one on the tree of preimages of all iterates up to n . The primary goal of the present paper is to show that for many, indeed in some sense most, f there are no large $MG_n(f)$ -blocks other than those coming from the tree structure. Here is a special case of our main results in this vein (Theorems 4.1, 4.3):

Theorem 1.1 *Let $f \in \mathbb{C}[x]$ have degree $d \geq 2$, $f^{\circ n}$ be the n th iterate of f , and let the monodromy group $MG_n(f)$ of $f^{\circ n}$ act naturally on set R_n of n th preimages of a generic point. Suppose that either d is prime or $MG(f)$ acts primitively (i.e. without non-trivial blocks) on R_1 . If $E \subseteq R_n$ is any set containing a, b such that $f^{\circ n-1}(a) \neq f^{\circ n-1}(b)$, then the smallest $MG_n(f)$ -block of R_n containing E is R_n itself.*

We also prove a result on the structure of $MG_n(f)$ -blocks in the case where d is a power of a prime (Theorem 4.1).

Our other primary goal is to find polynomials f whose monodromy groups have exceptional properties, such as R_n containing small $MG_n(f)$ -blocks not coming from the tree structure. To do this, we give a method for constructing examples of critically finite f with specified critical portrait (and often even specified monodromy groups of all iterates). The method relies on building real graphs with certain properties, and gives polynomials with real coefficients. For instance, in Theorem 4.6 we give a polynomial h of degree 6 such that R_2 contains an $MG_2(f)$ -block of size 4 containing a and b with $h(a) \neq h(b)$.

A more detailed outline of the paper is as follows. In Sect. 2 we discuss how the monodromy group of a polynomial f depends on the critical portrait of f , that is, the orbits $c \rightarrow f(c) \rightarrow f^{\circ 2}(c) \rightarrow \dots$ of the critical points of f . There we only look at the monodromy group of the first iterate and we show (Theorem 2.1, 2.2) that six is the smallest degree for which the critical portrait of a polynomial does not determine the monodromy group up to conjugacy. In fact we give two explicit polynomials of degree 6 with real coefficients and identical critical orbit but whose monodromy groups are not isomorphic. The method of construction is used also in Sect. 4. In Sect. 3 we fix notation and give a few definitions.

In Sect. 4 we prove the main results of this paper. We first show (Theorem 4.1) that if f is a polynomial whose degree is prime then the only $MG_n(f)$ -blocks are branches, and when the degree of f is a power of a prime then the only $MG_n(f)$ -blocks are unions of equal-height branches, contained in a branch of height one more. Motivated by this result we call such blocks *basic blocks*; see Definition 3.1. We then show (Theorem 4.3) that if $MG_1(f)$ acts primitively on R_1 , then any $MG_n(f)$ -block containing elements that map to distinct members of R_1 must be all of R_n . We apply this to the case where f is a conservative

polynomial, i.e. all critical points are also fixed points, and f has at least two critical points (Corollary 4.4).

Then we show (Theorem 4.6) that there exist polynomials of non-prime-power degree for which there are blocks that are not basic blocks. In fact, we give an explicit example of a degree 6 polynomial which has $MG_2(f)$ -blocks that are non-basic. This polynomial is quite similar to the examples used in the proof of Theorem 2.2.

In general it can be quite complicated to compute (iterated) monodromy groups of polynomials. However the polynomial examples presented in this paper have two properties in common that make it much easier to compute their monodromy groups. Firstly, the post-critical sets are finite, which means that only a finite number of generators have to be considered. Secondly, the post-critical sets are real, which means that the action of the generators can be determined by looking only at the real graph of the polynomials.

Let us finish the introduction by discussing the motivation for this paper. In [9] the following definition was introduced:

Definition 1.2 A polynomial f has a *constant weighted sum of iterates* (c.w.s.i.) near $z \in \mathbb{C}$ if there exist weights $a_0, a_1, \dots \in \mathbb{C}$ and a constant $c \in \mathbb{C}$ such that the finite sums

$$\sum_{n=0}^N a_n f^{\circ n}$$

converge uniformly to the constant function c in a neighborhood of z as $N \rightarrow \infty$.

The requirement that the maps $\sum_{n=0}^N a_n f^{\circ n}$ converge to the same constant for any point in the neighborhood of z is very strong and one would not expect convergence to a constant to be possible for many values of z . The only cases in which it is known that f has a constant weighted sum of iterates near z is when f is affine or when z lies in a Siegel disc of f . Indeed, it was shown in [9] that a generic polynomial f of degree at least 2 has a c.w.s.i. near z if and only if z lies in a Siegel disc of f .

The proof of this result relies on the fact that the monodromy groups of the iterates of a generic polynomial (a polynomial whose iterates $f^{\circ n}$ each have the maximal number of distinct critical values) are equal to the entire group of tree automorphisms. In general the monodromy groups of the iterates may be much smaller but in many cases the monodromy groups can still be used to prove the same result. In fact, if the following question can be answered affirmatively then it still follows that a polynomial of degree at least 2 has a c.w.s.i near z if and only if z lies in a Siegel disc:

Question 1.3 Let f be a polynomial of degree at least 2 and let V be a bounded Fatou component of f that is eventually mapped onto a periodic (super-) attracting basin or attracting petal. Let $N \in \mathbb{N}$.

In the case of an attracting basin or attracting petal, let $p \in F^{\circ N}(V)$ have d^N distinct pre-images and define the sets $S = f^{\circ -N}(\{p\})$ and $E = S \cap V$. In the case of a super-attracting basin, let $U \subset f^{\circ N}(V)$ be an arbitrarily small neighborhood of the super-attracting periodic point in $f^{\circ N}(V)$. Let $p \in U$ again have d^N distinct pre-images and define $S = f^{\circ -N}(p)$ as before. Now $E \subset S$ contains only points that lie in one chosen connected component of $f^{\circ -N}(U)$ in V .

Is it possible to choose N such that the smallest $MG(f^{\circ N})$ -block of S that contains E has more than d^{N-1} elements?

The orbit of a periodic Fatou component that is a (super-) attracting basin or an attracting petal must always contain a critical point. The integer N can be chosen such that $f^N(V)$

contains the corresponding critical value. As was explained in [9], we may assume that the neighborhood U of z in Definition 1.2 is large enough so that $f^{\circ N}(U)$ contains the critical value. It follows that when p is chosen close enough to this critical value, the set E will contain two points e_1, e_2 with the property that $f^{\circ N-1}(e_1) \neq f^{\circ N-1}(e_2)$. In the terminology of Sect. 3, the points e_1 and e_2 belong to different major branches.

Theorems 4.1 and 4.3 say that we can answer Question 1.3 affirmatively when the degree of f is a power of a prime number or when $MG(f)$ acts primitively on the roots of f , but negatively for certain polynomials of degree 6. We obtain the following corollary.

Corollary 1.4 *Suppose that f is a polynomial with complex coefficients, and that either f has degree p^k with p prime, or $MG(f)$ acts primitively on the f -preimages of a generic point. Then f has a constant weighted sum of iterates near $z \in \mathbb{C}$ if and only if z is eventually mapped into a Siegel disc of f .*

The negative answer to Question 1.3 only closes off one avenue of proof when f has degree 6. It does not imply that it is possible to have a constant weighted sum of iterates when z does not lie in a Siegel disc. Whether this can occur for a polynomial of degree at least 2 is still open.

2 Combinatorics of the post-critical set

Let f be a polynomial of degree d . Denote by C the set of critical points of f , and V the set of critical values of f , that is, $V = \{f(c) : c \in C\}$. Then $f : \mathbb{C} \setminus C \rightarrow \mathbb{C} \setminus V$ is a covering, and if $p \in \mathbb{C}$ has d distinct inverse images, then every closed loop at p contained in $\mathbb{C} \setminus V$ induces a permutation on the inverse images of p . The subgroup of S_d thus obtained is independent of the point p and is called the *monodromy group* of f , denoted $MG(f)$. Let $f^{\circ n}$ denote the n th iterate of f (i.e. the n -fold composition of f with itself). We denote by $MG_n(f)$ the monodromy group of $f^{\circ n}$. It is not a general subgroup of S_{dn} , as it must respect the natural tree structure on preimages of p under f (see Sect. 3 for details).

The critical portrait (namely the number and multiplicity of the critical values) of a polynomial f gives us a large amount of information about the monodromy groups of $f^{\circ n}$. For example, let v be a critical value of a polynomial f of degree d , and denote the inverse images of v by z_1, \dots, z_n . A small enough neighborhood U of v will have n disconnected inverse images V_1, \dots, V_n with $z_j \in v_j$. Let $\gamma \in U$ be a closed loop at $p \in U$ given by a circle centered at v . Then the inverse images of γ lie in the sets V_1, \dots, V_n . If z_j is a critical point of f with order k then γ induces a full cycle on the k preimages of p in V_j .

If γ is a closed loop given by a circle centered at the origin of large enough radius then it induces a full cycle. Indeed, if the radius of γ is larger than the modulus of all the (finite) critical points then we can view it as a loop around the point at infinity, a critical point of order equal to the degree of the map.

Theorem 2.1 *The monodromy group of a polynomial of degree at most 5 is completely determined by its critical portrait, that is, the number and multiplicity of its critical values.*

Proof This is merely a case of checking all possibilities. We will do so for degree 5 and leave the smaller degrees to the reader.

Since the degree is prime and $MG(f)$ is transitive, it must be a primitive subgroup of S_5 . It is well known (see e.g. [11]) that if G is a primitive subgroup of S_n that contains a transposition, then $G = S_n$, while if G contains a 3-cycle, then G contains A_n . Thus if f has

a critical value that is the image of a single critical point of multiplicity 2, then $MG(f) = S_n$. Otherwise, by the Riemann-Hurwitz formula, f must have either one or two critical values. In the former case, $MG(f)$ has only one generator and thus $MG(f) \cong C_5$. In the latter case, each critical value must either be the image of one critical point of multiplicity 3 or two critical points of multiplicity 2. Thus $MG(f)$ may be generated by (i) two 3-cycles, (ii) one 3-cycle and one element of type $(2, 2)$ or (iii) two elements of type $(2, 2)$. In all three cases we have $MG(f) \subseteq A_5$, and for (i) and (ii) we have that $MG(f)$ contains a 3-cycle, and so $MG(f) = A_5$. In case (iii) after conjugation we may assume the generators are (12)(34) and (23)(45) and we obtain $MG(f) \cong D_5$. \square

In degree 6 the situation is more complicated:

Theorem 2.2 *There exist polynomials f, g of degree 6 with identical critical portraits but different monodromy groups. Moreover, one can choose f and g to have real coefficients.*

Remark 2.3 One can provide a short proof of Theorem 2.2 using the Riemann Existence Theorem (see [3] for an exposition), which guarantees that there is $f \in \mathbb{C}[z]$ of degree 6 with $MG(f) = \langle (23)(45), (34), (12)(56) \rangle \neq S_6$, and $g \in \mathbb{C}[z]$ of degree 6, with identical critical portrait to f and $MG(g) = \langle (23)(45), (12), (34)(56) \rangle = S_6$. However, the Riemann Existence Theorem is non-constructive, and for our purposes it will be advantageous to have an explicit method to find such f and g , since in Theorem 4.6 we need to construct a polynomial whose second iterate has prescribed monodromy. In addition, the Riemann Existence Theorem does not guarantee that f and g can be chosen to have real coefficients.

Proof We construct f and g explicitly. Both have 5 distinct critical points that all lie on the real line. By choosing our base point also on the real line and choosing our generator loops close to the real line we can use the real graphs of f and g to determine the monodromy groups. For both f and g , two critical points map to a third critical point, and this third critical point gets mapped to a fourth critical point which is a fixed point. The fifth critical point is also fixed. Therefore the critical portraits of f and g are identical.

We now describe f . Its critical points are $0, c_1, 1, c_2$ and 2 , where $0 < c_1 < 1 < c_2 < 2, f(c_1) = f(c_2) = 2, f(0) = f(2) = 0$ and $f(1) = 1$. This guarantees that f has the critical portrait described above. See Fig. 1 for the real graph of f . The existence of f can be proved by checking the degrees of freedom, but in fact a formula for f can easily be found as follows. Start with an even polynomial h that has critical points at $0, \pm 1$ and $\pm \alpha$, where $h(0) = 0$ and $h(1) = -h(\alpha)$ (taking $\alpha = \sqrt{(2 + \sqrt{3})}$ works). The function f is then given by $f(z) = 1 + \frac{h(\alpha(z-1))}{h(1)}$; note that $c_1 = 1 - 1/\alpha, c_2 = 1 + 1/\alpha$.

The critical points of g are $c_1, c_2, 0, c_3$ and 1 , where $c_1 < c_2 < 0 < c_3 < 1$. Now we require that $g(c_1) = c_1, g(c_2) = g(c_3) = 1$ and $g(1) = g(0) = 0$. Again the existence of g can be proved by counting degrees of freedom.

Note that f is the composition of two polynomials of degree 3 and 2, respectively. Therefore, one immediately sees that the monodromy group of f must have three blocks with 2 elements each. Once we determine the monodromy group of g , which has no non-trivial blocks, this will be enough to prove the Theorem. However, since the computations of $MG(f)$ and $MG(g)$ are very similar, we will discuss them both.

To determine the monodromy group of f we choose a base point $1 < p < 2$ and we denote its (real) preimages by $x_1 < \dots < x_6$. The critical values of f are $0, 1$, and 2 , and we define the corresponding generating loops γ_0, γ_1 , and γ_2 as follows (see Fig. 1). The loop γ_2 moves along the real axis from p to $2 - \epsilon$, for $\epsilon > 0$ very small, then follows a full clockwise




Fig. 1 Real graph of f and generators of the corresponding fundamental group

circle centered at 2 and goes back to p along the real axis. When following the pre-images of this loop the points x_2 and x_3 first move very close to c_1 (thus moving upwards on the real graph of f), then switch and move back to x_3 and x_2 , respectively. Similarly, x_4 and x_5 go very close to c_2 , then switch and move back to respectively x_5 and x_4 . The preimages of γ_2 starting at x_1 and x_6 form closed loops. Hence assigning to x_i the number i , we have that γ_2 induces $(23)(45) \in S_6$.

The loop γ_1 is similar: starting at p follow the real axis to $1 + \epsilon$, then loop around 1 and follow the real axis back to p . By checking the real graph of f one sees that γ_1 induces (34) . Finally, the loop γ_0 starts at p , follows the real axis to $1 + \epsilon$, follows a small semi-circle centered at 1 and contained in the lower half plane, then follows the real axis again until very close to 0, follows a full circle around 0 and then takes the same path back to p . The graph of f shows that γ_0 induces $(12)(56)$.

Thus, the monodromy group of f is generated by the elements $(23)(45)$, (34) , and $(12)(56)$. Notice that the sets $E_1 = \{1, 6\}$, $E_2 = \{2, 5\}$ and $E_3 = \{3, 4\}$ form a partition of 1, 2, 3, 4, 5, 6 that is invariant under the action γ_0 , γ_1 , and γ_2 , and hence for all of $MG(f)$. In other words E_1 , E_2 and E_3 are $MG(f)$ -blocks and $MG(f) \neq S_6$.

For the monodromy group of g we take a base point $0 < p < 1$ and we denote its pre-images by $x_1 < \dots < x_6$ as before. The critical values of g are 0, 1 and $c := c_1$, so we define loops γ_c , γ_0 and γ_1 again following the real axis as much as possible as for f , and we obtain that γ_0 induces $(34)(56)$, γ_1 induces $(23)(45)$ and γ_c induces (12) . Since the elements $(34)(56)$, $(23)(45)$ and (12) generate S_6 we get $MG(g) = S_6 \neq MG(f)$ which completes the proof. \square

3 Tree structure

The inverse images of p under f have a natural tree structure. Indeed, let $R_n = \{x \in \mathbb{C} : f^{on}(x) = p\}$, and note that




Fig. 2 A basic block, *left*, consisting of branches of height 1 contained in a single branch of height 2. *Right*, a non-basic block

$$R_n = \bigsqcup_{x \in R_{n-1}} f^{-1}(x).$$

Thus R_n may be thought of as the top level of a tree T_n , where each vertex x is connected to $f(x)$ in the next level down, namely R_{n-1} . Similarly, each element of R_{n-1} is connected to an element of R_{n-2} , and so on down to the level 0, which consists only of p . Note that p connects to exactly d vertices, vertices in the top level connect to only one other vertex, and all vertices besides p and those not in the top level connect to $d + 1$ other vertices. Setting $R_1 = \{x_1, \dots, x_d\}$, we refer to the d sets $\{f^{-(n-1)}(x_1)\}, \dots, \{f^{-(n-1)}(x_d)\}$ as the *major branches* of R_n . Note that R_n is the disjoint union of the major branches. We call a *branch* of R_n any subset of the form $f^{-m}(x)$, where $x \in R_{n-m}$. We call m the *height* of such a branch. Hence the major branches are simply the branches of height $n - 1$ and we consider the entire tree a branch of height n .

By definition, the group $MG_n(f)$ acts on R_n , and indeed we may think of it as acting on T_n since any $\gamma \in \pi_1(\mathbb{C} \setminus \{\text{critical values of } f^{\circ n}\})$ may be lifted to any vertex of the tree. A key feature of this action is that it preserves the tree structure of T_n . Indeed, $x, y \in T_n$ are connected if and only if $y = f(x)$ or $x = f(y)$; assume without loss of generality that $x = f(y)$. If γ_y is the lift of γ under $f^{\circ n}$ that starts at y , then $f(\gamma_y)$ is a (and hence the) lift of γ under $f^{\circ n-1}$ starting at x . Since $\gamma(y)$ is the endpoint of γ_y and $\gamma(x)$ is the endpoint of $f(\gamma_y)$, we have $\gamma(x) = f(\gamma(y))$, implying that $\gamma(x)$ and $\gamma(y)$ are connected. Thus the tree structure is preserved, giving an injection $MG_n(f) \hookrightarrow \text{Aut}(T_n)$.

The preservation of the tree structure of T_n immediately gives that every branch of R_n is an $MG_n(f)$ -block. It can also occur that an $MG_n(f)$ -block is not a single branch but a union of branches of equal height, contained in a single branch of height one more. The easiest polynomial for which this can occur is $f : z \mapsto z^4$. Then we have that $MG(f)$ is a cyclic group of order 4, so this group has two blocks containing two elements. From now on we will refer to such blocks as basic blocks:

Definition 3.1 Let G be a subgroup of $\text{Aut}(T_n)$ and let $E \subset R_n$ be a G -block. If E is either a single branch or is a union of equal-height branches that are all contained in a single branch of height one more, then we say that E is a *basic block*.

See Fig. 2 for an illustration of a basic block and a non-basic block. The following example shows that there are subgroups $G < \text{Aut}(T_n)$ with G -blocks that are not basic blocks.

Example 3.2 Let G be the Klein 4-group acting on the complete binary tree of height 2. It is generated by the elements $(x_{00} \ x_{01})(x_{10} \ x_{11})$ and $(x_{00} \ x_{10})(x_{01} \ x_{11})$. Then G is a

subgroup of $\text{Aut}(T_2)$, yet $\{x_{00}, x_{10}\}$ and $\{x_{01}, x_{11}\}$ form an invariant partition. Both of these are unions of height 0 branches that are not contained in a height 1 branch, and so are not basic.

As noted in Sect. 2, p. 140, the monodromy group of a polynomial always contains a full cycle. Hence the group in Example 3.2 cannot occur as the monodromy group of a polynomial. In the next section we see that for many polynomials f we only need the existence of a full cycle to deduce strong consequences for the possible structures of $MG_n(f)$ -blocks.

4 G-blocks

In this section we prove results about automorphism groups of rooted trees (Theorems 4.1 and 4.3), and then derive Corollary 1.4. We also give a negative answer to Question 1.3 in Theorem 4.6. By slight abuse of notation we use the same notation $(T_n, R_n, \text{etc.})$ as the last section to refer to abstract trees.

Theorem 4.1 *Let T_n be the complete d -ary rooted tree of height n , and let R_n denote the vertices of level n . Suppose that $G \leq \text{Aut}(T_n)$ contains a full d^n -cycle σ . If d is a prime power then every G -block of R_n is a basic block. If d is prime then every G -block of R_n is a single branch.*

Remark 4.2 It follows that a G -block of R_n that contains points lying in distinct major branches must have at least $2d^{n-1}$ elements, which settles Question 1.3 for polynomials of prime-power degrees.

Proof Let E be a G -block of R_n , and note that if $g \in G$ satisfies $g(E) \cap E \neq \emptyset$, then $g(E) = E$, and it follows that the orbit under g of any subset of E is contained in E . By choosing an ordering of the vertices at level 1 of T_n , one obtains a natural representation of every element of R_n as a base- d string of length n . For elements $v = v_1 \dots v_n$ and $w = w_1 \dots w_n$ of R_n , define the distance between v and w to be

$$D(v, w) = \begin{cases} n & \text{if } v_1 \neq w_1 \\ n - m & \text{if } m \text{ is such that } v_{m+1} \neq w_{m+1} \text{ but } v_1 \dots v_m = w_1 \dots w_m \end{cases}$$

Choose $v, w \in E$ such that $D(v, w)$ is maximal. Then E is contained in a single branch of height $t := D(v, w)$. Let $s \in \mathbb{N}$ be such that $\sigma^s(v) = w$. Then E contains the orbit of any element in V under σ^s . Moreover, s is divisible by d^{n-t} (since σ^s fixes a branch of height t) but not by d^{n-t+1} (since σ^s does not fix branches of height $t - 1$). This gives $d^{t-1} \leq |\sigma^s| \leq d^t$. But since d is a power of a prime it follows that d^{t-1} divides the order of σ^s . Thus a power of σ^s is of order d^{t-1} and induces a permutation of maximal order on each of the branches of height $t - 1$. Since E is invariant under the action of σ it follows that E is a union of branches of height $t - 1$, which completes the proof. \square

Before stating our next result, we recall a group G acts *primitively* on a set S when the only G -blocks of S consist of either one point or all of S . For instance, the symmetric group on n letters acts primitively on $\{1, \dots, n\}$. Also, every doubly transitive action is primitive.

Theorem 4.3 *Let T_n be the complete d -ary rooted tree of height n , and let R_n denote the vertices of height n . Let $E \subseteq R_n$ be any set containing points in at least two major branches. Suppose that $G \leq \text{Aut}(T_n)$ contains a full d^n -cycle σ . Suppose also that the restriction G_1 of G to the height-1 vertices R_1 acts primitively. Then the smallest G -block containing E is all of R_n .*

Proof The idea of the proof is to use the presence of σ to show that any G_n -block of R_n restricts to a G_1 -block of R_1 , which by primitivity must be large. Again using σ one shows the only lift of such a block is all of R_n .

Let A be any G -block of R_n , and denote by A_1 the restriction of A to R_1 . Let $g_1 \in G_1$ satisfy $g_1(A_1) \cap A_1 \neq \emptyset$, implying that for some $\alpha, \beta \in A_1$ we have $g_1(\alpha) = \beta$. Let $g \in G$ be any lifting of g_1 , and let B_α, B_β be the major branches of R_n corresponding to α, β , whence $g(B_\alpha) = B_\beta$. Note also that there are elements $a \in B_\alpha, b \in B_\beta$ belonging to A . Now σ^d maps each major branch to itself, and acts on each as a d^{n-1} -cycle. Thus for some m we have $\sigma^{dm}g(a) = b$. Since A is a block, it follows that $\sigma^{dm}g$ maps A to itself. But σ^{dm} restricts to the identity on R_1 , meaning that $\sigma^{dm}g$ restricts to g_1 . Thus g_1 maps A_1 to itself, proving that A_1 is a G_1 -block.

Now let E be as in the statement of the theorem, let M be the minimal G -block of R_n containing E , and let M_1 be the restriction of M to R_1 . By the previous paragraph, M_1 is a G_1 -block of R_1 , and since E contains points in different major branches, $\#M_1 \geq 2$. By the primitivity of G_1 , $M_1 = R_1$. In particular, if B and B' are major branches of R_n with $\sigma(B) = B'$ then M contains some $a \in B$ and some $b \in B'$. Thus for some m , $\sigma^{1+dm}(a) = b$. Hence the orbit of a under σ^{1+dm} is contained in M . But since $1 + dm \equiv 1 \pmod d$, σ^{1+dm} acts on R_n as a d^n -cycle. Therefore $M = R_n$. □

We note that if $G := MG_n(f)$ is cyclic and f has non-prime power order, which occurs for instance when f is conjugate to x^{pq} for primes $p < q$, then for every $n \geq 2$ there are G -blocks that are not basic. Indeed, if σ generates G then an orbit of σ^{q^n} is a G -block consisting of p^n elements. This cannot be a union of equal-height branches that is contained in a branch of height one more, since each such branch has a number of elements that is a power of pq . However, a polynomial for which the groups $G := MG_n(f)$ are cyclic will have only a single critical point that is also a fixed point. For such a map it is easy to see that the answer to Question 1.3 is positive, since all preimages of the point p will lie in the same Fatou component. To give a negative answer we have to construct a more complicated map.

As noted above, polynomials conjugate to powers of x have the strong property of having a single critical point that is also a fixed point. As a first idea for making the map somewhat more complicated, one could simply require that all critical points of f also be fixed points. Such polynomials are called *conservative*, and they have been studied in some detail [8, 10]. In particular, their conjugacy classes are determined by relatively simple combinatorial data, and there is a natural faithful action of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ on these conjugacy classes [8]. However, our next result shows that to answer Question 1.3 in the negative, conservative polynomials will not do. In Theorem 4.6 we give an example of a degree-6 polynomial that does provide a negative answer to Question 1.3.

Corollary 4.4 *Suppose that $f \in \mathbb{C}[x]$ is conservative, and f has at least two critical points. Let $G = MG_n(f)$ act naturally on the tree T_n of preimages of a suitable point p , and let $E \subset R_n$ contain points in at least two major branches of R_n . Then the smallest G -block of R_n containing E is all of R_n .*

Corollary 4.4 is an immediate consequence of the following lemma. Recall that a group G acts doubly transitively on a set S if for any $a, b, c, d \in S$ with $a \neq c$ and $b \neq d$, there is $g \in G$ with $g(a) = b$ and $g(c) = d$. It is straightforward to show that a doubly-transitive action is primitive.

Lemma 4.5 *Suppose that $f \in \mathbb{C}[x]$ is conservative, and has at least two critical points. Then $MG(f)$ acts doubly-transitively on the set R_1 of preimages under f of a generic point.*

Proof It is enough to show that for $a, b, c \in R_1$ with $a \neq b, c \neq b$, there is $g \in MG(f)$ with $g(b) = b$ and $g(a) = c$.

By assumption the set of critical points of f coincides with the set of critical values of f , which we write $\{v_1, \dots, v_n\}$. Each v_i gives an element of $MG(f)$ that acts as a single d_i -cycle, where $d_i = 1 +$ (multiplicity of the critical value v_i). Call this cycle C_i , and note that the C_i generate $MG(f)$. Note that

$$\sum_{i=1}^n d_i = n + \sum_{i=1}^n \text{multiplicity of } v_i = n + d - 1.$$

Consider the graph Γ whose vertex set consists of R_1 , and where $u \neq v \in R_1$ are connected by one edge for each C_i with $C_i(u) = v$, and also by one edge for each C_i with $C_i(v) = u$. There are no edges from any vertices to themselves. Since the action of $MG(f)$ on R_1 must be transitive (e.g. since it contains a full cycle), Γ is connected. Hence one may think of the action of $MG(f)$ on R_1 as an interlinked system of circular conveyor belts, with the key property that each belt can be made to move independently of the others.

Now obtain a new graph Γ' by deleting one edge from each cycle, which preserves connectedness. The number of vertices of Γ' is d , and the number of edges is $\sum_{i=1}^n (d_i - 1)$, which is $n + d - 1 - n = d - 1$. Denoting by F the number of regions of the plane enclosed by the edges of the graph (including the region containing infinity), by Euler's formula we have $d - (d - 1) + F = 2$. Thus $F = 1$, whence Γ' is a tree. This implies that each cycle can have at most one point in common with any other cycle. Indeed, if C_i, C_j share two distinct points a and b , then $C_i^r(a) = b$ and $C_j^s(a) = b$ with C_i^r and C_j^s not the identity. This implies that in Γ there are four distinct paths from a to b , namely those given by $C_i^r, C_i^{-r}, C_j^s, C_j^{-s}$. Deleting any edge from C_i and any from C_j thus still leaves a cycle in Γ' .

Now let $a, b, c \in R_1$ be given, with $a \neq b, c \neq b$. In the graph Γ' mentioned above, consider a path from a to c . This path may be written as the action of

$$C_{i_t}^{e_t} C_{i_{t-1}}^{e_{t-1}} \dots C_{i_0}^{e_0} \tag{1}$$

where the action is on the left, each e_j is a nonzero integer, and $t \leq n$. Choose the path to be non-self-intersecting, and so that t is minimized. This implies all the cycles in (1) are distinct, since a cycle C occurring twice violates the minimality of t , as all cycles between occurrences of C could be eliminated. Since any two cycles meet in at most one point and the path given by (1) is non-self-intersecting, it follows that any element of R_1 may be moved by at most two of the cycles in (1). Moreover, these cycles must be consecutive. If b is left fixed by all the cycles in (1), then we have moved a to c while fixing b and are done. If not, we let k be maximal so that C_{i_k} does not fix b . Suppose that b is contained in two cycles, meaning it is the unique element moved by both C_{i_k} and $C_{i_{k-1}}$. We claim the product

$$C_{i_t}^{e_t} C_{i_{t-1}}^{e_{t-1}} \dots C_{i_{k+1}}^{e_{k+1}} C_{i_{k-1}}^{-e_{k-1}} C_{i_k}^{e_k} C_{i_{k-1}}^{e_{k-1}} \dots C_{i_0}^{e_0} \tag{2}$$

maps a to c and fixes b . To show (2) maps a to c , denote by a' the image of a under $C_{i_{k-2}}^{e_{k-2}} \dots C_{i_0}^{e_0}$, and note it is enough to show

$$C_{i_{k-1}}^{-e_{k-1}} C_{i_k}^{e_k} C_{i_{k-1}}^{e_{k-1}}(a') = C_{i_k}^{e_k} C_{i_{k-1}}^{e_{k-1}}(a')$$

But $C_{i_k}^{e_k} C_{i_{k-1}}^{e_{k-1}}(a')$ is a point moved by C_{i_k} , and thus cannot be moved by $C_{i_{k-1}}$ unless it is a' , which violates the non-self-intersecting property of the path corresponding to (1). To show (2) fixes b , we need only show

$$C_{i_{k-1}}^{-e_{k-1}} C_{i_k}^{e_k} C_{i_{k-1}}^{e_{k-1}} \tag{3}$$

fixes b . To see this, note that $C_{i_{k-1}}^{e_{k-1}}$ does not fix b , thus moving b away from the unique element moved both by $C_{i_{k-1}}$ and C_{i_k} . Hence $C_{i_k}^{e_k}$ fixes $C_{i_{k-1}}^{e_{k-1}}(b)$, implying that (3) fixes b .

In the case where b is contained in only one cycle C_{i_k} of (1) and $k \neq 0$, one verifies in a similar manner that

$$C_{i_t}^{e_t} C_{i_{t-1}}^{e_{t-1}} \dots C_{i_k}^{e_k} C_{i_{k-1}}^{e_{k-1}} C_{i_k}^{-e_k} C_{i_{k-2}}^{e_{k-2}} \dots C_{i_0}^{e_0}$$

moves a to c and fixes b . To see that b is fixed, let α denote the unique point contained in both C_{i_k} and $C_{i_{k-1}}$, and note that $C_{i_k}^{e_k}(\alpha)$ is either c (if $k = t$) or the unique point contained in both C_{i_k} and $C_{i_{k+1}}$. In either case, $C_{i_k}^{-e_k}(b) \neq \alpha$, implying that $C_{i_k}^{-e_k}(b)$ is fixed by $C_{i_{k-1}}$. If $k = 0$, one checks similarly that

$$C_{i_t}^{e_t} C_{i_{t-1}}^{e_{t-1}} \dots C_{i_2}^{e_2} C_{i_0}^{-e_0} C_{i_1}^{e_1} C_{i_0}^{e_0}$$

maps a to c and fixes b .

Finally, if (1) consists of only one cycle C_0 , then choose a cycle C' that moves a point α also moved by C_0 . This is possible since $n \geq 2$ by hypothesis. If j is such that $C_0^j(b) = \alpha$, then the element

$$C_0^{-j} C'^{-1} C_0^{e_0} C' C_0^j$$

maps a to c and fixes b . □

Theorem 4.6 *There exists a polynomial h of degree 6 such that R_2 contains an $MG_2(h)$ -block with only four elements, two each in two different major branches of R_n . Moreover, h can be chosen so that it provides a negative answer to Question 1.3.*

Proof The construction of the polynomial h is very similar to the constructions in the proof of Theorem 2.2. We will assume that the reader is familiar with the polynomial f from Theorem 2.2.

Here the polynomial h will again have critical points at $0 < c_1 < 1 < c_2 < 2$ and again $h(c_1) = h(c_2) = 2$ and $h(0) = h(2) = 0$. However this time $h(1)$ is not equal to 1 but make sure that $0 < h(1) < c_1$ in such a way that $h^{\circ 2}(1) = 1$. Hence there are 4 post-critical values, namely 0, a , 1, and 2, where $a = h(1)$, and the iterated monodromy group of h has four generators.

We denote the generator loops encircling 0, a , 1, and 2, by α , β , γ , and δ , respectively. The loops are again defined by paths that stay very close to the real axis as in the proof of Theorem 2.2 (see Fig. 3). To determine the action of these generators on T , we use the method of [5, Chapter 5]. For instance, consider α . We fix paths π_i connecting the basepoint p to each x_i , and then for each x_i we compute the lift ℓ_i of α beginning at x_i . Then given a vertex of T labeled with a word iw , α returns the word $kg(w)$, where x_j is the endpoint of the lift of ℓ_i and g is the element corresponding to the loop $\pi_i \ell_i \pi_i^{-1}$. The bottom part of Fig. 3 illustrates this. We thus see that the action of α on T is given by




Fig. 3 Real graph of h , generating loops of the corresponding fundamental group (all loops are counterclockwise), and computation of the action of α on T (all connecting paths move away from p)

$$\begin{aligned}
 \alpha \cdot 1 &= 2 \cdot \text{id} \\
 \alpha \cdot 2 &= 1 \cdot \alpha \\
 \alpha \cdot 3 &= 3 \cdot \text{id} \\
 \alpha \cdot 4 &= 4 \cdot \text{id} \\
 \alpha \cdot 5 &= 6 \cdot \text{id} \\
 \alpha \cdot 6 &= 5 \cdot \delta
 \end{aligned}$$

Hence in wreath recursion notation, we have $\alpha = \langle \text{id}, \alpha, \text{id}, \text{id}, \text{id}, \delta \rangle (1 \ 2)(5 \ 6)$. In many cases one can read the action of an element of the fundamental group directly from the real graph. For instance, to compute the action of γ , note that following preimages of γ is the same as moving the x_i down along the real graph of h towards 1. Since none of the x_i are near to each other when $y = 1$, the action of γ on the first level of T is trivial. When the y -values of these points reaches $1 + \epsilon$, a counterclockwise loop is executed (in the complex plane, and so not visibly on the real graph). One sees that since x_2 is near a , this loop must encircle a . For the others, the loops encircle no critical values, and so are homotopically trivial. Thus,

$$\gamma = \langle \text{id}, \beta, \text{id}, \text{id}, \text{id}, \text{id} \rangle.$$




Fig. 4 Action of $IMG(h)$ on the first two levels of T

Similar computations yield

$$\beta = \langle \text{id}, \text{id}, \text{id}, \gamma, \text{id}, \text{id} \rangle (3 \ 4)$$

$$\delta = \langle \text{id}, \text{id}, \text{id}, \text{id}, \text{id}, \text{id} \rangle (2 \ 3)(4 \ 5)$$

The action of $IMG(h)$ on the first two levels of T is pictured in Fig. 4. Note that the action of $IMG(h)$ on the first level of the tree is the same as the action of the monodromy group of f from Theorem 2.2.

Label a vertex on level two of T by x_{ij} where i denotes the major branch and j denotes the position within the i th major branch. Consider the partition consisting of

$$\{x_{ij}, x_{i(7-j)}, x_{(7-i)(4-j)}, x_{(7-i)(3+j)}\} \quad i = 2, 3, j = 1, 2, 3$$

as well as $\{x_{11}, x_{16}, x_{62}, x_{65}\}$, $\{x_{12}, x_{15}, x_{63}, x_{64}\}$, and $\{x_{13}, x_{14}, x_{61}, x_{66}\}$. One checks that all of $\alpha, \beta, \gamma, \delta$ respect this partition, and thus we have a set of G -blocks each containing four elements. Note that none of these blocks is basic.

To show that h provides a negative answer to Question 1.3, note that a is a super-attracting periodic point so it is contained in a Fatou component V that is a super-attracting basin. If N is chosen odd then p must be chosen in the Fatou component containing 1, which does not contain a critical value. One easily sees that all preimages $h^{\circ-N}(p)$ in V lie in the same major branch which is a $MG_N(h)$ -block and contains 6^{N-1} elements.

If $N = 2$ and thus $p \in V$ then the preimages $f^{\circ-2}(p)$ that lie in V correspond to the points x_{32}, x_{42} which lie in the $MG_2(h)$ -block $E = \{x_{32}, x_{35}, x_{42}, x_{45}\}$. So the smallest $MG_2(h)$ -block containing those two points must be contained in E (and is in fact equal to

E). E has only 4 elements which is fewer than $6^1 + 1 = 7$. If N is even but larger than 2 then again p must be chosen in V and all preimages of p in V are also preimages of points in E . That gives at most $4 \times 6^{N-2}$ elements and thus fewer than $6^{N-1} + 1$. This completes the proof. \square

Acknowledgments The authors wish to thank Laurent Bartholdi for reading a preliminary version of the article and for several helpful comments.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

1. Bartholdi, L., Virág, B.: Amenability via random walks. *Duke Math. J.* **130**(1), 39–56 (2005)
2. Bux, K.-U., Pérez, R.: On the growth of iterated monodromy groups. In: *Topological and Asymptotic Aspects of Group Theory. Contemp. Math.*, vol. 394, pp. 61–76. Amer. Math. Soc., Providence, RI, (2006)
3. Dèbes, P.: Méthodes topologiques et analytiques en théorie inverse de Galois: théorème d’existence de Riemann. In: *Arithmétique de revêtements algébriques (Saint-Étienne, 2000)*, Sémin. Congr. vol. 5, pp. 27–41. Soc. Math. France, Paris, (2001)
4. Kaimanovich, V.A.: “Münchhausen trick” and amenability of self-similar groups. *Internat. J. Algebra Comput.* **15**(5–6), 907–937 (2005)
5. Nekrashevych, V.: *Self-Similar Groups*, vol. 117. *Mathematical Surveys and Monographs*. American Mathematical Society, Providence (2005)
6. Nekrashevych, V.: A minimal Cantor set in the space of 3-generated groups. *Geom. Dedicata* **124**, 153–190 (2007)
7. Nekrashevych, V.: Combinatorics of polynomial iterations. In: *Complex Dynamics*, pp. 169–214. A K Peters, Wellesley (2009)
8. Pakovich, F.: Conservative polynomials and yet another action of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ on plane trees. *J. Théor. Nombres Bordeaux* **20**(1), 205–218 (2008)
9. Peters, H.: Constant weighted sums of iterates. *Complex Var. Elliptic Equ* **54**(3–4), 371–386 (2009)
10. Tischler, D.: Critical points and values of complex polynomials. *J. Complexity* **5**(4), 438–456 (1989)
11. Wielandt, H.: *Finite Permutation Groups*. Translated from the German by R. Bercov. Academic Press, New York (1964)