



## UvA-DARE (Digital Academic Repository)

### Unraveling the cold response in Draba

von Meijenfeldt, N.

**Publication date**  
2010

[Link to publication](#)

**Citation for published version (APA):**

von Meijenfeldt, N. (2010). *Unraveling the cold response in Draba*. [Thesis, fully internal, Universiteit van Amsterdam].

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

## ABBREVIATIONS USED:

<i>AP2-domain</i>	<i>APETALA2-domain</i> amino acid sequence thought to function as DNA-binding domain.
<i>CAB</i>	<i>Chlorophyll a/b binding factor</i> transcription factor encoding the light-harvesting chlorophyll a/b-binding proteins of photosystem II and is involved in the diurnal and circadian rhythm of plants. <i>CAB</i> levels peak in the late morning.
<i>CCR</i>	<i>Cold-circadian rhythm-RNA binding factor</i> transcription factor involved in the circadian rhythm of plants. <i>CCR</i> levels accumulate in the early evening.
<i>CRT/DRE</i>	<i>C-repeat (CRT) dehydration responsive element (DRE)</i> a DNA regulatory element with a conserved core sequence of CCGAC that imparts responsiveness to low temperature and dehydration.
<i>CBF</i>	<i>C-repeat binding factor</i> transcription factor that binds to the CRT/DRE element and is involved in the early cold response of plants.
<i>COR</i>	<i>Cold responsive genes</i> genes containing the CRT/DRE element, which are regulated by <i>CBF</i> and confer freezing tolerance.
<i>HOS</i>	<i>High expression of osmotically responsive genes</i> a constitutively expressed transcription factor, that through mediating <i>ICE</i> expression negatively regulates the expression of <i>CBF</i> .
<i>ICE</i>	<i>Inducer of CBF expression</i> a constitutively expressed transcription factor critical for the activation of <i>CBF</i> expression, but which is inactive under non-stress conditions.
<i>NLS</i>	<i>Nuclear localization signal</i> amino acid sequence used to target the protein to the cell nucleus.
<i>ZAT = STZ</i>	<i>Salt tolerance zinc finger</i> transcription factor induced by dehydration, high-salt, abscisic acid and cold stress.