



UvA-DARE (Digital Academic Repository)

Unraveling the cold response in Draba

von Meijenfeldt, N.

[Link to publication](#)

Citation for published version (APA):
von Meijenfeldt, N. (2010). *Unraveling the cold response in Draba*.

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

R

References

- Abbott RJ, Brochmann C (2003) History and evolution of the arctic flora: in the footsteps of Eric Hulten. *Molecular Ecology* 12: 299-313
- Al-Shehbaz IA (1991) Novelties in *Draba* (Brassicaceae) from Venezuela, Ecuador, and Peru. *Novon* 1: 67-70
- Al-Shehbaz IA, Beilstein MA, Kellogg EA (2006) Systematics and phylogeny of the Brassicaceae (Cruciferae): an overview. *Plant Systematics & Evolution* 259: 89
- Alonso-Blanco C, Gomez-Mena C, Llorente F, Koornneef M, Salinas J, Martinez-Zapater JM (2005) Genetic and molecular analyses of natural variation indicate *CBF2* as a candidate gene for underlying a freezing tolerance quantitative trait locus in *Arabidopsis*. *Plant Physiology* 139: 1304-1312
- Andel J van, Rozijn NAMG, Ernst WHO, Nelissen HJM (1986) Variability in growth and reproduction in F1-families of an *Erophila verna* population. *Oecologia* 69: 79-85
- Anisko T, Lindstrom OM (1995) Applying the Richards function in freezing tolerance determination with electrolyte and phenolic leakage techniques. *Physiologia Plantarum* 95: 281-287
- Artus NN, Uemura M, Steponkus PL, Gilmour SJ, Lin C, Thomashow MF (1996) Constitutive expression of the cold-regulated *Arabidopsis thaliana* *COR15a* gene affects both chloroplast and protoplast freezing tolerance. *PNAS* 93: 13404-13409
- Azocar A, Rada F, Goldstein G (1988) Freezing tolerance In *Draba chionophila*, a miniature caulescent rosette species. *Oecologia* 75: 156-160
- Baker SS, Wilhelm KS, Thomashow MF (1994) The 5'-region of *Arabidopsis thaliana* *COR15a* has cis-acting elements that confer cold-, drought- and ABA-regulated gene expression. *Plant Molecular Biology* 24: 701-713
- Beck E (1994) Cold tolerance in tropical alpine plants. In *Tropical Alpine Environments – Plant form and function*. Cambridge University Press, Cambridge, pp 77-110
- Beck E, Senser M, Scheibe R, Steiger H-M, Pongratz P (1982) Frost avoidance and freezing tolerance in Afroalpine 'giant rosette' plants. *Plant, Cell and Environment* 5: 215-222
- Beilstein MA, Windham MD (2003) A phylogenetic analysis of western North American *Draba* (Brassicaceae) based on nuclear ribosomal DNA sequences from the ITS region. *Systematic Botany* 28: 584-592
- Benderoth M, Textor S, Windsor AJ, Mitchell-Olds T, Gershenzon J, Kroymann J (2006) Positive selection driving diversification in plant secondary metabolism. *PNAS* 103: 9118-9123
- Benedict C, Skinner JS, Meng R, Chang Y, Bhalerao R, Huner NPA, Finn CE, Chen THH, Hurry V (2006) The *CBF1*-dependent low temperature signalling pathway, regulon and increase in freeze tolerance are conserved in *Populus* spp. *Plant, Cell & Environment* 29: 1259-1272
- Bieniawska Z, Espinoza C, Schlereth A, Sulpice R, Hinch DK, Hannah MA (2008) Disruption of the *Arabidopsis* circadian clock is responsible for extensive variation in

- the cold-responsive transcriptome. *Plant Physiology* 147: 263-279
- Billings WD (1974) Adaptations and origins of alpine plants. *Arctic and Alpine Research* 6: 129-142
- Brochmann C (1993) Reproductive strategies of diploid and polyploid populations of arctic *Draba* (Brassicaceae). *Plant Systematics and Evolution* 185: 55-83
- Brochmann C, Borgen L, Stedje B (1993) Crossing relationships and chromosome-numbers of Nordic populations of *Draba* (Brassicaceae), with emphasis on the *D. alpina* complex. *Nordic Journal of Botany* 13: 121-147
- Brochmann C, Elven R (1992) Ecological and genetic consequences of polyploidy in arctic *Draba* (Brassicaceae). *Evolutionary Trends in Plants* 6: 111-124
- Carpenter CD, Kreps JA, Simon AE (1994) Genes encoding glycine-rich *Arabidopsis thaliana* proteins with RNA-binding motifs are influenced by cold treatment and an endogenous circadian rhythm. *Plant Physiology* 104: 1015-1025
- Chinnusamy V, Ohta M, Kanrar S, Lee B-h, Hong X, Agarwal M, Zhu J-K (2003) *ICE1*: a regulator of cold-induced transcriptome and freezing tolerance in *Arabidopsis*. *Genes & Development* 17: 1043-1054
- Chinnusamy V, Zhu J, Zhu J-K (2007) Cold stress regulation of gene expression in plants. *Trends in Plant Science* 12: 444
- Choi D-W, Rodriguez EM, Close TJ (2002) Barley *CBF3* gene identification, expression pattern, and map location. *Plant Physiology* 129: 1781-1787
- Cook D, Fowler S, Fiehn O, Thomashow MF (2004) From the cover: A prominent role for the *CBF* cold response pathway in configuring the low-temperature metabolome of *Arabidopsis*. *PNAS* 101: 15243-15248
- Dong C-H, Agarwal M, Zhang Y, Xie Q, Zhu J-K (2006) The negative regulator of plant cold responses, *HOS1*, is a RING E3 ligase that mediates the ubiquitination and degradation of *ICE1*. *PNAS*: 8281-8286
- Doyle JJ, Doyle JL (1987) A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemistry Bulletin* 19: 11-15
- El Kayal W, Navarro M, Marque G, Keller G, Marque C, Teulieres C (2006) Expression profile of *CBF*-like transcriptional factor genes from *Eucalyptus* in response to cold. *Journal of Experimental Botany* 57: 2455-2469
- Fowler S, Thomashow MF (2002) *Arabidopsis* transcriptome profiling indicates that multiple regulatory pathways are activated during cold acclimation in addition to the *CBF* cold response pathway. *Plant Cell* 14: 1675-1690
- Fowler SG, Cook D, Thomashow MF (2005) Low temperature induction of *Arabidopsis CBF1*, 2, and 3 is gated by the circadian clock. *Plant Physiology* 137: 961-968
- Gao M-J, Allard G, Byass L, Flanagan AM, Singh J (2002) Regulation and characterization of four *CBF* transcription factors from *Brassica napus*. *Plant Molecular Biology* 49: 459-471
- Gavel Y, von Heijne G (1990) A conserved cleavage-site motif in chloroplast transit peptides. *FEBS Letters* 261: 455-458
- Gilmour SJ, Artus NN, Thomashow MF (1992) cDNA sequence analysis and expression of two cold-regulated genes of *Arabidopsis thaliana*. *Plant Molecular Biology* 18: 13-21
- Gilmour SJ, Fowler SG, Thomashow MF (2004) *Arabidopsis* transcriptional activators *CBF1*, *CBF2*, and *CBF3* have matching functional activities. *Plant Molecular Biology* 54: 767-781

- Gilmour SJ, Sebolt AM, Salazar MP, Everard JD, Thomashow MF (2000) Overexpression of the *Arabidopsis* *CBF3* transcriptional activator mimics multiple biochemical changes associated with cold acclimation. *Plant Physiology* 124: 1854-1865
- Gilmour SJ, Zarka DG, Stockinger EJ, Salazar MP, Houghton JM, Thomashow MF (1998) Low temperature regulation of the *Arabidopsis* *CBF* family of AP2 transcriptional activators as an early step in cold-induced *COR* gene expression. *Plant Journal* 16: 433-442
- Griffith M, Timonin M, Wong ACE, Gray GR, Akhter SR, Saldanha M, Rogers MA, Weretilnyk EA, Moffatt B (2007) *Thellungiella*: an *Arabidopsis*-related model plant adapted to cold temperatures. *Plant, Cell & Environment* 30: 529-538
- Grundt HH, Kjolner S, Borgen L, Rieseberg LH, Brochmann C (2006) High biological species diversity in the arctic flora. *PNAS* 103: 972-975
- Grundt HH, Obermayer R, Borgen LIV (2005) Ploidal levels in the arctic-alpine polyploid *Draba lactea* (Brassicaceae) and its low-ploid relatives. *Botanical Journal of the Linnean Society* 147: 333-347
- Grundt HH, Popp M, Brochmann C, Oxelman B (2004) Polyploid origins in a circumpolar complex in *Draba* (Brassicaceae) inferred from cloned nuclear DNA sequences and fingerprints. *Molecular Phylogenetics and Evolution* 32: 695-710
- Guindon S, Gascuel O (2003) A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Systematic Biology* 52: 696
- Guy CL (1990) Cold acclimation and freezing stress tolerance: role of protein metabolism. *Annual Review of Plant Physiology and Plant Molecular Biology* 41: 187-223
- Guy CL, Huber JLA, Huber SC (1992) Sucrose phosphate synthase and sucrose accumulation at low temperature. *Plant Physiology* 100: 502-508
- Haake V, Cook D, Riechmann JL, Pineda O, Thomashow MF, Zhang JZ (2002) Transcription factor *CBF4* is a regulator of drought adaptation in *Arabidopsis*. *Plant Physiology* 130: 639-648
- Hajela RK, Horvath DP, Gilmour SJ, Thomashow MF (1990) Molecular cloning and expression of *COR* (cold-regulated) genes in *Arabidopsis thaliana*. *Plant Physiology* 93: 1246-1252
- Hannah MA, Heyer AG, Hinch DK (2005) A global survey of gene regulation during cold acclimation in *Arabidopsis thaliana*. *PLoS Genetics* 1: e26
- Hannah MA, Wiese D, Freund S, Fiehn O, Heyer AG, Hinch DK (2006) Natural genetic variation of freezing tolerance in *Arabidopsis*. *Plant Physiology* 142: 98-112
- Harmer SL, Hogenesch JB, Straume M, Chang HS, Han B, Zhu T, Wang X, Kreps JA, Kay SA (2000) Orchestrated transcription of key pathways in *Arabidopsis* by the circadian clock. *Science* 290: 2110
- Hughes C, Eastwood R (2006) From the Cover: Island radiation on a continental scale: Exceptional rates of plant diversification after uplift of the Andes. *PNAS* 103: 10334-10339
- Ito Y, Katsura K, Maruyama K, Taji T, Kobayashi M, Seki M, Shinozaki K, Yamaguchi-Shinozaki K (2006) Functional analysis of rice *DREB1/CBF*-type transcription factors involved in cold-responsive gene expression in transgenic rice. *Plant Cell Physiology* 47: 141-153
- Jaglo-Ottosen KR, Gilmour SJ, Zarka DG, Schabenberger O, Thomashow MF (1998) *Arabidopsis* *CBF1* overexpression induces *COR* genes and enhances freezing tolerance. *Science* 280: 104-106

- Jaglo KR, Kleff S, Amundsen KL, Zhang X, Haake V, Zhang JZ, Deits T, Thomashow MF (2001) Components of the *Arabidopsis* C-repeat/dehydration-responsive element binding factor cold-response pathway are conserved in *Brassica napus* and other plant species. *Plant Physiology* 127: 910-917
- Janska A, Marsik P, Zelenkova S, Ovesna J (2010) Cold stress and acclimation – what is important for metabolic adjustment? *Plant Biology* 12: 395-405
- John UP, Polotnianka RM, Sivakumaran KA, Chew O, Mackin L, Kuiper MJ, Talbot JP, Nugent GD, Mautord J, Schrauf GE, Spangenberg GC (2009) Ice recrystallization inhibition proteins (IRIPs) and freeze tolerance in the cryophilic Antarctic hair grass *Deschampsia antarctica* E. Desv. *Plant, Cell & Environment* 32: 336-348
- Jordon-Thaden I, Hase I, Al-Shehbaz I, Koch MA (2010) Molecular phylogeny and systematics of the genus *Draba* (Brassicaceae) and identification of its most closely related genera. *Molecular Phylogenetics and Evolution* 55: 524-540
- Jordon-Thaden I, Koch M (2008) Species richness and polyploid patterns in the genus *Draba* (Brassicaceae): a first global perspective. *Plant Ecology & Diversity* 1: 255-263
- Jørgensen PM, Léon-Yanez S (1999) Catalogue of the vascular plants of Ecuador. In W³CEC; Web version: <http://www.mobot.org/MOBOT/research/ecuador/search.shtml>. Missouri Botanical Garden Press
- Jørgensen PM, Ulloa Ulloa C (1994) Seed Plants of the High Andes of Ecuador – a checklist, Vol 34. Aarhus University Press, Aarhus
- Kaplan F, Kopka J, Sung DY, Zhao W, Popp M, Porat R, Guy CL (2007) Transcript and metabolite profiling during cold acclimation of *Arabidopsis* reveals an intricate relationship of cold-regulated gene expression with modifications in metabolite content. *The Plant Journal* 50: 967-981
- Karlsón DT, Zeng Y, Stirm VE, Joly RJ, Ashworth EN (2003) Photoperiodic regulation of a 24-kD dehydrin-like protein in red-osier dogwood (*Cornus sericea* L.) in relation to freeze-tolerance. *Plant and Cell Physiology* 44: 25-34
- Kasuga M, Liu Q, Miura S, Yamaguchi-Shinozaki K, Shinozaki K (1999) Improving plant drought, salt, and freezing tolerance by gene transfer of a single stress-inducible transcription factor. *Nature Biotechnology* 17: 287-291
- Katoh K, Toh H (2008) Recent developments in the MAFFT multiple sequence alignment program. *Brief Bioinform* 9: 286-298
- Kimura M, Yamamoto YY, Seki M, Sakurai T, Sato M, Abe T, Yoshida S, Manabe K, Shinozaki K, Matsui M (2009) Identification of *Arabidopsis* genes regulated by high light-stress using cDNA microarray. *Photochemistry and Photobiology* 77: 226-233
- Klotke J, Kopka J, Gatzke N, Heyer AG (2004) Impact of soluble sugar concentrations on the acquisition of freezing tolerance in accessions of *Arabidopsis thaliana* with contrasting cold adaptation – evidence for a role of raffinose in cold acclimation. *Plant, Cell & Environment* 27: 1395-1404
- Knight H, Veale EL, Warren GJ, Knight MR (1999) The *sfr6* mutation in *Arabidopsis* suppresses low-temperature induction of genes dependent on the CRT/DRE sequence motif. *Plant Cell* 11: 875-886
- Knight H, Zarka DG, Okamoto H, Thomashow MF, Knight MR (2004) Abscisic acid induces *CBF* gene transcription and subsequent induction of cold-regulated genes via the CRT promoter element. *Plant Physiology* 135: 1710-1717

- Koch M, Al-Shehbaz IA (2002) Molecular data indicate complex intra- and intercontinental differentiation of American *Draba* (Brassicaceae). *Annals of the Missouri Botanical Garden* 89: 88
- Körner C (2003) Climatic stress. *In* *Alpine Plant Life; Functional Plant Ecology of High Mountain Ecosystems*. Springer, pp 101-119
- Körner C (2003) Life under snow: protection and limitation. *In* *Alpine Plant Life; Functional Plant Ecology of High Mountain Ecosystems*. Springer, pp 47-62
- Kreps JA, Simon AE (1997) Environmental and genetic effects on circadian clock-regulated gene expression in *Arabidopsis*. *Plant Cell* 9: 297-304
- Labarga A, Valentin F, Anderson M, Lopez R (2007) Web services at the European Bioinformatics Institute. *Nucleic Acids Research*: 1-6
- Lambrecht SC, Shattuck AK, Loik ME (2007) Combined drought and episodic freezing effects on seedlings of low- and high-elevation subspecies of sagebrush (*Artemisia tridentata*). *Physiologia Plantarum* 130: 207-217
- Larcher W (2003) Plants under stress. *In* *Physiological plant ecology: ecophysiology and stress physiology of functional groups*, Ed 4th. Springer, pp 345-450
- Le MQ, Engelsberger WR, Hinch DK (2008) Natural genetic variation in acclimation capacity at sub-zero temperatures after cold acclimation at 4°C in different *Arabidopsis thaliana* accessions. *Cryobiology* 57: 104-112
- Lee H, Guo Y, Ohta M, Xiong LM, Stevenson B, Zhu JK (2002) *LOS2*, a genetic locus required for cold-responsive gene transcription encodes a bi-functional enolase. *Embo Journal* 21: 2692-2702
- Lee H, Xiong L, Gong Z, Ishitani M, Stevenson B, Zhu J-K (2001) The *Arabidopsis HOS1* gene negatively regulates cold signal transduction and encodes a RING finger protein that displays cold-regulated nucleo cytoplasmic partitioning. *Genes & Development* 15: 912-924
- Lee YP, Fleming AJ, Ch K, F. Meins J (2009) Differential expression of the *CBF* pathway and cell cycle-related genes in *Arabidopsis* accessions in response to chronic low-temperature exposure. *Plant Biology* 11: 273-283
- León-Yáñez S, Pitman N (2003) *Draba hookeri*. *In* IUCN Red List of Threatened Species, Vol Version 2009.2. IUCN
- Librado P, Rozas J (2009) DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25: 1451-1452
- Lin C, Thomashow MF (1992) A cold-regulated *Arabidopsis* gene encodes a polypeptide having potent cryoprotective activity. *Biochemical and Biophysical Research Communications* 183: 1103-1108
- Lin C, Thomashow MF (1992) DNA sequence analysis of a complementary DNA for cold-regulated *Arabidopsis* gene *cor15* and characterization of the COR15 polypeptide. *Plant Physiology* 99: 519-525
- Lin Y-H, Hwang S-Y, Hsu P-Y, Chiang Y-C, Huang C-L, Wang C-N, Lin T-P (2008) Molecular population genetics and gene expression analysis of duplicated *CBF* genes of *Arabidopsis thaliana*. *BMC Plant Biology* 8: 111
- Liu Q, Kasuga M, Sakuma Y, Abe H, Miura S, Yamaguchi-Shinozaki K, Shinozaki K (1998) Two transcription factors, *DREB1* and *DREB2*, with an EREBP/AP2 DNA binding domain separate two cellular signal transduction pathways in drought- and low-temperature-responsive gene expression, respectively, in *Arabidopsis*. *Plant Cell* 10: 1391-1406

- Liu S, Wang X, Fan Z, Pang Y, Sun X, Wang X, Tang K (2004) Molecular cloning and characterization of a novel cold-regulated gene from *Capsella bursa-pastoris*. *DNA Sequence* 15: 262-268
- Loik ME, Redar SP (2003) Microclimate, freezing tolerance, and cold acclimation along an elevation gradient for seedlings of the Great Basin Desert shrub, *Artemisia tridentata*. *Journal of Arid Environments* 54: 769
- Luteyn JL (1999) Introduction to the páramo ecosystem. In *Páramos – a checklist of plant diversity, geographical distribution and botanical literature*. The New York Botanical Garden, Bronx, New York, pp 1-39
- Marcotte EM, Pellegrini M, Yeates TO, Eisenberg D (1999) A census of protein repeats. *Journal of Molecular Biology* 293: 151-160
- Maruyama K, Sakuma Y, Kasuga M, Ito Y, Seki M, Goda H, Shimada Y, Yoshida S, Shinozaki K, Yamaguchi-Shinozaki K (2004) Identification of cold-inducible downstream genes of the *Arabidopsis DREB1A/CBF3* transcriptional factor using two microarray systems. *The Plant Journal* 38: 982-993
- McKhann H, Gery C, Berard A, Leveque S, Zuther E, Hinch D, De Mita S, Brunel D, Teoule E (2008) Natural variation in *CBF* gene sequence, gene expression and freezing tolerance in the Versailles core collection of *Arabidopsis thaliana*. *BMC Plant Biology* 8: 105
- Medina J, Bargues M, Terol J, Perez-Alonso M, Salinas J (1999) The *Arabidopsis CBF* gene family is composed of three genes encoding AP2 domain-containing proteins whose expression is regulated by low temperature but not by abscisic acid or dehydration. *Plant Physiology* 119: 463-470
- Michelmore RW, Meyers BC (1998) Clusters of resistance genes in plants evolve by divergent selection and a birth-and-death process. *Genome Research* 8: 1113-1130
- Millar AJ, Kay SA (1991) Circadian control of *Cab* gene-transcription and messenger-RNA accumulation in *Arabidopsis*. *Plant Cell* 3: 541-550
- Millar AJ, Kay SA (1996) Integration of circadian and phototransduction pathways in the network controlling *CAB* gene transcription in *Arabidopsis*. *PNAS* 93: 15491-15496
- Murray MB, Cape JN, Fowler D (1989) Quantification of frost damage in plant tissues by rates of electrolyte leakage. *New Phytologist* 113: 307-311
- Nakayama K, Okawa K, Kakizaki T, Honma T, Itoh H, Inaba T (2007) *Arabidopsis Cor15am* is a chloroplast stromal protein that has cryoprotective activity and forms oligomers. *Plant Physiology* 144: 513-523
- Navarro M, Marque G, Ayax C, Keller G, Borges JP, Marque C, Teulieres C (2009) Complementary regulation of four *Eucalyptus CBF* genes under various cold conditions. *Journal of Experimental Botany* 60: 2713-2724
- Novillo F, Alonso JM, Ecker JR, Salinas J (2004) *CBF2/DREB1C* is a negative regulator of *CBF1/DREB1B* and *CBF3/DREB1A* expression and plays a central role in stress tolerance in *Arabidopsis*. *PNAS* 101: 3985-3990
- Novillo F, Medina J, Salinas J (2007) *Arabidopsis CBF1* and *CBF3* have a different function than *CBF2* in cold acclimation and define different gene classes in the *CBF* regulon. *PNAS* 104: 21002-21007
- Ouellet F (2007) Cold acclimation and freezing tolerance in plants. *Encyclopedia of Life Sciences*: 1-6
- Pfitsch WA (1988) Microenvironment and the distribution of two species of *Draba* (Brassicaceae) in a Venezuelan páramo. *Arctic and Alpine Research* 20: 333-341

- Pfitsch WA (1994) Morphological and physiological radiation in páramo *Draba*. In PW Rundel, AP Smith, FC Meinzer, eds, Tropical Alpine Environments – Plant Form and Function. Cambridge University Press, Cambridge
- Ramsay PM, Oxley ERB (1997) The growth form composition of plant communities in the ecuadorian páramos. *Plant Ecology* 131: 173-192
- Rapacz M, Wolanin B, Hura K, Tyrka M (2008) The effects of cold acclimation on photosynthetic apparatus and the expression of *COR14b* in four genotypes of barley (*Hordeum vulgare*) contrasting in their tolerance to freezing and high-light treatment in cold conditions. *Annals of Botany* 101: 689-699
- Ratcliffe D (1960) *Draba muralis* L. *Journal of Ecology* 48: 737-744
- Reyes-Diaz M, Ulloa N, Zuniga-Feest A, Gutierrez A, Gidekel M, Alberdi M, Corcuera LJ, Bravo LA (2006) *Arabidopsis thaliana* avoids freezing by supercooling. *Journal of Experimental Botany* 57: 3687-3696
- Ristic Z, Ashworth EN (1993) Changes in leaf ultrastructure and carbohydrates in *Arabidopsis thaliana* L. (Heyn) cv. Columbia during rapid cold acclimation. *Protoplasma* 172: 111
- Rizzon C, Ponger L, Gaut BS (2006) Striking similarities in the genomic distribution of tandemly arrayed genes in *Arabidopsis* and rice. *PLoS Computational Biology* 2: e115
- Roche P, Diaz-Burlinson N, Gachet S (2004) Congruency analysis of species ranking based on leaf traits: which traits are the more reliable? *Plant Ecology* 174: 37-48
- Rohde P, Hinch DK, Heyer AG (2004) Heterosis in the freezing tolerance of crosses between two *Arabidopsis thaliana* accessions (Columbia-0 and C24) that show differences in non-acclimated and acclimated freezing tolerance. *Plant Journal* 38: 790-799
- Ruelland E, Vaultier MN, Zachowski A, Hurry V (2009) Cold signalling and cold acclimation in plants. In *Advances in Botanical Research*, Vol 49, pp 35-150
- Sakamoto H, Maruyama K, Sakuma Y, Meshi T, Iwabuchi M, Shinozaki K, Yamaguchi-Shinozaki K (2004) *Arabidopsis* Cys2/His2-type zinc-finger proteins function as transcription repressors under drought, cold, and high-salinity stress conditions. *Plant Physiology* 136: 2734-2746
- Sakuma Y, Liu Q, Dubouzet JG, Abe H, Shinozaki K, Yamaguchi-Shinozaki K (2002) DNA-binding specificity of the ERF/AP2 domain of *Arabidopsis* DREBs, transcription factors involved in dehydration- and cold-inducible gene expression. *Biochemical and Biophysical Research Communications* 290: 998
- Savitch LV, Allard G, Seki M, Robert LS, Tinker NA, Huner NPA, Shinozaki K, Singh J (2005) The effect of overexpression of two *Brassica* CBF/DREB1-like transcription factors on photosynthetic capacity and freezing tolerance in *Brassica napus*. *Plant and Cell Physiology* 46: 1525-1539
- Schranz ME, Song BH, Windsor AJ, Mitchell-Olds T (2007) Comparative genomics in the Brassicaceae: a family-wide perspective. *Current Opinion in Plant Biology* 10: 168-175
- Seki M, Kamei A, Yamaguchi-Shinozaki K, Shinozaki K (2003) Molecular responses to drought, salinity and frost: common and different paths for plant protection. *Current Opinion in Biotechnology* 14: 194-199
- Seki M, Narusaka M, Abe H, Kasuga M, Yamaguchi-Shinozaki K, Carninci P, Hayashizaki Y, Shinozaki K (2001) Monitoring the expression pattern of 1300 *Arabidopsis* genes under drought and cold stresses by using a full-length cDNA microarray. *Plant Cell* 13: 61-72

- Sharma N, Cram D, Huebert T, Zhou N, Parkin IAP (2007) Exploiting the wild crucifer *Thlaspi arvense* to identify conserved and novel genes expressed during a plant's response to cold stress. *Plant Molecular Biology* 63: 171-184
- Shinozaki K, Yamaguchi-Shinozaki K (2000) Molecular responses to dehydration and low temperature: differences and cross-talk between two stress signaling pathways. *Current Opinion in Plant Biology* 3: 217-223
- Shinwari ZK, Nakashima K, Miura S, Kasuga M, Seki M, Yamaguchi-Shinozaki K, Shinozaki K (1998) An *Arabidopsis* gene family encoding DRE/CRT binding proteins involved in low-temperature-responsive gene expression. *Biochemical and Biophysical Research Communications* 250: 161-170
- Sklenár P, Balslev H (2005) Superpáramo plant species diversity and phytogeography in Ecuador. *Flora – Morphology, Distribution, Functional Ecology of Plants* 200: 416-433
- Sklenar P, Ramsay PM (2001) Diversity of zonal páramo plant communities in Ecuador. *Diversity and Distributions* 7: 113-124
- Skrede I, Borgen L, Brochmann C (2008) Genetic structuring in three closely related circumpolar plant species: AFLP versus microsatellite markers and high-arctic versus arctic-alpine distributions. *Heredity* 102: 293-302
- Slater RJ (1984) The extraction of total RNA by the detergent and phenol method. *In Nucleic Acids*, Vol 2, pp 101-108
- Smallwood M, Bowles DJ (2002) Plants in a cold climate. *Philosophical transactions of the Royal Society of London. B* 357: 831-847
- Smith JMB, Cleef AM (1988) Composition and origins of the world's tropical alpine floras. *Journal of Biogeography* 15: 631-645
- Squeo FA, Rada F, Azocar A, Goldstein G (1991) Freezing tolerance and avoidance in high tropical Andean plants: Is it equally represented in species with different plant height? *Oecologia* 86: 378
- Squeo FA, Rada F, García C, Ponce M, Rojas A, Azócar A (1996) Cold resistance mechanisms in high desert Andean plants. *Oecologia* 105: 552
- Steponkus PL (1984) Role of the plasma-membrane in freezing-injury and cold-acclimation. *Annual Review of Plant Physiology and Plant Molecular Biology* 35: 543-584
- Steponkus PL, Uemura M, Joseph RA, Gilmour SJ, Thomashow MF (1998) Mode of action of the *COR15a* gene on the freezing tolerance of *Arabidopsis thaliana*. *PNAS* 95: 14570-14575
- Stockinger EJ, Gilmour SJ, Thomashow MF (1997) *Arabidopsis thaliana CBF1* encodes an AP2 domain-containing transcriptional activator that binds to the C-repeat/DRE, a cis-acting DNA regulatory element that stimulates transcription in response to low temperature and water deficit. *PNAS* 94: 1035-1040
- Strand Å, Foyer CH, Gustafsson P, Gardeström P, Hurry V (2003) Altering flux through the sucrose biosynthesis pathway in transgenic *Arabidopsis thaliana* modifies photosynthetic acclimation at low temperatures and the development of freezing tolerance. *Plant, Cell & Environment* 26: 523-535
- Thomashow MF (1998) Role of cold-responsive genes in plant freezing tolerance. *Plant Physiology* 118: 1-8
- Thomashow MF (1999) Plant cold acclimation: freezing tolerance genes and regulatory mechanisms. *Annual Review of Plant Physiology and Plant Molecular Biology* 50: 571-599
- Thomashow MF (2001) So what's new in the field of plant cold acclimation? Lots! *Plant Physiology* 125: 89-93

- Tropicos.org (2010) Tropicos.org. 06 April 2010. In <<http://www.Tropicos.org>>. Missouri Botanical Garden.
- Uemura M, Joseph RA, Steponkus PL (1995) Cold acclimation of *Arabidopsis thaliana* (Effect on plasma membrane lipid composition and freeze-induced lesions). *Plant Physiology* 109: 15-30
- Ulloa Ulloa C, Álvarez Molina S, Jørgensen PM, Minga D (2004) Cien plantas silvestres del páramo. Parque Nacional Cajas-ETAPA & Missouri Botanical Garden, Cuenca, Ecuador
- Vogel JT, Zarka DG, Van Buskirk HA, Fowler SG, Thomashow MF (2005) Roles of the *CBF2* and *ZAT12* transcription factors in configuring the low temperature transcriptome of *Arabidopsis*. *The Plant Journal* 41: 195-211
- Wang H, Cutler AJ (1995) Promoters from *KINI* and *COR6.6*, two *Arabidopsis thaliana* low-temperature-inducible and ABA-inducible genes, direct strong beta-glucuronidase expression in guard-cells, pollen and young developing seeds. *Plant Molecular Biology* 28: 619-634
- Wang X, Liu L, Liu S, Sun X, Deng Z, Pi Y, Tang K (2004) Isolation and molecular characterization of a new CRT binding factor gene from *Capsella bursa-pastoris*. *Journal of Biochemistry and Molecular biology* 37: 538-545
- Wang Y, Hua J (2009) A moderate decrease in temperature induces *COR15a* expression through the *CBF* signaling cascade and enhances freezing tolerance. *The Plant Journal* 60: 340-349
- Wang Z, Triezenberg SJ, Thomashow MF, Stockinger EJ (2005) Multiple hydrophobic motifs in *Arabidopsis CBF1* COOH-terminus provide functional redundancy in trans-activation. *Plant Molecular Biology* 58: 543
- Wanner LA, Junttila O (1999) Cold-induced freezing tolerance in *Arabidopsis*. *Plant Physiology* 120: 391-400
- Warwick S, Al-Shehbaz I (2006) Brassicaceae: Chromosome number index and database on CD-Rom. *Plant Systematics and Evolution* 259: 237-248
- Webb MS, Uemura M, Steponkus PL (1994) A comparison of freezing injury in oat and rye: two cereals at the extremes of freezing tolerance. *Plant Physiology* 104: 467-478
- Weeda EJ, Westra R, Westra C, Westra T (2003) Wilde planten en hun relaties. In *Nederlandse Oecologische Flora*. KNNV Uitgeverij / IVN, pp 33-34
- Welling A, Palva ET (2008) Involvement of *CBF* transcription factors in winter hardiness in birch. *Plant Physiology* 147: 1199-1211
- Wilhelm KS, Thomashow MF (1993) *Arabidopsis thaliana cor15b*, an apparent homologue of *cor15a*, is strongly responsive to cold and ABA, but not drought. *Plant Molecular Biology* 23: 1073-1077
- Xiao H, Siddiqua MK, Braybrook S, Nassuth A (2006) Three grape *CBF/DREB1* genes respond to low temperature, drought and abscisic acid. *Plant, Cell and Environment* 29: 1410-1421
- Xin Z, Browse J (1998) *eskimo1* mutants of *Arabidopsis* are constitutively freezing-tolerant. *PNAS* 95: 7799-7804
- Xin Z, Browse J (2000) Cold comfort farm: the acclimation of plants to freezing temperatures. *Plant, Cell & Environment* 23: 893-902
- Xin Z, Mandaokar A, Chen J, Last RL, Browse J (2007) *Arabidopsis ESK1* encodes a novel regulator of freezing tolerance. *The Plant Journal* 49: 786-799
- Xiong L, Schumaker KS, Zhu J-K (2002) Cell signaling during cold, drought, and salt stress. *Plant Cell* 14: 165-183

- Xiong Y, Fei S-Z (2006) Functional and phylogenetic analysis of a *DREB/CBF*-like gene in perennial ryegrass (*Lolium perenne* L.). *Planta* 224: 878
- Yamaguchi-Shinozaki K, Shinozaki K (1994) A novel cis-acting element in an *Arabidopsis* gene is involved in responsiveness to drought, low-temperature, or high-salt stress. *Plant Cell* 6: 251-264
- Yamaguchi-Shinozaki K, Shinozaki K (2006) Transcriptional regulatory networks in cellular responses and tolerance to dehydration and cold stresses. *Annual Review of Plant Biology* 57: 781-803
- Yuanyuan M, Yali Z, Jiang L, Hongbo S (2009) Roles of plant soluble sugars and their responses to plant cold stress. *African Journal of Biotechnology* 8: 2004-2010
- Zarka DG, Vogel JT, Cook D, Thomashow MF (2003) Cold induction of *Arabidopsis CBF* genes involves multiple ICE (Inducer of *CBF* Expression) promoter elements and a cold-regulatory circuit that is desensitized by low temperature. *Plant Physiology* 133: 910-918
- Zhang X, Fowler SG, Cheng H, Lou Y, Rhee SY, Stockinger EJ, Thomashow MF (2004) Freezing-sensitive tomato has a functional *CBF* cold response pathway, but a *CBF* regulon that differs from that of freezing-tolerant *Arabidopsis*. *Plant Journal* 39: 905-919
- Zhen Y, Ungerer MC (2008) Clinal variation in freezing tolerance among natural accessions of *Arabidopsis thaliana*. *New Phytologist* 177: 419-427
- Zhen Y, Ungerer MC (2008) Relaxed selection on the *CBF/DREB1* regulatory genes and reduced freezing tolerance in the southern range of *Arabidopsis thaliana*. *Molecular Biology and Evolution* 25: 2547-2555
- Zhou D, Zhou J, Meng L, Wang Q, Xie H, Guan Y, Ma Z, Zhong Y, Chen F, Liu J (2009) Duplication and adaptive evolution of the *COR15* genes within the highly cold-tolerant *Draba* lineage (Brassicaceae). *Gene* 441: 36-44
- Zhou N, Robinson SJ, Huebert T, Bate NJ, Parkin IAP (2007) Comparative genome organization reveals a single copy of *CBF* in the freezing tolerant crucifer *Thlaspi arvense*. *Plant Molecular Biology* 65: 693-705
- Zhu J, Shi H, Lee B-h, Damsz B, Cheng S, Stirm V, Zhu J-K, Hasegawa PM, Bressan RA (2004) An *Arabidopsis* homeodomain transcription factor gene, *HOS9*, mediates cold tolerance through a *CBF*-independent pathway. *PNAS* 101: 9873-9878
- Zuther E, Büchel K, Hundertmark M, Stitt M, Hinch DK, Heyer AG (2004) The role of raffinose in the cold acclimation response of *Arabidopsis thaliana*. *FEBS Letters* 576: 169-173