UvA-DARE (Digital Academic Repository)

Optical counterpart of the LMXB GX 13+1

Groot, P.J.; van der Klis, M.; van Paradijs, J.A.; Augusteijn, T.; Berger, M.

Publication date
1996

Published in
Cataclysmic variables and related objects

[Link to publication](#)

Citation for published version (APA):

Groot, P. J., van der Klis, M., van Paradijs, J. A., Augusteijn, T., & Berger, M. (1996). Optical counterpart of the LMXB GX 13+1. In J. H. Wood, & A. Evans (Eds.), *Cataclysmic variables and related objects* (pp. 367-368)

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

OPTICAL COUNTERPART OF THE LMXB GX 13+1

P. J. GROOT¹, M. VAN DER KLIS¹, J. VAN PARADIJS^{1,2},
T. AUGUSTEIJN³, M. BERGER¹

*1. Astronomical Institute 'Anton Pannekoek'/CHEAF,
University of Amsterdam, Amsterdam, The Netherlands*

2. Physics Department, University of Alabama, Alabama, USA

3. ESO, La Silla, Chile

We have observed the IR counterpart of the Atoll type LMXB GX 13+1, (Naylor, Charles & Longmore 1991), from 1993 July 11...30, with the Dutch 90 cm telescope at ESO using a Gunn z ($0.9\text{--}1.0\ \mu\text{m}$) filter. Over this period 110 usable images (integration time 12 m) were obtained. The data were reduced using the ESO-MIDAS/DAPHOT reduction package. A Lomb-Scargle period search was performed with the PERIOD package.

Fig. 1 shows our Gunn z band image with stars labeled as in Naylor et al. (1991). They propose their star 101 as the IR counterpart. Fig. 2 shows the light curve we obtained for this star. The star is clearly variable on a time-scale of ~ 10 d. To estimate any period more accurately we have made a Lomb-Scargle statistics periodogram (see Fig. 3); maximum power is at 12.6 d. We estimate the error to be ~ 1 day.

The time interval covered by our observations (18 d) is too short to establish the periodicity of the variation. If the orbital period is long as is suggested by our results it may be difficult to understand the difference between Atoll and Z sources on the basis of simple evolutionary arguments (see e.g. Lewin, van Paradijs & van der Klis 1988).

Acknowledgements. We thank Jeroen de Jong, Vik Dhillon and Gene Magnier for the use of their programs.

References

- Naylor, T., Charles, P.A., Longmore, A.J., 1991, MNRAS, **252**, 203
Lewin, H.G., van Paradijs, J., van der Klis, M., 1988, Sp. Sc. Rev., **46**, 273

Figure 1. Gunn z band image of GX 13+1. Star 101 is the IR counterpart.

Figure 2. Light curve of star 101 in Fig. 1. Figure 3. Lomb-Scargle statistics power spectrum of the light curve in Fig. 2. Magnitudes in units of average magnitude of comparison stars.