



## UvA-DARE (Digital Academic Repository)

### Clinical measures trump neurocognition in predicting long-term outcome for adolescents at ultra-high risk for psychosis

Ziermans, T.; de Wit, S.; Schothorst, P.; Sprong, M.; van Engeland, H.; Kahn, R.; Durston, S.

**DOI**

[10.1016/S0924-9338\(13\)75938-8](https://doi.org/10.1016/S0924-9338(13)75938-8)

**Publication date**

2013

**Document Version**

Final published version

**Published in**

European psychiatry

[Link to publication](#)

**Citation for published version (APA):**

Ziermans, T., de Wit, S., Schothorst, P., Sprong, M., van Engeland, H., Kahn, R., & Durston, S. (2013). Clinical measures trump neurocognition in predicting long-term outcome for adolescents at ultra-high risk for psychosis. *European psychiatry*, 28(Supplement 1), Article 717. [https://doi.org/10.1016/S0924-9338\(13\)75938-8](https://doi.org/10.1016/S0924-9338(13)75938-8)

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

*UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)*

## CLINICAL MEASURES TRUMP NEUROCOGNITION IN PREDICTING LONG-TERM OUTCOME FOR ADOLESCENTS AT ULTRA-HIGH RISK FOR PSYCHOSIS

*T. Ziermans*<sup>1,2</sup>, *S. de Wit*<sup>2</sup>, *P. Schothorst*<sup>2</sup>, *M. Sprong*<sup>2</sup>, *H. van Engeland*<sup>2</sup>, *R. Kahn*<sup>2</sup>, *S. Durston*<sup>2</sup>

<sup>1</sup>Karolinska Institutet, Stockholm, Sweden, <sup>2</sup>University Medical Center Utrecht, Utrecht, The Netherlands

**Background:** Most studies aiming to predict transition to psychosis for individuals at ultra-high risk (UHR) have focused on either neurocognitive or clinical variables and have made little effort to combine the two. We aimed to investigate the relative value of neurocognitive and clinical variables for predicting transition to psychosis as well as long-term functional outcome.

**Methods:** Sixty-seven adolescents at UHR and 72 controls completed an extensive clinical and neurocognitive assessment. Forty-three UHR individuals and 47 controls participated in long-term follow-up approximately six years later. UHR adolescents who had converted to psychosis (UHR-P) were compared to individuals who had not (UHR-NP) and controls on clinical and neurocognitive variables. Regression analyses were performed to determine which baseline measures best predicted transition to psychosis and long-term functional outcome for UHR individuals.

**Results:** Low IQ was the single neurocognitive parameter that discriminated UHR-P individuals from UHR-NP individuals and controls. The severity of attenuated positive symptoms was the only significant predictor of a transition to psychosis and disorganized symptoms were highly predictive of functional outcome.

**Conclusions:** IQ was lowest for those individuals at ultra high risk for psychosis who later went on to have a psychotic episode. However, IQ was not a good predictor of either transition or functional outcome. Rather, clinical measures proved to be the most important vulnerability markers for long-term outcome.