UvA-DARE (Digital Academic Repository)

Forest ecosystem development on degraded and reclaimed sites.

Fanta, J.

Publication date
1996

Published in
Ecological Engineering

[Link to publication](#)

Citation for published version (APA):

Fanta, J. (1996). Forest ecosystem development on degraded and reclaimed sites. *Ecological Engineering*, 3(spec.issue).

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Editorial

Forest ecosystem development on degraded and reclaimed sites

J. Fanta ^{a,b}

^a Dept. of Physical Geography and Soil Science, University of Amsterdam, Netherlands

^b DLO Institute for Forestry and Nature Research, Wageningen, Netherlands

Forests are irreplaceable as stabilizing elements in landscapes degraded by grazing, mining, severe acid deposition and other anthropogenic activities. Afforestation has been and still is one of the most usual measures applied in land reclamation in these degraded areas.

Forest development on degraded and reclaimed sites is a difficult problem for present-day forestry. The application of conventional silvicultural measures is limited and in many cases impossible because of the specific site and vegetation conditions and phenomena such as destabilized artificial relief, undeveloped raw soil substratum or absence of soil, absence of a proper hydrological regime, chemically altered soils, absence of soil biota, etc. Under such circumstances, technical measures are often necessary (although they often prove insufficient as well) to establish new forests and to ensure their survival and stabilizing function in degraded landscapes.

Until recently, the processes of the spontaneous development of forest ecosystems on degraded and reclaimed sites have not received adequate attention from researchers, although the knowledge of these processes is necessary for functional landscape reclamation, forestry and nature conservation. In recent reclamation plans, however, more attention has been given to ecological criteria. Hence, knowledge of the spontaneous processes in forests on these sites is becoming ever more important.

The workshop “Forest ecosystems development on degraded and reclaimed sites”, organized by the IUFRO working group “Forest dynamics” (2–4 September 1992) in the framework of the Centennial IUFRO Congress in Berlin, Germany, explored this difficult problem. The workshop focused especially on the “Black Triangle”, the area straddling the borders of Saxony, Bohemia and Silesia, with its very topical problems of extensive forest decline and chemical degradation of forest sites caused by acid deposition (Fig. 1).

A selection of the main papers presented during the workshop is brought together in this issue of *Ecological Engineering*.

Fig. 1. The "Black Triangle" in Central Europe along the borders of Germany, Czech Republic and Poland.

The papers focused on:

- processes of chemical soil degradation and spontaneous soil development on reclaimed sites,
- spontaneous forest succession on degraded and reclaimed sites, and
- forest growth and development in heavily polluted areas.

The authors J. Hruška and P. Krám investigated the mechanism of chemical degradation of shallow granitic soils caused by acid deposition. In the long run, the upper soil horizons became depleted of mobilisable cations (bases and aluminium), input of acids was unbuffered, and the main cation in soil solution remained the hydrogen ion. As a result, the soil solution was strongly acid and surface runoff from the catchment was high in aluminium concentration. This mechanism can explain the overall decline of vitality of the second generation Norway spruce forests in the area.

The effect of heavy metal pollution on forest soils is another factor contributing to their degradation. J. Zwoliński established a significant negative correlation between the heavy metal content and the rates of dead organic material in Scots pine forests in the surroundings of a zinc smelter and a copper plant. The toxic effect of heavy metals on decomposition increased with the depth of the soil

profile (L–F–H). The results clearly document the negative influence of heavy metals on soil biota in the organic profile, resulting at least in an extensive decline of forest vitality.

In contrast, soil microbiologists A. Lettl and J. Hýsek document a revival of soil microbiological activity during spontaneous successional processes following forest dieback and resulting in a strong development of grassy vegetation (*Calamagrostis villosa*, *Deschampsia flexuosa*) and, later on, spontaneous invasion of pioneering tree species (*Betula pendula*, *Sorbus aucuparia*). The microbial communities increased greatly in the organic horizons under grasses and in spontaneous stands of birch and mountain ash, and the mineralization of carbon and nitrogen compounds was more intensive than under the dead spruce forest.

P. Pyšek investigated successional changes in grassy vegetation on clearings following forest dieback. In the beginning, *Calamagrostis villosa* was the most successful colonizer on sites rich or poor in organic matter. After 5 years of dominance it was gradually replaced by *Deschampsia flexuosa* on humus-poor sites but retained its dominance on sites with a thick layer of organic matter. On these sites, this species can prevent immigration of both plant and tree species, causing a delay in spontaneous succession.

The contribution of K. Prach deals with the potential for spontaneous succession of woody species on degraded sites and derelict land in Central Europe. The most successful colonizers are anemochorous and zoochorous tree species. Succession of trees is site-specific and highly variable because many adverse abiotic and biotic factors influence the process of establishment, survival and initial growth of seedlings and saplings (e.g. dryness, toxicity of substratum and/or competition, animal browsing). Despite this, the spontaneous regeneration of woody species might often play an important role in reclamation.

In the last contribution, P. Moravčík analyses young stands of *Picea excelsa* and *P. pungens* (both planted), *Betula verrucosa* (sown) and *Sorbus aucuparia* (spontaneous) in an area with a high acid deposition. Compared to commercial conifers, the pioneering broadleaves do not suffer from industrial immissions, have a high biomass production and encourage soil development. This result indicates that spontaneous regeneration of pioneering broadleaves and their role in reclamation of sites heavily affected by acid deposition should be given more attention.

The contributions clearly indicate that degradation of forests and forest sites in areas heavily affected by industrial pollution is a complicated process. They testify to the need for an integrated, fundamental research to this difficult problem. Understanding the mechanisms and dynamics of degradation is the crucial condition for a successful reclamation of degraded forests and sites. An ecosystem approach is necessary to get insight into ecological relations in ecosystems under stress and to reveal the role of particular components and ecosystem compartments in the process. Processes of spontaneous succession and development of forests deserve much more attention than they have received to date, especially because of their role in developing a higher ecological complexity of forest ecosystems and in supporting the stability of forests and sites in polluted areas.