



UvA-DARE (Digital Academic Repository)

Multiple faces of *Fusarium oxysporum* effector protein Avr2

Di, X.

Publication date

2017

Document Version

Other version

License

Other

[Link to publication](#)

Citation for published version (APA):

Di, X. (2017). *Multiple faces of Fusarium oxysporum effector protein Avr2*. [Thesis, fully internal, Universiteit van Amsterdam].

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Chapter 1

General introduction

General introduction

Vascular wilt disease and its control

Vascular wilt diseases caused by soil-borne fungi are among the most devastating plant diseases worldwide (Tjamos and Beckman, 1989). After penetrating the root and colonizing the vasculature, the pathogen is shielded from most fungicides prohibiting curative treatments of infected plants. Due to the presence of persistent resting structures, such as chlamydospores, vascular fungi exhibit extreme longevity in the soil making crop-rotation futile (Michielse and Rep, 2009). One control measure is soil sterilization, which is expensive and harmful to the soil, making it applicable only on small scales such as in greenhouses. An alternative method of disease control is the use of resistant plant varieties generated by plant breeding. However, when resistances are introduced into elite varieties, they are often overcome by the pathogen in the course of time (Michielse and Rep, 2009). The high economic impact of wilt diseases, combined with the lack of sustainable control treatments, substantiates the need for alternative control strategies. To design novel strategies a thorough understanding of the molecular mechanism underlying disease-susceptibility and -resistance to vascular pathogens is of fundamental importance and the main aim of this thesis is to contribute to this.

Plant innate immune response

To defend themselves against pathogens plants rely on a sophisticated innate immune system (de Wit, 2007). Two distinct layers can be distinguished: the first is based on the perception of conserved pathogen-associated molecular patterns (PAMPs) from pathogens by host pattern-recognition receptors (PRR) (Nurnberger et al., 2004; Zipfel and Felix, 2005). One of the best-characterized PAMP/PRR pair is the FLAGELLIN SENSING 2 (FLS2) receptor mediating recognition of bacterial flagellin (or the elicitor-active peptide flg22 derived from it). PAMP recognition and subsequent activation of the PRR, triggers a basal defence response (also called pattern-triggered immunity - PTI) that wards off attacks by most non-adapted pathogens (Jones and Dangl, 2006). Adapted pathogens can evade or subvert PTI by secretion of so-called "effector" proteins that manipulate host-targets involved in basal defence (Dodds and Rathjen, 2010). Some plants however, carry genes encoding "resistance" proteins that recognise specific effectors allowing them to initiate a second layer of inducible defences, which is called effector-triggered immunity (ETI) (Jones and Dangl, 2006). Pathogens that overcome ETI have often shed, or mutated, the cognate effector thereby evading recognition by the corresponding resistance (R) protein. The outcomes of ETI include an oxidative burst, production of anti-microbial compounds, expression of pathogenesis-related (PR) genes, and often a rapid programmed localized cell death (PCD) response

at the site of infection (Spoel and Dong, 2012). The ETI reaction as a whole is denoted the hypersensitive response (HR) (Mur et al., 2008; Hofius et al., 2011). Notably, the defense responses that are activated during ETI and PTI are partly overlapping, including alterations of phytohormone levels, production of reactive oxygen species (ROS), changes in intracellular calcium levels, transcriptional reprogramming and synthesis of antimicrobial compounds (Coll et al., 2011).

The *Fusarium*-tomato pathosystem

Fusarium oxysporum (*Fo*) is a ubiquitous occurring fungus that inhabits the soil and rhizosphere or even colonizes and lives within plant tissues without causing disease symptoms (Swarupa et al., 2014). Actually, many *Fo* strains confer beneficial effects to their host (Alabouvette et al., 2009; Edel-Hermann et al., 2015; Imazaki and Kadota, 2015). In these cases *Fo* colonization remains symptomless and has beneficial effects on plant growth and/or tolerance to biotic and abiotic stresses. However, a small number of forms evolved into pathogens that cause disease on specific host plants (Michielse and Rep, 2009). In many of these pathogenic interactions a lengthy asymptomatic phase precedes disease development (Demers et al., 2015). Thus, based on its lifestyle pathogenic *Fo* strains are typically classified as hemi-biotrophs, shifting from a biotrophic phase early in infection to necrotrophy at later stages. These pathogenic strains typically produce toxins at later stages of disease development in order to kill the host cells, as to complete their life cycle on dead tissues (Horbach et al., 2011). Pathogenic *Fo* strains have been classified in over 120 formae speciales, which refer to their respective plant hosts, as a particular *forma specialis* (f.sp.) typically produces disease in one or a limited range of host species only (Armstrong and Armstrong, 1981; Katan and Di Primo, 1999). Over past decades the interaction between *Fo* f.sp. *lycopersici* (*Fol*) and tomato (*Solanum lycopersicum*) has evolved into an excellent model to study the molecular mechanisms underlying disease and resistance (Takken and Rep, 2010). Upon recognition *Fol* attaches to the roots, colonizes the cortex and subsequently penetrates the endodermis. It then rapidly colonizes the xylem vessels and eventually the xylem-adjacent tissues (di Pietro et al., 2003). Successful infection causes typical disease symptoms such as yellowing, wilting, stunted growth and in some cases plant death (di Pietro et al., 2003; Agrios, 2005).

Phytohormones such as salicylic acid (SA), ethylene (ET), jasmonic acid (JA), abscisic acid (ABA) and auxin are signaling molecules that are not only essential for regulation of plant growth, development and reproduction, but they also play a vital role in adaptive responses to a wide variety of biotic and abiotic stresses (Bari and Jones, 2009; Grant and Jones, 2009; Pieterse et al., 2009; Pieterse et al., 2012). Plants typically respond to pathogen infection with a complex scenario of sequential, antagonistic or synergistic

action of different hormone signals (Robert-Seilaniantz et al., 2011). In *Arabidopsis thaliana*, various mutants and transgenic lines impaired in hormone biosynthesis, perception or signaling display a severe alteration in their level of susceptibility to *Fo* (Berrocal-Lobo and Molina, 2008; Pieterse et al., 2009; Thatcher et al., 2009; Trusov et al., 2009). For instance, *Arabidopsis NahG* (containing a bacterial salicylate hydroxylase transgene) and *sid2* (salicylic acid induction deficient 2) lines, in which SA accumulation is impaired, exert increased susceptibility to *Fo* f.sp. *conglutinans* (*Focn*) pointing at the involvement of SA in disease susceptibility (Berrocal-Lobo and Molina, 2004; Diener and Ausubel, 2005). ET is also involved in susceptibility, but its role is opposite to that of SA. *Arabidopsis* mutants *ein2* (ethylene insensitive 2) and *etr1-1* (ethylene receptor 1), both ET insensitive, show a reduction in disease symptoms as compared to Col-0 plants, when inoculated with *Focn* (Trusov et al., 2009; Pantelides et al., 2013). The involvement of phytohormones in plant susceptibility following *Fo* infection has mostly been studied in the model *Arabidopsis thaliana*. Their role in modulating susceptibility of crops, such as tomato, in relation to *Fo* infection has not systematically been studied yet. As their activity can be different in different plant species, it is worthwhile to investigate involvement of these hormones in the *Fol*-tomato pathosystem.

***Fol* effectors and tomato R proteins**

During colonization of its host *Fol* secretes many small proteins into the xylem sap (Rep, 2005). Using mass-spectrometry our lab identified 14 putative effector proteins in the xylem sap of infected susceptible tomato plants, named "Secreted In Xylem" (SIX) proteins (Rep et al., 2005; Houterman et al., 2007; Houterman et al., 2009; Schmidt et al., 2013). For Six1, Six3, Six5 and Six6 a virulence function has been revealed (Rep et al., 2005; Houterman et al., 2009; Gawehns et al., 2014; Ma et al., 2015). Besides, some effectors have been found to act as avirulence factors, triggering resistance gene-mediated immunity in tomato. The resistance thus brought about, fits the "gene-for-gene" model, in which a plant *R* gene requires a "matching" avirulence (*Avr*) gene in the pathogen (Figure 1). So far, three *R* genes against *Fol* have been introgressed into commercially cultivated tomato varieties: genes *I* and *I-2* from *S. pimpinellifolium*, conferring resistance against *Fol* races 1 and 2, respectively, and the *I-3* gene from *S. pennellii*, which confers resistance to *Fol* race 3 (Huang and Lindhout, 1997). Recently, all these three *R* genes have been cloned (Simons et al., 1998; Catanzariti et al., 2015; Catanzariti et al., 2016). The three genes encoding the *Fol* effector proteins Six4 (*Avr1*), Six3 (*Avr2*) and Six1 (*Avr3*), which are recognized by R proteins I, I-2 and I-3, respectively, had already been cloned before (Rep et al., 2004; Houterman et al., 2008; Houterman et al., 2009). The *Avr3* gene encodes a 32kDa protein, which includes a 2kDa signal peptide, an 8kDa N-terminal prodomain and a 22kDa mature protein that represents the C-terminal part of the protein (Rep et al., 2005; van der Does et al., 2008). *Avr2*

encodes a small 15.7kDa protein that is processed after cleavage of its signal peptide (Houterman et al., 2007; Houterman et al., 2009). *Avr3* is expressed when the fungus is in contact with living plant cells (van der Does et al., 2008), while *Avr2* is predominantly expressed in xylem-colonizing hyphae and to some extent in hyphae that reside in the root cortex (Ma et al., 2013). Both *Avr3* and *Avr2* contribute to pathogenicity as deletion of either *Avr2* or *Avr3* in *Fol* greatly reduced virulence of the knockout strains on susceptible tomato plants (Rep et al., 2005; Houterman et al., 2009). Like *Avr3*, the *Avr1* gene encodes a 27kDa product with a signal peptide and a predicated prodomain (Houterman et al., 2007; Houterman et al., 2008). After processing a 20kDa mature protein remains. Notably, *Avr1* does not contribute to fungal virulence on susceptible plants, but it suppresses *I-2* and *I-3* mediated resistance allowing the fungus to cause disease on plants carrying the cognate resistance genes (Houterman et al., 2008). None of the three *Avr* proteins shares significant homology to any known protein and their mode of actions in the host are unknown (Fraser-Smith et al., 2014).

Avr2 is expressed by the fungus following penetration of the root cortex and during colonization of the xylem vessels in which the encoded protein can be readily detected (Houterman et al., 2009; Ma et al., 2013). *I-2* encodes a coiled coil (CC), nucleotide-binding (NB), ARC domain (for Apaf1, R proteins, and CED4) and leucine-rich repeat (LRR) protein (Simons et al., 1998). *Avr2* is recognized inside plant cells by *I-2* and a nuclear localization is required to trigger *I-2*-dependent cell death (Ma et al., 2013) (Figure 1). *Avr2* is found in all *Fol* isolates. However, race 3 isolates carry specific point mutations in *Avr2* or a deletion of one triplet (*Avr2*^{V41M}, *Avr2*^{R45H}, *Avr2*^{R46P} and *Avr2*^{T50-}, respectively) that allows *Fol* to overcome *I-2* mediated resistance (Houterman et al., 2009; Chellappan et al., 2016). Interestingly, the point mutations neither the deletion in the race 3 *Avr2* variants do negatively affect virulence of race 3 *Fol* strains, which shows that avirulence of *Avr2* can be uncoupled from its virulence function (Houterman et al., 2009; Chellappan et al., 2016).




Figure 1. A schematic representation of the “gene-for-gene” resistance in the *Fusarium*-tomato interaction. *Fol* secretes *Avr2* into the xylem vessels and apoplastic spaces of tomato plants. Subsequently, it is translocated into the cell by a hitherto unknown mechanism. *Avr2* acts inside the plant cell to exert its virulence function. In a resistant plant, *I-2* containing plant immune responses are triggered upon recognition of nuclear-localized *Avr2* by *I-2*.

I-2 is specifically expressed in the parenchyma cells adjacent to the xylem vessels (Mes et al., 2000). Currently it is unknown how *Avr2* present in the xylem sap is translocated into *I-2* expressing plant cells allowing its perception by the host. *I-2* mediated resistance does not lead to a typical HR response in infected roots, as cell death is not induced (Beckman, 2000). Nevertheless, *I-2* can induce cell death upon the systemic expression of *Avr2* using a Potato virus X (PVX)-based expression system or when *Avr2* and *I-2* are transiently co-expressed in leaves of the non-host *Nicotiana benthamiana* (Houterman et al., 2008; Houterman et al., 2009; Ma et al., 2013). The well-characterized *Avr2-I-2* gene pair makes the *Fol*-tomato interaction a perfect model to study the molecular basis of disease and resistance to wilt diseases.

Outline of the thesis

In **Chapter 2**, the current knowledge on the role of the major phytohormones including SA, JA, ET, ABA and auxin on the interaction between *Fo* and its diverse hosts is summarized. We discuss how phytohormones determine the *Fo* lifestyle, and how phytohormones and *Fo* effectors together act to control the balance between a beneficial and a pathogenic interaction and *vice versa*. **Chapter 3** investigated the role of SA, JA and ET in tomato susceptibility to *FoI*. Thereto, mutants affected in the biosynthesis and perception of SA, ET and JA were inoculated with wild-type *FoI* or a less pathogenic mutant derived from *FoI007* in which the *Avr2* gene was deleted (*FoIΔAvr2*). Finally, a model for the role of SA, ET and JA signaling in the susceptibility of tomato against *FoI* is presented and compared to the involvement of these hormones in the susceptibility of Arabidopsis plant to *Focn*. In **Chapter 4** the virulence function of *Avr2* was investigated. To identify whether *Avr2* acts inside or outside host cells, transgenic tomato plants stably expressing full-length *Avr2* or a cytosolic *Avr2* (Δ *spAvr2*) variant that lacks the signal peptide for secretion were generated. The observation that pathogenicity of an *Avr2* knockout *Fusarium* (*FoIΔAvr2*) strain was fully complemented in Δ *spAvr2* transgenic tomato lines suggests that *Avr2* exerts its virulence functions inside host cells. This hypothesis is in line with the observation that *Avr2* recognition by *I-2* occurs inside plant cells (Ma et al., 2013). Surprisingly, plant-produced full length *Avr2* can also fully complement *FoIΔAvr2* virulence. Western blot analysis revealed that *Avr2*, in contrast to Δ *spAvr2*, was readily detected in the xylem sap and apoplastic fluids, implying that *Avr2* is taken up by plant cells from the extracellular spaces. Additional experiments using other pathogens revealed that *Avr2* uptake is not a host autonomous event and relies on the presence of a pathogen. In **Chapter 5** it is shown that tomato plants expressing Δ *spAvr2* become hyper-susceptible towards various plant pathogens, such as *Fusarium oxysporum*, *Verticillium dahliae* and *Pseudomonas syringae*. This observation prompted us to examine whether *Avr2* specifically interferes with PTI responses. Moreover, to understand how *Avr2* functions at a molecular level its crystal structure was solved in

collaboration with M. Banfield at the John Innes Centre, UK. The structural homology that Avr2 shares with some proteins with known activities allowed prediction of residues that might be involved in the interaction of Avr2 with partner proteins. Two of these putative important residues were found to be required for virulence activity of the protein and enabled us to uncouple virulence from avirulence function. The results described in this thesis are summarized and discussed in **Chapter 6**.

References

- Agrios, G. N. (2005). Plant pathology. St. Louis, MO: Academic Press.
- Alabouvette, C., Olivain, C., Migheli, Q., and Steinberg, C. (2009). Microbiological control of soil-borne phytopathogenic fungi with special emphasis on wilt-inducing *Fusarium oxysporum*. *New Phytol* 184, 529-544. doi: 10.1111/j.1469-8137.2009.03014.x.
- Armstrong, G. M., and Armstrong, J. K. (1981). *Formae speciales and races of Fusarium oxysporum causing wilt diseases*. Pennsylvania State University Press In: Nelson, P.E., Toussoun, T.A. & Cook, R.J. (Eds), 391-399.
- Bari, R., and Jones, J. D. (2009). Role of plant hormones in plant defence responses. *Plant Mol Biol* 69, 473-488. doi: 10.1007/s11103-008-9435-0.
- Beckman, C. H. (2000). Phenolic-storing cells: keys to programmed cell death and periderm formation in wilt disease resistance and in general defence responses in plants? *Physiol Mol Plant P* 57, 101-110. doi: 10.1006/pmpp.2000.0287.
- Berrocal-Lobo, M., and Molina, A. (2004). Ethylene response factor 1 mediates Arabidopsis resistance to the soilborne fungus *Fusarium oxysporum*. *Mol Plant Microbe In* 17, 763-770. doi: Doi 10.1094/Mpmi.2004.17.7.763.
- Berrocal-Lobo, M., and Molina, A. (2008). Arabidopsis defense response against *Fusarium oxysporum*. *Trends Plant Sci* 13, 145-150. doi: 10.1016/j.tplants.2007.12.004.
- Catanzariti, A. M., Do, H. T., Bru, P., de Sain, M., Thatcher, L. F., Rep, M., et al. (2016). The tomato *I* gene for *Fusarium* wilt resistance encodes an atypical leucine-rich repeat receptor-like protein whose function is nevertheless dependent on SOBIR1 and SERK3/BAK1. *Plant J*. doi: 10.1111/tpj.13458.
- Catanzariti, A. M., Lim, G. T., and Jones, D. A. (2015). The tomato *I-3* gene: a novel gene for resistance to *Fusarium* wilt disease. *New Phytol* 207, 106-118. doi: 10.1111/nph.13348.
- Chellappan, B. V., Fokkens, L., Houterman, P. M., Rep, M., and Cornelissen, B. J. C. (2016). Multiple evolutionary trajectories have led to the emergence of races in *Fusarium oxysporum* f. sp. *lycopersici*. *Appl Environ Microbiol*. doi: 10.1128/AEM.02548-16.
- Coll, N. S., Epple, P., and Dangl, J. L. (2011). Programmed cell death in the plant immune system. *Cell Death Differ* 18, 1247-1256. doi: 10.1038/cdd.2011.37.
- de Wit, P. J. G. M. (2007). Visions & reflections (minireview) - How plants recognize pathogens and defend themselves. *Cell Mol Life Sci* 64, 2726-2732. doi: 10.1007/s00018-007-7284-7.
- Demers, J. E., Gugino, B. K., and Jimenez-Gasco, M. D. (2015). Highly diverse endophytic and soil *Fusarium oxysporum* populations associated with field-grown tomato plants. *Appl Environ Microbiol* 81, 81-90. doi: 10.1128/Aem.02590-14.
- di Pietro, A., Madrid, M. P., Caracuel, Z., Delgado-Jarana, J., and Roncero, M. I. (2003). *Fusarium oxysporum*: exploring the molecular arsenal of a vascular wilt fungus. *Mol Plant Pathol* 4, 315-325. doi: 10.1046/j.1364-3703.2003.00180.x.
- Diener, A. C., and Ausubel, F. M. (2005). Resistance to *Fusarium oxysporum* 1, a dominant Arabidopsis disease-resistance gene, is not race specific. *Genetics* 171, 305-321. doi: Doi 10.1534/Genetics.105.042218.
- Dodds, P. N., and Rathjen, J. P. (2010). Plant immunity: towards an integrated view of plant-pathogen interactions. *Nat Rev Genet* 11, 539-548. doi: 10.1038/nrg2812.
- Edel-Hermann, V., Gautheron, N., Mounier, A., and Steinberg, C. (2015). *Fusarium* diversity in soil using a specific molecular approach and a cultural approach. *Journal of Microbiological Methods* 111, 64-71. doi: 10.1016/j.mimet.2015.01.026.

- Fraser-Smith, S., Czişlowski, E., Meldrum, R. A., Zander, M., O'Neill, W., Balali, G. R., et al. (2014). Sequence variation in the putative effector gene *SIX8* facilitates molecular differentiation of *Fusarium oxysporum* f. sp. *cubense*. *Plant Pathol* 63, 1044-1052. doi: 10.1111/ppa.12184.
- Gawehns, F., Houterman, P. M., Ichou, F. A., Michielse, C. B., Hijdra, M., Cornelissen, B. J. C., et al. (2014). The *Fusarium oxysporum* effector Six6 contributes to virulence and suppresses I-2-mediated cell death. *Mol Plant Microbe In* 27, 336-348. doi: 10.1094/MPMI-11-13-0330-R.
- Grant, M. R., and Jones, J. D. G. (2009). Hormone (dis)harmony moulds plant health and disease. *Science* 324, 750-752. doi: 10.1126/science.1173771.
- Hofius, D., Munch, D., Bressendorff, S., Mundy, J., and Petersen, M. (2011). Role of autophagy in disease resistance and hypersensitive response-associated cell death. *Cell Death Differ* 18, 1257-1262. doi: 10.1038/cdd.2011.43.
- Horbach, R., Navarro-Quesada, A. R., Knogge, W., and Deising, H. B. (2011). When and how to kill a plant cell: Infection strategies of plant pathogenic fungi. *J Plant Physiol* 168, 51-62. doi: 10.1016/j.jplph.2010.06.014.
- Houterman, P. M., Cornelissen, B. J. C., and Rep, M. (2008). Suppression of plant resistance gene-based immunity by a fungal effector. *PLoS Pathog* 4, e1000061. doi: 10.1371/journal.ppat.1000061.
- Houterman, P. M., Ma, L., van Ooijen, G., de Vroomen, M. J., Cornelissen, B. J. C., Takken, F. L. W., et al. (2009). The effector protein Avr2 of the xylem-colonizing fungus *Fusarium oxysporum* activates the tomato resistance protein I-2 intracellularly. *Plant J* 58, 970-978. doi: 10.1111/j.1365-313X.2009.03838.x.
- Houterman, P. M., Speijer, D., Dekker, H. L., C.G., d. K., Cornelissen, B. J. C., and Rep, M. (2007). The mixed xylem sap proteome of *Fusarium oxysporum*-infected tomato plants. *Mol Plant Pathol* 8, 215-221. doi: 10.1111/j.1364-3703.2007.00384.x.
- Huang, C. C., and Lindhout, P. (1997). Screening for resistance in wild *Lycopersicon* species to *Fusarium oxysporum* f. sp. *lycopersici* race 1 and race 2. *Euphytica* 93, 145-153. doi: Doi 10.1023/A:1002943805229.
- Imazaki, I., and Kadota, I. (2015). Molecular phylogeny and diversity of *Fusarium* endophytes isolated from tomato stems. *Fems Microbiology Ecology* 91, fiv098. doi: 10.1093/femsec/fiv098.
- Jones, J. D., and Dangl, J. L. (2006). The plant immune system. *Nature* 444, 323-329. doi: 10.1038/nature05286.
- Katan, T., and Di Primo, P. (1999). Current status of vegetative compatibility groups in *Fusarium oxysporum*: Supplement (1999). *Phytoparasitica* 27, 273-277. doi: Doi 10.1007/Bf02981483.
- Ma, L., Cornelissen, B. J. C., and Takken, F. L. W. (2013). A nuclear localization for Avr2 from *Fusarium oxysporum* is required to activate the tomato resistance protein I-2. *Front Plant Sci* 4, 94. doi: 10.3389/fpls.2013.00094.
- Ma, L., Houterman, P. M., Gawehns, F., Cao, L., Sillo, F., Richter, H., et al. (2015). The AVR2-SIX5 gene pair is required to activate I-2-mediated immunity in tomato. *New Phytol* 208, 507-518. doi: 10.1111/nph.13455.
- Mes, J. J., van Doorn, A. A., Wijbrandi, J., Simons, G., Cornelissen, B. J., and Haring, M. A. (2000). Expression of the *Fusarium* resistance gene I-2 colocalizes with the site of fungal containment. *Plant J* 23, 183-193.
- Michielse, C. B., and Rep, M. (2009). Pathogen profile update: *Fusarium oxysporum*. *Mol. Plant Pathol.* 10, 311-324.
- Mur, L. A. J., Kenton, P., Lloyd, A. J., Ougham, H., and Prats, E. (2008). The hypersensitive response; the centenary is upon us but how much do we know? *J Exp Bot* 59, 501-520. doi: 10.1093/

- jxb/erm239.
- Nurnberger, T., Brunner, F., Kemmerling, B., and Piater, L. (2004). Innate immunity in plants and animals: striking similarities and obvious differences. *Immunol Rev* 198, 249-266.
- Pantelides, I. S., Tjamos, S. E., Pappa, S., Kargakis, M., and Paplomatas, E. J. (2013). The ethylene receptor ETR1 is required for *Fusarium oxysporum* pathogenicity. *Plant Pathol* 62, 1302-1309. doi: 10.1111/ppa.12042.
- Pieterse, C. M. J., Leon-Reyes, A., Van der Ent, S., and Van Wees, S. C. M. (2009). Networking by small-molecule hormones in plant immunity. *Nat Chem Biol* 5, 308-316. doi: 10.1038/nchembio.164.
- Pieterse, C. M. J., Van der Does, D., Zamioudis, C., Leon-Reyes, A., and Van Wees, S. C. M. (2012). Hormonal modulation of plant immunity. *Annu Rev Cell Dev Bi* 28, 489-521. doi: 10.1146/annurev-cellbio-092910-154055.
- Rep, M. (2005). Small proteins of plant-pathogenic fungi secreted during host colonization. *FEMS Microbiol. Lett.* 253, 19-27.
- Rep, M., Meijer, M., Houterman, P. M., van der Does, H. C., and Cornelissen, B. J. C. (2005). *Fusarium oxysporum* evades *I*-3-mediated resistance without altering the matching avirulence gene. *Mol Plant Microbe In* 18, 15-23. doi: 10.1094/MPMI-18-0015.
- Rep, M., van der Does, H. C., Meijer, M., van Wijk, R., Houterman, P. M., Dekker, H. L., et al. (2004). A small, cysteine-rich protein secreted by *Fusarium oxysporum* during colonization of xylem vessels is required for *I*-3-mediated resistance in tomato. *Mol Microbiol* 53, 1373-1383. doi: 10.1111/j.1365-2958.2004.04177.x.
- Robert-Seilaniantz, A., Grant, M., and Jones, J. D. G. (2011). Hormone crosstalk in plant disease and defense: more than just jasmonate-salicylate antagonism. *Annual Review of Phytopathology*, Vol 49 49, 317-343. doi: 10.1146/annurev-phyto-073009-114447.
- Schmidt, S. M., Houterman, P. M., Schreiver, I., Ma, L., Amyotte, S., Chellappan, B., et al. (2013). MITEs in the promoters of effector genes allow prediction of novel virulence genes in *Fusarium oxysporum*. *Bmc Genomics* 14, 119. doi: 10.1186/1471-2164-14-119.
- Simons, G., Groenendijk, J., Wijbrandi, J., Reijans, M., Groenen, J., Diergaarde, P., et al. (1998). Dissection of the *Fusarium I2* gene cluster in tomato reveals six homologs and one active gene copy. *Plant Cell* 10, 1055-1068.
- Spoel, S. H., and Dong, X. N. (2012). How do plants achieve immunity? Defence without specialized immune cells. *Nat Rev Immunol* 12, 89-100. doi: 10.1038/nri3141.
- Swarupa, V., Ravishankar, K. V., and Rekha, A. (2014). Plant defense response against *Fusarium oxysporum* and strategies to develop tolerant genotypes in banana. *Planta* 239, 735-751. doi: 10.1007/s00425-013-2024-8.
- Takken, F. L. W., and Rep, M. (2010). The arms race between tomato and *Fusarium oxysporum*. *Mol Plant Pathol* 11, 309-314. doi: 10.1111/j.1364-3703.2009.00605.x.
- Thatcher, L. F., Manners, J. M., and Kazan, K. (2009). *Fusarium oxysporum* hijacks COI1-mediated jasmonate signaling to promote disease development in Arabidopsis. *Plant Journal* 58, 927-939. doi: 10.1111/j.1365-313X.2009.03831.x.
- Tjamos, E. C., and Beckman, C. H. (1989). Vascular wilt diseases of plants. in: Springer-Verlag, Berlin.
- Trusov, Y., Sewelam, N., Rookes, J. E., Kunkel, M., Nowak, E., Schenk, P. M., et al. (2009). Heterotrimeric G proteins-mediated resistance to necrotrophic pathogens includes mechanisms independent of salicylic acid-, jasmonic acid/ethylene- and abscisic acid-mediated defense signaling. *Plant J* 58, 69-81. doi: 10.1111/j.1365-313X.2008.03755.x.
- van der Does, H. C., Lievens, B., Claes, L., Houterman, P. M., Cornelissen, B. J. C., and Rep, M.

- (2008). The presence of a virulence locus discriminates *Fusarium oxysporum* isolates causing tomato wilt from other isolates. *Environ Microbiol* 10, 1475-1485. doi: 10.1111/j.1462-2920.2007.01561.x.
- Zipfel, C., and Felix, G. (2005). Plants and animals: a different taste for microbes? *Curr Opin Plant Biol* 8, 353-360. doi: 10.1016/j.pbi.2005.05.004.