



UvA-DARE (Digital Academic Repository)

Polyamine metabolism and activation of lipid signalling pathways in *Arabidopsis thaliana*

Zarza Cubero, J.

Publication date

2017

Document Version

Other version

License

Other

[Link to publication](#)

Citation for published version (APA):

Zarza Cubero, J. (2017). *Polyamine metabolism and activation of lipid signalling pathways in Arabidopsis thaliana*. [Thesis, fully internal, Universiteit van Amsterdam].

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Chapter 2

***Polyamine Oxidase 5* loss-of-function mutations in *Arabidopsis thaliana* trigger metabolic and transcriptional reprogramming and promote salt stress tolerance**

Xavier Zarza^{1,2}, Kostadin E Atanasov¹, Francisco Marco³, Vicent Arbona⁴, Pedro Carrasco⁵, Joachim Kopka⁶, Vasileios Fotopoulos⁷, Teun Munnik², Aurelio-Gómez Cadenas⁴, Antonio F. Tiburcio¹ and Rubén Alcázar¹

¹Department of Natural Products, Plant Biology and Soil Science, Laboratory of Plant Physiology, Faculty of Pharmacy, University of Barcelona, Barcelona, Spain.

²Department of Plant Physiology, Swammerdam Institute for Life Sciences, University of Amsterdam, Amsterdam, The Netherlands.

³Departamento de Biología Vegetal, Universidad de Valencia, Facultad de Farmacia, Burjassot, Spain.

⁴Departamento de Ciencias Agrarias y del Medio Natural, Universitat Jaume I, Campus Riu Sec, E- 12071, Castelló de la Plana, Spain.

⁵Departamento de Bioquímica y Biología Molecular, Universidad de Valencia, Facultad de Ciencias Biológicas, Burjassot, Spain.

⁶Max Planck Institute for Molecular Plant Physiology, Golm, Germany.

⁷Department of Agricultural Sciences, Biotechnology and Food Science, Cyprus University of Technology, P.O. Box 50329, Limassol, Cyprus.

Published as:

Zarza X, Atanasov KE, Marco F, Arbona V, Carrasco P, Kopka J, Fotopoulos V, Munnik T, Gómez-Cadenas A, Tiburcio AF, and Alcázar R (2016) *Polyamine Oxidase 5* loss-of-function mutations in *Arabidopsis thaliana* trigger metabolic and transcriptional reprogramming and promote salt stress tolerance. *Plant Cell Environ* doi: 10.1111/pce.12714.

ABSTRACT

The family of polyamine oxidases (PAO) in *Arabidopsis* (*AtPAO1-5*) mediates polyamine (PA) back-conversion, which reverses the PA biosynthetic pathway from spermine, and its structural isomer thermospermine (tSpm), into spermidine and then putrescine. Here, we have studied the involvement of PA back-conversion in *Arabidopsis* salinity tolerance. *AtPAO5* is the *Arabidopsis* PAO gene member most transcriptionally induced by salt stress. Two independent loss-of-function mutants (*atpao5-2* and *atpao5-3*) were found to exhibit constitutively higher tSpm levels, with associated increased salt tolerance. Using global transcriptional and metabolomic analyses, the underlying mechanisms were studied. Stimulation of abscisic acid and jasmonate (JA) biosynthesis and accumulation of important compatible solutes, such as sugars, polyols and proline, as well as TCA cycle intermediates were observed in *atpao5* mutants under salt stress. Expression analyses indicate that tSpm modulates the transcript levels of several target genes, including many involved in the biosynthesis and signaling of JA, some of which are already known to promote salinity tolerance. Transcriptional modulation by tSpm is isomer-dependent, thus demonstrating the specificity of this response. Overall, we conclude that tSpm triggers metabolic and transcriptional reprogramming that promotes salt stress tolerance in *Arabidopsis*.

INTRODUCTION

Salinity affects more than 800 million hectares of arable land around the world and it has a detrimental impact on crop yield (Hasegawa 2013). To cope with saline stress, plants have evolved mechanisms that involve discrimination of ions taken up by roots, osmotic adjustment, sequestering of ions into separate cellular compartments, and tolerance against ionic toxicity (Munns & Tester 2008). Polyamines (PAs) are organic amines the levels of which increase in response to various stress conditions, including salinity. Genetic approaches indicate that these compounds are required to cope with abiotic constraints, although their modes-of-action are not fully elucidated (Cona *et al.* 2006; Alcázar *et al.* 2006b; Kusano *et al.* 2008; Alcázar *et al.* 2010; Tiburcio *et al.* 2014). In the glycophyte species, *Arabidopsis thaliana* (hereafter referred to as *Arabidopsis*), the most abundant polyamines (PAs) are the diamine, putrescine (Put), the triamine, spermidine (Spd), and the tetraamines, spermine (Spm) and its structural isomer thermospermine (tSpm). Biosynthesis of PAs requires the presence of arginine or ornithine, from which Put is synthesized. Both amino acids are derived from glutamate, a derivative of the tricarboxylic acid cycle (TCA) intermediate 2-oxoglutarate, which is also precursor of the compatible solute proline (Alcázar *et al.* 2010). Spd is synthesized from Put via co-action of Spd synthase and S-adenosylmethionine decarboxylase. The latter is also involved in the biosynthesis of Spm and tSpm, by Spm synthase (SPMS) and tSpm synthase (*ACAULIS5*, *ACL5*), respectively. Despite their structural similarity, Spm and tSpm isomers exhibit strikingly different biological functions and their biosynthesis is differentially regulated (Takahashi & Kakehi 2010; Vera-Sirera *et al.* 2010; Takano *et al.* 2012).

In *Arabidopsis*, salt stress up-regulates *SPMS* expression (Urano *et al.* 2003; Bagni *et al.* 2006; Tassoni *et al.* 2008; Naka *et al.* 2010). Long-term salt stress in this and other species leads to increased Spm levels (Naka *et al.* 2010; Alet *et al.* 2012). Genetic engineering of the PA pathway and correlational studies using cultivars with contrasting salinity tolerance, suggest that Spm accumulation is a metabolic hallmark often associated with enhanced salt tolerance (Yamaguchi *et al.* 2006). However, sensitivity to salinity of *Arabidopsis* mutants impaired in tSpm biosynthesis (*acl5-1*) or both tSpm and Spm (*acl5-1 spms-1*) is higher than that of Spm deficient (*spms-1*) or wild-type plants (Yamaguchi *et al.* 2006; Alet *et al.* 2012). Therefore, a potential contribution of tSpm to salt stress tolerance could not be ruled out. Evidence indicate that PA-triggered salinity tolerance is independent of the salt overly sensitive (SOS) signalling pathway (Yamaguchi *et al.* 2006; Alet *et al.* 2012) and may involve calcium signalling (Yamaguchi *et al.* 2006), modulation of various cation and K⁺ channels (Pottosin & Shabala 2014), H⁺/ATPase activities (Pottosin *et al.* 2014), protein post-translational modifications such as S-nitrosylation, carbonylation and tyrosine nitration (Tanou *et al.* 2014), enzymatic and non-enzymatic reactive oxygen species (ROS) detoxification (Saha *et al.* 2015), among others (Alcázar *et al.* 2010). However, the

overall picture remains largely undefined and global transcriptional and metabolic approaches are, to our knowledge, not yet reported.

Free PA levels are not only determined by their biosynthesis but also by their oxidation (Cona et al. 2006). In recent years, several genes have been characterized that are involved in PA oxidation (Moschou et al. 2012; Tavladoraki et al. 2012). PA oxidation requires the activity of copper-containing amine oxidases (CuAOs) or FAD-containing polyamine oxidases (PAOs). Both types of amine oxidases exhibit different substrate preferences. While Put is preferentially oxidized by CuAOs, higher molecular weight PAs (Spd, Spm and tSpm) are PAO substrates (Cona et al. 2006). PAOs are classified into two main families, depending on whether they terminally oxidize PAs or catalyse the PA back-conversion, which reverses the PA biosynthetic pathway releasing H₂O₂ (Moschou et al. 2012; Tavladoraki et al. 2012). *Arabidopsis* carries five *AtPAO* genes (*AtPAO1-5*), all known to be involved in PA back-conversion (Tavladoraki et al. 2006; Kamada-Nobusada et al. 2008; Moschou et al. 2008c; Takahashi et al. 2010; Fincato et al. 2011; Ahou et al. 2014; Kim et al. 2014). Among them, *AtPAO5* encodes a cytosolic PAO, the expression of which is restricted to vascular tissue, which is also the site of synthesis of tSpm by *ACL5* (Kim et al. 2014). *AtPAO5* preferentially catalyses the conversion of tSpm to Spd, but not of Spd to Put (Ahou et al. 2014; Kim et al. 2014). Consistent with the proposed catalytic activity of *AtPAO5*, *atpao5* mutants accumulate higher levels of tSpm under normal growth conditions (Ahou et al. 2014; Kim et al. 2014). Therefore, *AtPAO5* and *ACL5* activities can fine-tune the balance between tSpm and Spm.

PA mechanisms of action might involve cross-regulation with different hormones such as gibberellins (Alcázar et al. 2005), ethylene (Bitrián et al. 2012), auxins (Tong et al. 2014) or abscisic acid (ABA) (Alcázar et al. 2006a; Cuevas et al. 2008). Recently, an apoplastic CuAO (*AtAO1*) that is expressed early during root vascular tissue differentiation has been shown to mediate jasmonic acid-triggered early protoxylem differentiation (Ghuge et al. 2015). In addition to the involvement of jasmonates (JAs) in plant development, wounding and biotic stress interactions, recent evidence indicate that JA contributes to salt stress tolerance (Kazan 2015). Overexpression of the JA biosynthesis gene *allene oxide cyclase 1* (*AOC1*) from wheat (*Triticum aestivum*) in *Arabidopsis* leads to higher JA levels, and this is associated with increased salt tolerance independent of ABA and dependent of *MYC2* (Zhao et al. 2014b). Furthermore, overexpression of the wheat 12-oxophytodienoate reductase 1 in *Arabidopsis* promotes ABA-dependent salt tolerance and enhanced ROS detoxification (Dong et al. 2013). Hence, JA triggers both ABA dependent and independent salinity tolerance in this species.

Overexpression of PA biosynthetic genes has been successfully applied for the manipulation of PA levels in plants (Alcázar et al. 2006b). However, isolation of PA catabolism mutants emerges as an alternative approach for the accumulation of PAs. It is expected that such approaches should help to further define PA-related

biological functions. In this work, we have studied the involvement of PA back-conversion during salt stress in *Arabidopsis*. Expression analyses indicate that *AtPAO5* is the *Arabidopsis* PAO gene family member most transcriptionally responsive to salinity. Phenotypic characterization of two *atpao5* mutant alleles evidences that *AtPAO5* loss of function leads to salinity tolerance. Through global transcriptional and metabolic analyses, we identify that salt tolerance in *atpao5* associates with early boosting of ABA biosynthesis and higher JA levels, followed by the accumulation of important compatible solutes, TCA cycle intermediates, and transcriptional modulation of several target genes by tSpm. Collectively, the results indicate that tSpm has a protective role against salt stress in *Arabidopsis*.

RESULTS

Expression of *AtPAO1-5* gene family members during salt stress

To identify which *AtPAO* gene family members are transcriptionally responsive to salinity, we determined the expression of all five (*AtPAO1-5*) members by qRT-PCR in 4-week-old wild-type *Arabidopsis* plants treated with 50 mM NaCl between 0 and 96 h (**Fig. 1**). The expression of most *AtPAO* genes (*AtPAO1*, *AtPAO3*, *AtPAO4*) remained unaltered by the saline treatment. In contrast, the expression of *AtPAO5* progressively increased up to 4.5-fold after 96 h of salt treatment. These analyses also revealed an early, but transient, peak of *AtPAO2* expression after 3 h (**Fig. 1**). We concluded that *AtPAO5* is the *PAO* family member that is most transcriptionally responsive to saline conditions in *Arabidopsis*, which might represent its potential involvement during salt stress.




Figure 1. Transcript profiles of *AtPAO* gene family members during salt stress. qRT-PCR expression analyses of *AtPAO1–5* genes in 4-week-old *Arabidopsis* wild-type and *atpao5-3* loss-of-function mutants exposed to 50 mM NaCl for 0–96 h. Values are relative to wild type (0 h) and are the mean \pm SD of three biological replicates, each with three technical replicates. nd, not detectable. Letters indicate values that are significantly different according to Student–Newman–Keuls test at P value < 0.05.

Salt tolerance phenotype of *atpao5-2* and *atpao5-3* loss-of-function mutants

To investigate the biological significance of the transcriptional activation of *AtPAO5* by salt stress, two *atpao5* T-DNA insertion mutants were isolated: *atpao5-2* (Kim et al. 2014) and *atpao5-3* (Fig. S1a). *AtPAO5* expression was undetectable in both *atpao5-2* and *atpao5-3*, indicating that they are loss-of-function mutants (Fig. S1b). Loss of *AtPAO5* expression in *atpao5-3* did not lead to consistent changes in the expression of *AtPAO1–4* genes during salt stress (Fig. 1). These results point to the absence of transcriptional compensatory mechanisms between *AtPAO5* and *AtPAO1–4* gene family members during the saline treatment.

The salt-tolerant phenotype of *atpao5-2*, *atpao5-3* and wild-type plants was investigated. Four-week-old plants grown in the absence of stress were exposed to gradual increases of NaCl (50, 100, 150, 200 and 250 mM NaCl; refer to Materials and methods). *atpao5-2* and *atpao5-3* mutants maintained higher percentage of water content and chlorophyll levels than wild-type plants, consistent with a better performance under salt stress conditions (**Fig. 2a-c**). Salt stress tolerance was also phenotypically evident in *atpao5-2* and *atpao5-3* seedlings (**Fig. 2d,e**). We concluded that *AtPAO5* loss of function leads to enhanced salt tolerance in *Arabidopsis*.




Figure 2. Salt stress tolerance of *atpao5* compared with wild-type plants. (a) Phenotype of *atpao5-2*, *atpao5-3* and wild-type plants after 96 h of 200 mM NaCl treatment with gradual increases of salt (refer to Materials and methods). (b) Percentage of water content and (c) chlorophyll levels in *atpao5-2*, *atpao5-3* and wild-type plants exposed to gradual increases of salt (+NaCl) and untreated controls (-NaCl). (d) Phenotype and (e) chlorophyll levels of *atpao5-2*, *atpao5-3* and wild-type seedlings exposed to gradual increases of salt (refer to Materials and methods). Letters indicate values that are significantly different according to Student-Newman-Keuls test at *P* value <0.05.

PA levels in *atpao5* loss-of-function mutants during salt stress

Free Put, Spd, Spm and tSpm levels were quantified in *atpao5-2*, *atpao5-3* and wild-type plants exposed to 0, 50, 100 and 150 mM NaCl for 48 h (**Fig. 3**). In the absence of stress (0 mM NaCl), PA levels remained constant over the time of analysis (**Fig. 3**). Put levels significantly increased in the wild type and both *atpao5-2* and *atpao5-3* mutant alleles exposed to 150 mM NaCl, while less or no Put increases were observed at lower NaCl concentrations (50 and 100 mM NaCl, **Fig. 3**). Conversely, Spd levels declined in response to 100 and 150 mM NaCl in the wild type and both *atpao5* mutants (**Fig. 3**). No remarkable differences in the levels of Put, Spd or Spm were observed between *atpao5* mutants and the wild type (**Fig. 3**). However, the levels of tSpm were consistently higher in *atpao5-2* and *atpao5-3* than the wild type and remained unaltered during the saline treatment (**Fig. 3**). These results are in agreement with previous observations that tSpm is preferential substrate for AtPAO5 activity (Ahou *et al.* 2014; Kim *et al.* 2014). Because both *atpao5-2* and *atpao5-3* mutants exhibited identical salt tolerance phenotypes and PA profiles (**Figs. 2 & 3**), *atpao5-3* was selected for further characterization.




Figure 3. Free Put, Spd, Spm and tSpm levels in 4-week-old wild-type, *atpao5-2* and *atpao5-3* mutants treated with 0, 50, 100 and 150 mM NaCl between 0 and 48 h. Values are the mean \pm SD of three biological replicates, each with three technical replicates. FW, fresh weight. Letters indicate values that are significantly different according to Student–Newman–Keuls test at P value < 0.05 .

Sodium ion levels in *atpao5-3* during salt stress

Quantification of Na^+ in leaves of *atpao5-3* and wild type during salt stress evidenced the occurrence of lower shoot sodium accumulation in *atpao5-3* (**Fig. S2**). This observation suggested that Na^+ exclusion might contribute to the salt-tolerant phenotype of *atpao5*, which is in line with a potential role for PAs in regulating various ion channels (Pottosin & Shabala 2014). However, because shoot Na^+ content in *atpao5-3* almost reached wild-type levels after 96 h of 150 mM NaCl treatment, without evident symptoms of ion toxicity (**Figs. 2 & S2**), we reasoned that other mechanisms, in addition to transient Na^+ exclusion, might contribute to the salt tolerance phenotype induced by *AtPAO5* loss of function.

Quantification of ABA in *atpao5-3* during salt stress

ABA levels were determined during early (osmotic) and late (ionic, approximately 24 to 48 h after 100 mM NaCl) phases of salt treatment (**Fig. 4a**) (Munns & Tester 2008). Under basal conditions, no differences in ABA contents were observed between *atpao5-3* and wild-type plants. Treatment with 50 mM NaCl induced ABA accumulation in both genotypes after 3 h of treatment. However, ABA levels in *atpao5-3* were moderately but significantly higher than the wild type up to 48 h of salt treatment (**Fig. 4a**). Quantitative expression analyses of the ABA biosynthesis gene *NCED3* and ABA-inducible gene *RD29B* (Uno *et al.* 2000) in *atpao5-3* and wild type correlated with ABA levels (**Fig. 4b**). Conversely, *atpao5-3* accumulated less ABA than the wild type at later stages of salt treatment (**Fig. 4a**). This observation is consistent with the salinity tolerance in *atpao5-3* that is associated with delayed sodium accumulation in leaves (**Fig. S2**). We concluded that, although ABA biosynthesis is moderately promoted in *atpao5-3* during the osmotic phase of salt stress, this is translated into higher ABA-dependent *RD29B* gene expression compared with the wild type.




Figure 4. (a) Determination of ABA levels in *atpao5-3* and wild-type plants exposed to gradual increases of NaCl between 50 to 150 mM. Values are the mean \pm SD of five technical replicates. The analysis was repeated in two additional independent experiments showing similar results. (b) Quantitation of *NCED3* and *RD29B* expression in *atpao5-3* and wild-type plants exposed to 50 mM NaCl between 0 and 96 h. Values are the mean \pm SD from three biological replicates, each with three technical replicates. Letters indicate values that are significantly different according to Student-Newman-Keuls test at P value <0.05 .

Hydrogen peroxide levels in *atpao5-3* during salt stress

To determine the ROS detoxification capacity, hydrogen peroxide (H_2O_2) levels in *atpao5-3* and wild-type plants were quantified during salt stress (**Fig. S3**). Interestingly, H_2O_2 levels did not differ between both genotypes and accumulated to similar levels throughout the saline treatment (**Fig. S3**). Major increases in hydrogen peroxide were observed during the ionic phase of salt stress in both the wild type and *atpao5-3* (**Fig. S3**). These results suggest that the salt tolerance of the *atpao5-3* mutant is not caused by enhanced ROS detoxification capacity. It also demonstrates that contribution of AtPAO5 to the overall generation of H_2O_2 during salt stress in *Arabidopsis* is rather limited.

Transcriptome analyses of *atpao5-3* during salt stress

To get further insight into mechanisms underlying *atpao5-3* salt tolerance, we determined the genome wide-expression profiles of *atpao5-3* and wild-type plants before and after 3 h of 50 mM NaCl treatment using Affymetrix microarrays. Genes that exhibited significant expression differences ($P < 0.05$; fold change ≥ 2) were detected when comparing genotypes (*atpao5-3* 0 h versus wild-type 0 h, referred to as '*p0w0*'; *atpao5-3* 3 h versus wild-type 3 h, '*p3w3*') and treatments (wild-type 3 h versus wild-type 0 h, '*w3w0*'; *atpao5-3* 3 h versus *atpao5-3* 0 h, '*p3p0*') (**Fig. 5a**; **Tables S2–S15**).

In the absence of stress, already 201 genes were differentially expressed between *atpao5-3* and the wild type (*p0w0*, **Table S4**). GO analyses identified a significant number of genes related to signaling (**Fig. 5b**), which were mainly related to JA, protein kinases and transcription factors (**Table S4**). Interestingly, *atpao5-3* exhibited higher basal expression of JA metabolism genes *lipoxygenase 4 (LOX4)*, *AOC1*, *jasmonoyl-isoleucine-12-hydroxylase (CYP94B3)*, *jasmonate ZIM-domain 1 (JAZ1)*, *JAZ5*, *JAZ6*, *JAZ7* transcriptional repressors and the JAZ-target transcription factor *MYC4* (**Table S16**). Overall, these results suggest that *atpao5-3* has increased basal expression of several JA biosynthesis and signalling genes. Remarkably, most JA genes up-regulated in *atpao5-3* were also induced by salt stress in the wild type (*w3w0* in **Table S16**).




Figure 5. (a) Venn diagrams and (b) heat map of gene ontology terms from differentially expressed genes in *atpao5-3* and wild-type plants treated with 50 mM NaCl between 0 and 3 h. Genes in the different comparisons (*w3w0*, wild-type 3 h versus wild-type 0 h of 50 mM NaCl treatment; *p3p0*, *atpao5-3* 3 h versus *atpao5-3* 0 h of 50 mM NaCl treatment; *p0w0*, *atpao5-3* 0 h versus wild-type 0 h; and *p3w3*, *atpao5-3* 3 h versus wild-type 3 h) are listed in **Tables S2–15**.

In the comparison between treatments (*p3p0* and *w3w0*), 805 differentially expressed genes in *p3p0* were detected and 622 in *w3w0*, of which 356 were common between *p3p0* and *w3w0* (**Fig. 5a**; **Tables S2, S3 and S10**). Common genes were similarly up- or down-regulated and were mostly related to abiotic stimulus but also response to wounding, further reinforcing a potential role for JA on salt stress (**Table S10**; **Fig. 5b**). In the set of genes whose expression was exclusively changed in *atpao5-3* by 50 mM NaCl treatment (*p3p0*; **Table S6**), a significant enrichment of protein kinases was identified (**Table S6**; **Fig. 5b**). We also found that salt treatment led to increased expression of the Na⁺/pyruvate symporter *bile acid:sodium symporter family protein 2* (*BASS2*, *At2g26900*) and Na⁺/H⁺ antiporter *sodium:hydrogen antiporter 1* (*NHD1*, *At3g19490*) in *atpao5-3*, which are sodium transporters localized to the chloroplast envelope that mediate the plastidial influx or efflux of Na⁺, respectively (Furumoto *et al.* 2011; Müller *et al.* 2014) (**Table**

S6). Enhanced expression of *NHD1* in *atpao5-3* might contribute to its salt tolerance, because *nhd1* mutants exhibit reduced growth, less chlorophyll levels and lower photosynthetic performance because of decreased capacity of sodium export from chloroplasts (Müller *et al.* 2014). Interestingly, exogenous application of 100 μ M tSpm to 7-day-old wild-type seedlings quickly induced the expression of *NHD1*, and this response was not triggered by Spm (**Fig. S4**). This early peak of *NHD1* expression was followed by a 2.3-fold expression peak after 8 h of treatment with tSpm or Spm, which overlapped with *BASS2* expression kinetics (**Fig. S4**). These data suggest that tSpm rapidly induces *NHD1* expression, thus potentially promoting Na^+ efflux from the chloroplast. Conversely, Spm and tSpm exhibit the same effect on *BASS2* transcriptional up-regulation after 8 h of exogenous treatment (**Fig. S4**).

Taken together, the data suggest that *AtPAO5* loss of function impacts cell signalling through the transcriptional modulation of protein phosphorylation. This reinforces the hypothesis of a PA modulon in plants that involves transcriptional regulation of certain target genes by PAs (Tiburcio *et al.* 2014).

Transcriptional regulation of the JA pathway by tSpm JA levels during salt stress.

To investigate the effect of higher PAs on the transcriptional regulation of the JA pathway, we applied 100 μ M Spm or tSpm to 7-day-old wild-type seedlings and monitored the expression kinetics of JA biosynthesis genes (*LOX1*, *LOX2*, *LOX3*, *LOX4*, *AOS*, *AOC1*, *AOC2*, *OPR3*, *OPCL1*, *CYP94B3*; **Fig. 6a**), JA signalling genes (*JAZ1*, *JAZ5*, *JAZ6*, *JAZ7*, *JAZ10*, *MYC2*, *MYC3* and *MYC4*; **Fig. 6b**) and JA response marker genes (*VSP2* and *PDF1.2*; **Fig. 6b**). A significant up-regulation of JA marker gene *JAZ10* was observed by exogenous application of tSpm, whereas Spm induced less, but still significant (≥ 2 -fold), up-regulation. Conversely, *PDF1.2* and *VSP2* expressions were only significantly induced by tSpm and not by Spm (**Fig. 6b**). Exogenously supplied tSpm also increased the expression of 13-lipoxygenases *LOX1-4* (**Fig. 6a**). LOX encode enzymes oxidizing α -linoleic acid to 13(S)-hydroperoxy-octadecatrienoic acid, which is precursor of different oxylipins. tSpm induced the transcriptional up-regulation of *LOX1* and *LOX4* expression, whereas Spm had little or no effect (**Fig. 6a**). The transcript levels of *AOS*, which is the first gene committed to JA biosynthesis, allene oxide cyclases *AOC1* and *AOC2*, OPC-8:CoA ligase 1 (*OPCL1*) and *CYP94B3*, were also increased by tSpm treatment and less by Spm. Conversely, *OPR3* expression was more induced by Spm than tSpm (**Fig. 6a**).




Figure 6. Quantitative real-time gene expression analyses of JA biosynthesis (a) and signalling (b) genes from 7-day-old wild-type seedlings treated with 100 μM tSpm or 100 μM Spm during 0-24 h. Values are the mean from three biological replicates \pm SD, each with three technical replicates. Significant differences in gene expression relative to time 0 h were determined using Student's *t*-test and indicated by asterisks: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.005$.

Consistent with the transcriptional activation of the JA pathway by tSpm, the expression of transcription factors *JAZ1*, *JAZ5*, *JAZ6*, *JAZ7* and *JAZ10* was highly induced by exogenously supplied tSpm and less by Spm (**Fig. 6b**). Among *MYC2*, *MYC3* and *MYC4* gene family members, tSpm significantly induced *MYC4* expression after 24 h of treatment (**Fig. 6b**). Based on DNA binding specificity, *MYC4* has been suggested to have different target genes than *MYC2* (Fernández-Calvo *et al.* 2011). Remarkably, *MYC4*, *ACL5* and *AtPAO5* expression are found in the vasculature, which is the site of tSpm biosynthesis (Fernández-Calvo *et al.* 2011; Kim *et al.* 2014). Dose-response expression kinetics were determined for some tSpm and/or Spm-induced genes after 15 min of treatment, using different concentrations of tSpm and Spm (0, 5, 10, 20 and 100 μM ; **Fig. S5**). Expression up-regulation was already observed using as low as 5 μM tSpm or Spm, thus suggesting the specificity of this

response (**Fig. S5**). Transcriptional responses increased linearly with the concentration of exogenously supplied PAs, with the highest differences observed at 100 μ M tSpm or Spm.

Finally, and consistently with the transcriptional activation of the JA pathway by tSpm, we observed higher levels of JA in *atpao5-3* than in the wild type during the saline treatment (**Fig. 7**). Overall, JA levels increased during saline stress but accumulated to higher levels in *atpao5-3* than the wild type (**Fig. 7**).




Figure 7. Quantification of JA levels in *atpao5-3* and wild-type plants exposed to gradual increases of NaCl between 50 and 150 mM. Values are the mean \pm SD of five technical replicates. The analysis was repeated in two additional independent experiments showing similar results. Letters indicate values that are significantly different according to Student-Newman-Keuls test at P value < 0.05 .

Altogether, these results indicate that tSpm, and to a lower extent Spm, promote the transcriptional activation of genes involved in JA biosynthesis and signalling. Furthermore, these data indicate that transcriptional up-regulation of the JA pathway by PAs is isomer-specific with major contribution by tSpm and minor contribution by Spm.

Global metabolite profiles of *atpao5-3* during salt stress

To get further insight into additional mechanisms underlying *atpao5-3* salt stress tolerance from a metabolic view, we performed global metabolite profiling of *atpao5-3* and wild-type plants during salt stress by GC-MS. Our analyses identified 84 metabolites in 36 samples analysed, which were annotated according to the MPIMP-

Golm inventory list (Kopka *et al.* 2005) (**Table S17**). The chemical nature of 56 of such metabolites was known, and this set was used for PCA. For simplification, we used 0 and 24 h time points in which most metabolic changes were evident (**Fig. 8a**). PC1, which explains 72.8% of total variance, revealed the effect of salt stress on metabolic profiles. PC2 (13.3% of total variance) reflected the differences between genotypes (*atpao5-3* and wild type), which increased after exposure to salt stress (**Fig. 8a**). *atpao5-3* 0 h was shifted to the right in PC1, which suggests that *atpao5-3* loss of function triggers a metabolic response similar to the salt treatment (**Fig. 8a**). Furthermore, metabolic responses to salt in *atpao5-3* were earlier and more intense, consistent with a metabolic pre-acclimation hypothesis (PC1 in Fig. 8a). PC1 and PC2 loadings with highest values corresponded to galactinol and melibiose (**Table S18**). HCA Pearson's correlation and average linkage of metabolites and samples revealed the occurrence of four major metabolite clusters (**Fig. 8b**). Sample groupings evidenced contrasted differences between *atpao5-3* and wild-type plants under salt stress. Heat map representation clearly indicated that some metabolic responses to salt were more intense in *atpao5-3* than the wild type, in line with PCA observations (**Fig. 8b**).




Figure 8. (a) Principal component analysis (PCA) and (b) HCA Pearson's correlation and average linkage of metabolites and samples from *atpao5-3* and wild-type plants exposed to salt stress.

Based on metabolite profiles of the wild type, we defined 'early' and 'late' salt-responsive metabolites. Late salt-responsive metabolites were those that exhibited significant ($P < 0.05$) differences in the wild type after 24 and 48 h of 150 mM NaCl treatment. Early salt-responsive metabolites corresponded to those showing significant ($P < 0.05$) differences in the wild type after 8 and 24 h of 50 mM NaCl. The sum of Put and agmatine levels was the only early salt-responsive metabolite detected, but its levels were not consistently different between the wild type and *atpao5-3* during salt stress (**Fig. S6**). Conversely, 28 late salt-responsive metabolites were detected, of which 22 were increased and 6 reduced by salt treatment (**Tables S19** and **S17**). Eighteen of the 28 late salt-responsive metabolites exhibited significant differences between *atpao5-3* and wild type (**Table S19**). Among these, we identified polyols (galactinol, *myo*-inositol), amino acids (proline, alanine, serine), TCA cycle intermediates (succinate, malate), sugars (lyxose), dehydroascorbic acid and shikimic acid (**Fig. 9** and **Table S17**).




Figure 9. Schematic representation of TCA cycle intermediates and primary metabolites in associated pathways. Log₂ ratios ± SE values are relative to wild-type 0 h. Values are shown for *atpao5-3* (red) and wild-type (blue) 24 h after 50, 100 and 150 mM NaCl treatment. Significant differences between genotypes at the same time point and salt concentration were determined using Student's *t*-test and indicated by asterisks: **P* < 0.05, ***P* < 0.01, ****P* < 0.005.

A number of additional metabolites were exclusively changed in *atpao5-3* by salt stress and not in the wild type. Such 'elicited' metabolites in *atpao5-3* corresponded to amino acids (alanine), sugar alcohols (arabitol, erythritol), additional TCA cycle intermediates (fumarate, citrate), sugars (glucose, sucrose, rhamnose) and ethanolamine, which is precursor of choline and glycine betaine (Chen & Murata 2011) (**Fig. 9 and Tables S20 and S17**). Increased levels of glucose in *atpao5-3* suggested activation of glycolysis that would feed the TCA cycle. Indeed, depletion of starch content and accumulation of soluble sugars seem to be a general response to salt stress (Kempa *et al.* 2008).

Altogether, metabolic profile analyses indicate that *atpao5-3* mutant, which exhibits constitutively higher tSpm levels, responds earlier and more intense to salt stress and accumulates higher levels of important compatible solutes. Overall, we conclude that tSpm triggers metabolic reprogramming in *Arabidopsis* that allows a faster and more intense response to salt stress.

DISCUSSION

In this work, we have studied the potential involvement of the PA back-conversion pathway in *Arabidopsis* salt stress tolerance. *Arabidopsis* carries five PAO genes (*AtPAO1-5*) that encode proteins with distinct subcellular localizations and different substrate preferences. *AtPAO1* and *AtPAO5* are cytosolic enzymes, whereas *AtPAO2-4* are localized in peroxisomes (Møller & McPherson 1998; Cona *et al.* 2006; Reumann *et al.* 2009; Planas-Portell *et al.* 2013). *Arabidopsis* PAOs mediate PA back-conversion, which is coupled to H₂O₂ production. High levels of H₂O₂ generated by apoplastic PAO activity can trigger programmed cell death under salt stress in tobacco (Moschou *et al.* 2008b; Moschou *et al.* 2008a). However, in *Arabidopsis*, most PAOs are peroxisomal and apoplastic oxidation seems to rely on CuAO activity (Planas-Portell *et al.* 2013). Among *AtPAO* gene family members, *AtPAO5* is the most transcriptionally responsive to salt stress (**Fig. 1**). Consistent with a potential role for *AtPAO5* on salt stress, two independent *atpao5* loss-of-function mutants (*atpao5-2* and *atpao5-3*) exhibit better performance under salinity (**Fig. 2**). Our results indicate that *AtPAO5* activity is detrimental for salinity tolerance. Tolerance to abiotic stress has also been generated by antisense down-regulation of PAO expression in tobacco (Moschou *et al.* 2008b). However, *atpao5* tolerance does not correlate with diminished ROS production because the levels of H₂O₂ during salt stress are barely

affected by *atpao5* mutation (**Fig. S3**). Our analyses suggest that tSpm, which levels are constitutively higher in *atpao5* mutants under salt stress (Fig. 3), triggers transcriptional and metabolic reprogramming that is isomer-specific. These results support an intrinsic role for PAs providing salt stress tolerance (Moschou *et al.* 2008b).

Hyperosmolarity is the primary effect of salt stress, causing turgor loss and ABA induction (Hasegawa 2013). To achieve osmotic adjustment, different compounds such as sugars, sugar alcohols, proline and glycine betaine accumulate (Chen & Murata 2002). We observed that ABA biosynthesis is moderately promoted in *atpao5-3* compared with the wild type early during the osmotic phase of salt stress (Munns & Tester 2008), but this is translated into significant differences in the expression of the ABA-inducible *RD29B* gene (**Fig. 4**) (Uno *et al.* 2000). Metabolic profiling by GC-MS of polar metabolites evidenced contrasted differences between *atpao5-3* and wild-type plants during salt stress (**Fig. 8a**). Most metabolic changes were observed after 24 h of NaCl treatments and involved higher levels of sugars, polyols, proline and TCA cycle intermediates in *atpao5-3* compared with the wild type (**Figs. 8b & 9**). Most of such metabolites were also found increased in *Arabidopsis* wild-type plants exposed to 150 mM NaCl between 6 h to 5 d (Kempa *et al.* 2008). The higher levels of some osmoprotective compounds in *atpao5-3* are compatible with early boost of ABA biosynthesis (Kempa *et al.* 2008). Compatible solutes maintain cell turgor by osmotic adjustment, and redox balance by removing excess of ROS, thus contributing to abiotic stress tolerance (Munns & Tester 2008; Székely *et al.* 2008; Krasensky & Jonak 2012). Interactions between PAs and ABA have been described during drought and cold stresses, and lower ABA levels have been shown to underlie the freezing sensitivity of *adc1* mutants (Alcázar *et al.* 2006a; Cuevas *et al.* 2008; Toumi *et al.* 2010). Conversely, ABA levels in *atpao5-3* are lower than the wild type at later stages of the saline treatment, which is consistent with better performance of *atpao5-3* under salt stress that leads to reduced stress perception. Reduction in the levels of organic acids and TCA cycle intermediates are often observed in metabolite profiling of glycophyte species exposed to saline stress (Gong *et al.* 2005; Sánchez *et al.* 2008; Hill *et al.* 2013; Zhao *et al.* 2014a). These observations suggest that energy production and/or plant growth are repressed by salinity (Zhao *et al.* 2014a). In this work, we have observed that *atpao5-3* accumulates higher levels of TCA cycle intermediates during salt stress (**Fig. 9**) and this associates with tolerance. Higher levels of soluble sugars suggest activation of glycolysis that would feed the TCA cycle. Taken together, *atpao5-3* mutation impacts cell metabolism through promotion of the accumulation of stress-protective compounds and TCA cycle intermediates.

Despite salt stress tolerance was observed at later stages of treatment (**Fig. 2**), we determined transcriptional changes early during salt stress (3 h after 50 mM NaCl treatment; **Fig. 5**) to distinguish between direct effects from *atpao5* mutation

from late, possibly indirect, transcriptional changes derived from long-term salt exposure. Microarray expression analyses revealed an enhanced transcriptional steady state of the JA pathway in *atpao5-3* mutants (**Table S16**). Quantitative expression of several JA biosynthesis, signalling and marker genes from wild-type seedlings treated with 100 μ M tSpm or Spm evidenced that this effect is isomer-specific (**Fig. 6**). The expression of most JA-related genes analysed responded to exogenous tSpm within 15 min of treatment, which is consistent with a direct effect of tSpm on transcriptional regulation. In addition, *atpao5-3* exhibited higher JA levels than the wild type during salt stress (**Fig. 7**). JAs are known to accumulate under saline conditions (**Fig. 7**) (Moons *et al.* 1997; Pedranzani *et al.* 2003), and correlations between high JA levels and salinity tolerance have been observed in different rice cultivars (Kang *et al.* 2005). Furthermore, most transcriptionally up-regulated JA genes detected in the microarray were responsive to salt treatment (**Table S16**). Additional recent evidence support that JA are positive regulators of salt tolerance (Dong *et al.* 2013; Zhao *et al.* 2014a, 2014b; Kazan 2015). Based on these evidences, we suggest that tSpm-triggered transcriptional up-regulation of JA biosynthesis and signalling genes, as well as higher JA levels, contribute to the metabolic reprogramming and salt tolerance phenotypes of *atpao5* mutants. We suggest that both ABA and JA pathways are involved in tSpm-triggered salt tolerance in *Arabidopsis*.

Microarray analyses also identified that the expression of *BASS2* and *NHD1* genes were up-regulated by salt stress in *atpao5-3*. *BASS2* encodes a plastid localized Na^+ /pyruvate symporter, the activity of which is coupled with that of the Na^+/H^+ antiporter *NHD1* (Furumoto *et al.* 2011). *BASS2* functions in C4 photosynthesis and provides pyruvate for the plastidic methylerythritol phosphate pathway in C3 plants (Furumoto *et al.* 2011). In *Arabidopsis*, *NHD1* (Na^+/H^+ antiporter 1) mutation leads to higher plastidic Na^+ , which is associated with reduced growth and lower chlorophyll levels (Müller *et al.* 2014). Interestingly, tSpm rapidly induces *NHD1* expression (**Fig. S4**) that may allow higher efflux of chloroplastic Na^+ , thus contributing to enhanced tolerance against salt stress. In addition, PAs with higher net charge have stronger inhibitory effect on the vacuolar non-selective cation channels, hence, potentially assisting vacuolar Na^+ sequestration (Pottosin & Shabala 2014).

Altogether, we propose that mechanisms underlying salt stress tolerance by PAs are diverse and additive and involve transcriptional and metabolic reprogramming by tSpm. This view is consistent with co-evolution of PAs with stress pathways that has allowed efficient integration of environmental signals into transcriptional and metabolic responses. Furthermore, we conclude that tSpm is not only involved in plant development but also plays an important role in abiotic stress tolerance.

MATERIALS AND METHODS

Plant material and growth conditions

Arabidopsis thaliana wild-type (Col-0), *atpao5-2* (N553110) and *atpao5-3* (N509671) T-DNA insertion mutants were obtained from the Nottingham *Arabidopsis* Stock Centre (University of Nottingham, Loughborough, UK). Seeds were sown on a mixture of soil and vermiculite (3:1) and stratified 4 d in the dark at 4 °C to synchronize germination. Plants were grown under 16/8 h light/dark cycles, 100-125 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ of light intensity, 21 ± 2 °C and $70 \pm 10\%$ relative humidity. Salt treatments in adult plants were performed by irrigation of 4-week-old plants grown on soil with incremental concentrations (50, 100 and 150 mM) of NaCl every 4 d (Gaxiola *et al.* 2001). Samples from rosette leaves were harvested at different time points, frozen in liquid nitrogen and stored at -80 °C until further analysis. Salt treatment of *Arabidopsis* seedlings was performed as follows: Different genotypes were germinated on a nylon mesh (\varnothing 43 μm) placed on top of 0.5 \times MS-MES medium (Murashige & Skoog, 0.5 mM MES, 1% sucrose, 1% agar, pH 5.7 with KOH) under sterile conditions. Ten days after germination, the nylon mesh with seedlings was transferred to the same medium supplemented with 50 mM NaCl for 3 d. Seedlings were then transferred to 0.5 \times MS-MES medium supplemented with 225 mM NaCl. Chlorophyll quantification was determined in the different genotypes 5 d after 225 mM NaCl treatment.

Identification of *atpao5-2* and *atpao5-3* mutants

Confirmation of the identity of the mutants, and isolation of homozygous lines, was performed by PCR amplification and sequencing using a combination of gene- and T-DNA-specific (SALK-LB) primers (**Table S1**). Briefly, genomic DNA from individual plants was extracted using DNeasy Plant Minikit (Qiagen) and used as template for PCR amplification with ExTaq DNA polymerase (Takara). Amplified bands from SALK-LB and gene-specific combinations were separated in 1% (w/v) agarose gels stained with EtBr and purified by gel scission using Qiaquick Gel Extraction Kit (Qiagen). Purified DNA was used for sequencing using the same primer set. The position of T-DNA insertion was confirmed by local alignment with the *AtPAO5* gene (www.ebi.ac.uk/Tools/msa/muscle) and BLAST query (www.arabidopsis.org). Transcript levels of *AtPAO5* in the homozygous lines were determined by real-time PCR (RT-PCR) from cDNA of *atpao5-2* and *atpao5-3* mutants using the *AtPAO5*-specific primers and the following PCR conditions: 95 °C 5 min, 30 cycles (95 °C 15 s, 55 °C 20 s, 72 °C 1 min), 72 °C 10 min, and primers specified in **Table S1**. Single T-DNA insertion lines were confirmed by the occurrence of 1:3 segregation for the presence of kanamycin resistance gene (*NPTII*) in the progeny of segregating *atpao5-2* and *atpao5-3* mutants.

Determination of PA levels

The levels of free Put, Spd and Spm were determined by high-performance liquid chromatography (HPLC) separation of dansyl chloride-derived PAs as described (Alcázar *et al.* 2005). Free tSpm levels were determined by PA benzoylation according to Kim *et al.* 2014.

Determination of water content

Water content (WC) was determined using the fresh weight (FW) and dry weight (DW) and calculated as $\% \text{ WC} = (\text{FW} - \text{DW}) / \text{FW}$. Dry weight was obtained after drying samples at 60 °C for 48 h. The shoots of at least 10 individual plants were used to calculate % WC in every point of analysis and genotype.

Determination of chlorophyll levels

Chlorophyll extracts were obtained from leaves of plants grown on soil or seedlings grown *in vitro* at indicated time points, using buffered acetone (acetone/Tris-HCl buffer 80:20 vol pH 7.8). Total chlorophyll levels were calculated according to Porra (2002).

Determination of sodium levels

For the determination of Na⁺ content, lyophilized shoot tissue was digested in concentrated nitric acid at 80 °C for 1 h. After digestion, the acid was diluted with double deionized water. Mineral element concentrations in the supernatants were determined by inductively coupled plasma optical emission spectroscopy (Perkin Elmer Optima 2000).

Determination of ABA and JA levels

ABA and JA levels were analysed by HPLC coupled to tandem mass spectrometry (Durgbanshi *et al.* 2005). Briefly, the plant tissue was extracted in ultrapure water using a tissue homogenizer. Before extraction, samples were spiked with 100 ng of *d6*-ABA and *dh*-JA (Gómez-Cadenas *et al.* 2002; Durgbanshi *et al.* 2005). After extraction and centrifugation, the pH of the supernatant was adjusted to 3.0 and partitioned twice against diethyl ether. The organic layers were combined and evaporated. The dry residue was resuspended in a solution of water:methanol (9:1, v/v), filtered and injected into a HPLC system (Alliance 2695, Waters Corp.). The phytohormone fraction was separated on a reversed-phase C18 column using methanol and 0.01% acetic acid in water as solvents. The mass spectrometer, a triple quadrupole (Quattro LC, Micromass Ltd.), was operated in negative ionization electrospray mode, and the different plant hormones were detected according to their specific transitions using a multi-residue mass spectrometric method as described in detail by Durgbanshi *et al.* 2005.

Determination of hydrogen peroxide levels

Hydrogen peroxide levels were determined as described by Loreto & Velikova (2001). Frozen leaf material (~0.1 g) was homogenized on ice with 0.1% (w/v) trichloroacetic acid. The homogenate was centrifuged at 15 000×g for 15 min at 4 °C. Then, 0.5 mL of the supernatant was mixed with 0.5 mL of 10 mM potassium phosphate buffer (pH 7.0) and 1 mL of 1 M KI. The absorbance of the assay mixture was determined at 390 nm and the content of H₂O₂ calculated based on standard curves of known hydrogen peroxide concentrations.

Microarray expression analyses

Four-week-old *atpao5-3* and wild-type plants were used for microarray analyses. RNA samples were extracted in triplicate from pooled samples of five plants per genotype using TRIzol reagent (Invitrogen). Transcriptomes were analysed using 1 µg of total RNA as starting material. Targets were prepared with MessageAmp II-Biotin Enhanced Single Round aRNA Amplification Kit (Ambion) and hybridized to ATH1 gene chips (Affymetrix). Microarray data and gene ontology (GO) classification were analysed using GeneSpring GX 10 software (Agilent Technologies) and Database for Annotation, Visualization and Integrated Discovery v6.7 (Huang da *et al.* 2009). A threshold of significance at $P = 0.05$ and a cut-off value of $\geq \pm 2$ -fold change was used for the identification of genes that were differentially expressed. Microarray data has been deposited in ArrayExpress under accession number E-MTAB-3817.

Quantitative real-time PCR gene expression analyses

Total RNA isolated from 4-week-old plants or 7-day-old seedlings, as indicated, was extracted using TRIzol reagent (Invitrogen). Two micrograms of RNA was treated with DNase I (Invitrogen) and first-strand cDNA synthesized using Superscript II (Invitrogen) and oligo dT. Quantitative real-time PCR using SYBR Green I dye method was performed on Roche LightCycler 480 II detector system following the PCR conditions: 95 °C 2 min, 40 cycles (95 °C, 15 s; 60 °C, 30 s; 68 °C, 20 s). Standard curves were performed for quantification. Primers used for gene expression analyses are listed in Table S1. Quantitative RT-PCR (qRT-PCR) analyses were always performed on three biological replicates with three technical replicates each using *ACTIN2* (*AT3G18780*) as housekeeping gene.

Gas chromatography/mass spectrometry metabolite profiling

Metabolite profiling was performed from leaf samples (120 mg) of wild-type and *atpao5-3* plants using seven independent biological replicates for each time point under study. Samples were shock frozen in liquid nitrogen, and the fraction enriched in polar primary metabolites was prepared and processed as described (Erban *et al.* 2007). Gas chromatography coupled to electron impact ionization-time of flight-mass spectrometry (GC/EI-TOF-MS) was performed using an Agilent 6890N24 gas

chromatograph hyphenated to a Pegasus III time-of-flight mass spectrometer (LECO, St. Joseph, USA) (Wagner *et al.* 2003). Chromatograms were acquired and processed by CHROMATOF software 1.00, Pegasus driver 1.61 (LECO; <http://www.leco.de>). Selective peak heights representing arbitrary mass spectral ion currents were normalized by the dry weight of the sample and to an internal standard, which was added upon extraction of the polar metabolite fraction. GC-TOF-MS chromatography data processing was performed using the TagFinder software (Luedemann *et al.* 2008). Metabolites were identified under manual supervision using the TagFinder, the NIST08 software (<http://chemdata.nist.gov>) and the mass spectral and retention time index reference collection of the Golm Metabolome Database (Kopka *et al.* 2005; Hummel *et al.* 2010). Only metabolites identified in at least four of seven replicates were considered. Wild type (0 h) was used as reference condition for the calculation of Log₂ ratios of metabolites. Principal component analysis (PCA) was performed using *R* (www.r-project.org). Hierarchical clustering with Pearson correlation was performed using the software MultiExperiment Viewer (www.tm4.org/mev; version 27 4.8.1).

ACKNOWLEDGEMENTS

We thank Prof. Tomonobu Kusano for providing advice in the determination of thermospermine levels in *Arabidopsis thaliana*. Research in this work has been supported by the Spanish Ministerio de Ciencia e Innovación (BIO2011-29683 and CSD2007-00036), and the Generalitat de Catalunya (SGR2009-1060) to A.F.T. R.A. acknowledges support from the Ramón y Cajal Program (RYC-2011-07847) of the Ministerio de Ciencia e Innovación (Spain).

REFERENCES

- Ahou A., Martignago D., Alabdallah O., Tavazza R., Stano P., Macone A., ... Tavladoraki P. (2014) A plant spermine oxidase/dehydrogenase regulated by the proteasome and polyamines. *Journal of Experimental Botany* **65**, 1585–1603.
- Alcázar R., Altabella T., Marco F., Bortolotti C., Reymond M., Koncz C., ... Tiburcio A.F. (2010) Polyamines: molecules with regulatory functions in plant abiotic stress tolerance. *Planta* **231**, 1237–1249.
- Alcázar R., Cuevas J.C., Patrón M., Altabella T., & Tiburcio A.F. (2006a) Abscisic acid modulates polyamine metabolism under water stress in *Arabidopsis thaliana*. *Physiologia Plantarum* **128**, 448–455.
- Alcázar R., García-Martínez J.L., Cuevas J.C., Tiburcio A.F., & Altabella T. (2005) Overexpression of *ADC2* in *Arabidopsis* induces dwarfism and late-flowering through GA deficiency. *The Plant Journal* **43**, 425–436.

- Alcázar R., Marco F., Cuevas J.C., Patron M., Ferrando A., Carrasco P., ... Altabella T. (2006b) Involvement of polyamines in plant response to abiotic stress. *Biotechnology Letters* **28**, 1867–1876.
- Alet A.I., Sánchez D.H., Cuevas J.C., Marina M., Carrasco P., Altabella T., ... Ruiz O.A. (2012) New insights into the role of spermine in *Arabidopsis thaliana* under long-term salt stress. *Plant Science* **182**, 94–100.
- Bagni N., Ruiz-Carrasco K., Franceschetti M., Fornalè S., Fornasiero R.B., & Tassoni A. (2006) Polyamine metabolism and biosynthetic gene expression in *Arabidopsis thaliana* under salt stress. *Plant Physiology and Biochemistry* **44**, 776–786.
- Bitrián M., Zarza X., Altabella T., Tiburcio A.F., & Alcázar R. (2012) Polyamines under abiotic stress: metabolic crossroads and hormonal crosstalks in plants. *Metabolites* **2**, 516–528.
- Chen T.H. & Murata N. (2002) Enhancement of tolerance of abiotic stress by metabolic engineering of betaines and other compatible solutes. *Current Opinion in Plant Biology* **5**, 250–257.
- Chen T.H.H. & Murata N. (2011) Glycinebetaine protects plants against abiotic stress: mechanisms and biotechnological applications. *Plant, Cell & Environment* **34**, 1–20.
- Cona A., Rea G., Angelini R., Federico R., & Tavladoraki P. (2006) Functions of amine oxidases in plant development and defence. *Trends in Plant Science* **11**, 80–88.
- Cuevas J.C., López-Cobollo R., Alcázar R., Zarza X., Koncz C., Altabella T., ... Ferrando A. (2008b) Putrescine is involved in *Arabidopsis* freezing tolerance and cold acclimation by regulating abscisic acid levels in response to low temperature. *Plant Physiology* **148**, 1094–1105.
- Dong W., Wang M., Xu F., Quan T., Peng K., Xiao L., & Xia G. (2013) Wheat oxophytodienoate reductase gene *TaOPR1* confers salinity tolerance via enhancement of abscisic acid signaling and reactive oxygen species scavenging. *Plant Physiology* **161**, 1217–1228.
- Durgbanshi A., Arbona V., Pozo O., Miersch O., Sancho J. V., & Gómez-Cadenas A. (2005) Simultaneous determination of multiple phytohormones in plant extracts by liquid chromatography-electrospray tandem mass spectrometry. *Journal of Agricultural and Food Chemistry* **53**, 8437–8442.
- Erban A., Schauer N., Fernie A.R., & Kopka J. (2007) Nonsupervised construction and application of mass spectral and retention time index libraries from time-of-flight gas chromatography-mass spectrometry metabolite profiles. *Methods in Molecular Biology* **358**, 19–38.
- Fernández-Calvo P., Chini A., Fernández-Barbero G., Chico J.-M., Gimenez-Ibanez S., Geerinck J., ... Solano R. (2011) The *Arabidopsis* bHLH transcription factors MYC3 and MYC4 are targets of JAZ repressors and act additively with MYC2 in the activation of jasmonate responses. *Plant Cell* **23**, 701–715.
- Fincato P., Moschou P.N., Spedaletti V., Tavazza R., Angelini R., Federico R., ... Tavladoraki P. (2011) Functional diversity inside the *Arabidopsis* Polyamine Oxidase gene family. *Journal of Experimental Botany* **62**, 1155–1168.
- Furumoto T., Yamaguchi T., Ohshima-Ichie Y., Nakamura M., Tsuchida-Iwata Y., Shimamura M., ... Izui K. (2011) A plastidial sodium-dependent pyruvate transporter. *Nature* **476**, 472–475.
- Gaxiola R.A., Li J., Undurraga S., Dang L.M., Allen G.J., Alper S.L., & Fink G.R. (2001) Drought- and salt-tolerant plants result from overexpression of the *AVP1* H⁺-pump.

- Proceedings of the National Academy of Sciences of the United States of America* **98**, 11444–9.
- Ghuge S.A., Carucci A., Rodrigues Pousada R.A., Tisi A., Franchi S., Tavladoraki P., ... Cona A. (2015) The apoplastic copper amine oxidase *AtAO1* mediates jasmonic acid-induced protoxylem differentiation in *Arabidopsis* roots. *Plant Physiology* **168**, 690–707.
- Gómez-Cadenas A., Pozo O.J., García-Augustín P., & Sancho J. V (2002) Direct analysis of abscisic acid in crude plant extracts by liquid chromatography–electrospray/tandem mass spectrometry. *Phytochemical Analysis* **13**, 228–234.
- Gong Q., Li P., Ma S., Indu Rupassara S., & Bohnert H.J. (2005) Salinity stress adaptation competence in the extremophile *Thellungiella halophila* in comparison with its relative *Arabidopsis thaliana*. *The Plant Journal* **44**, 826–839.
- Hasegawa P.M. (2013) Sodium (Na⁺) homeostasis and salt tolerance of plants. *Environmental and Experimental Botany* **92**, 19–31.
- Hill C.B., Jha D., Bacic A., Tester M., & Roessner U. (2013) Characterization of ion contents and metabolic responses to salt stress of different *Arabidopsis AtHKT1;1* genotypes and their parental strains. *Molecular Plant* **6**, 350–368.
- Huang da W., Sherman B.T., Lempicki R.A. (2009) Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature Protocols* **4**, 44–57.
- Hummel J., Strehmel N., Selbig J., Walther D., & Kopka J. (2010) Decision tree supported substructure prediction of metabolites from GC-MS profiles. *Metabolomics* **6**, 322–333.
- Kamada-Nobusada T., Hayashi M., Fukazawa M., Sakakibara H., & Nishimura M. (2008) A putative peroxisomal polyamine oxidase, *AtPAO4*, is involved in polyamine catabolism in *Arabidopsis thaliana*. *Plant and Cell Physiology* **49**, 1272–1282.
- Kang D.-J., Seo Y.-J., Lee J.-D., Ishii R., Kim K.U., Shin D.H., ... Lee I.-J. (2005) Jasmonic acid differentially affects growth, ion uptake and abscisic acid concentration in salt-tolerant and salt-sensitive rice cultivars. *Journal of Agronomy and Crop Science* **191**, 273–282.
- Kazan K. (2015) Diverse roles of jasmonates and ethylene in abiotic stress tolerance. *Trends in Plant Science* **20**, 219–229.
- Kempa S., Krasensky J., Dal Santo S., Kopka J., & Jonak C. (2008) A central role of abscisic acid in stress-regulated carbohydrate metabolism. *PLoS One* **3**, e3935.
- Kim D.W., Watanabe K., Murayama C., Izawa S., Niitsu M., Michael A.J., ... Kusano T. (2014) Polyamine Oxidase5 regulates *Arabidopsis* growth through thermospermine oxidase activity. *Plant Physiology* **165**, 1575–1590.
- Kopka J., Schauer N., Krueger S., Birkemeyer C., Usadel B., Bergmüller E., ... Steinhauser D. (2005) GMD@CSB.DB: the Golm Metabolome Database. *Bioinformatics* **21**, 1635–1638.
- Krasensky J. & Jonak C. (2012) Drought, salt, and temperature stress-induced metabolic rearrangements and regulatory networks. *Journal of Experimental Botany* **63**, 1593–1608.
- Kusano T., Berberich T., Tateda C., & Takahashi Y. (2008) Polyamines: essential factors for growth and survival. *Planta* **228**, 367–381.
- Loreto F. & Velikova V. (2001) Isoprene produced by leaves protects the photosynthetic apparatus against ozone damage, quenches ozone products, and reduces lipid peroxidation of cellular membranes. *Plant Physiology* **127**, 1781–1787.
- Luedemann A., Strassburg K., Erban A., & Kopka J. (2008) TagFinder for the quantitative analysis of gas chromatography–mass spectrometry (GC-MS)-based metabolite profiling experiments. *Bioinformatics* **24**, 732–737.

- Møller S.G. & McPherson M.J. (1998) Developmental expression and biochemical analysis of the *Arabidopsis* *ATAO1* gene encoding an H₂O₂-generating diamine oxidase. *The Plant Journal* **13**, 781–791.
- Moons A., Prinsen E., Bauw G., & Van Montagu M. (1997) Antagonistic effects of abscisic acid and jasmonates on salt stress-inducible transcripts in rice roots. *Plant Cell* **9**, 2243–2259.
- Moschou P.N., Delis I.D., Paschalidis K.A., & Roubelakis-Angelakis K.A. (2008a) Transgenic tobacco plants overexpressing *Polyamine Oxidase* are not able to cope with oxidative burst generated by abiotic factors. *Physiologia Plantarum* **133**, 140–156.
- Moschou P.N., Paschalidis K. a, Delis I.D., Andriopoulou A.H., Lagiotis G.D., Yakoumakis D.I., & Roubelakis-Angelakis K.A. (2008b) Spermidine exodus and oxidation in the apoplast induced by abiotic stress is responsible for H₂O₂ signatures that direct tolerance responses in tobacco. *Plant Cell* **20**, 1708–1724.
- Moschou P.N., Sanmartin M., Andriopoulou A.H., Rojo E., Sanchez-Serrano J.J., & Roubelakis-Angelakis K.A. (2008c) Bridging the gap between plant and mammalian polyamine catabolism: a novel peroxisomal polyamine oxidase responsible for a full back-conversion pathway in *Arabidopsis*. *Plant Physiology* **147**, 1845–1857.
- Moschou P.N., Wu J., Cona A., Tavladoraki P., Angelini R., & Roubelakis-Angelakis K.A. (2012) The polyamines and their catabolic products are significant players in the turnover of nitrogenous molecules in plants. *Journal of Experimental Botany* **63**, 5003–5015.
- Müller M., Kunz H.-H., Schroeder J.I., Kemp G., Young H.S., & Neuhaus H.E. (2014) Decreased capacity for sodium export out of *Arabidopsis* chloroplasts impairs salt tolerance, photosynthesis and plant performance. *The Plant Journal* **78**, 646–658.
- Munns R. & Tester M. (2008) Mechanisms of salinity tolerance. *Annual Review of Plant Biology* **59**, 651–681.
- Naka Y., Watanabe K., Sagor G.H.M., Niitsu M., Pillai M.A., Kusano T., & Takahashi Y. (2010) Quantitative analysis of plant polyamines including thermospermine during growth and salinity stress. *Plant Physiology and Biochemistry* **48**, 527–533.
- Pedranzani H., Racagni G., Alemano S., Miersch O., Ramírez I., Peña-Cortés H., ... Abdala G. Salt tolerant tomato plants show increased levels of jasmonic acid. *Plant Growth Regulation* **41**, 149–158.
- Porra R.J. (2002) The chequered history of the development and use of simultaneous equations for the accurate determination of chlorophylls *a* and *b*. *Photosynthesis Research* **73**, 149–156.
- Pottosin I. & Shabala S. (2014) Polyamines control of cation transport across plant membranes: implications for ion homeostasis and abiotic stress signaling. *Frontiers in Plant Science* **5**, 154.
- Pottosin I., Velarde-Buendía A.M., Bose J., Fuglsang A.T., & Shabala S. (2014) Polyamines cause plasma membrane depolarization, activate Ca²⁺, and modulate H⁺-ATPase pump activity in pea roots. *Journal of Experimental Botany* **65**, 2463–2472.
- Reumann S., Quan S., Aung K., Yang P., Manandhar-Shrestha K., Holbrook D., ... Hu J. (2009) In-depth proteome analysis of *Arabidopsis* leaf peroxisomes combined with in vivo subcellular targeting verification indicates novel metabolic and regulatory functions of peroxisomes. *Plant Physiology* **150**, 125–143.

- Saha J., Brauer E.K., Sengupta A., Popescu S.C., Gupta K., & Gupta B. (2015) Polyamines as redox homeostasis regulators during salt stress in plants. *Frontiers in Environmental Science* **3**, 1–13.
- Sánchez D.H., Siahpoosh M.R., Roessner U., Udvardi M., & Kopka J. (2008) Plant metabolomics reveals conserved and divergent metabolic responses to salinity. *Physiologia Plantarum* **132**, 209–219.
- Székely G., Abrahám E., Csépló A., Rigó G., Zsigmond L., Csiszár J., ... Szabados L. (2008) Duplicated *P5CS* genes of *Arabidopsis* play distinct roles in stress regulation and developmental control of proline biosynthesis. *The Plant Journal* **53**, 11–28.
- Takahashi T. & Kakehi J.-I. (2010) Polyamines: ubiquitous polycations with unique roles in growth and stress responses. *Annals of Botany* **105**, 1–6.
- Takahashi Y., Cong R., Sagor G.H.M., Niitsu M., Berberich T., & Kusano T. (2010) Characterization of five polyamine oxidase isoforms in *Arabidopsis thaliana*. *Plant Cell Reports* **29**, 955–965.
- Takano A., Kakehi J.-I., & Takahashi T. (2012) Thermospermine is not a minor polyamine in the plant kingdom. *Plant and Cell Physiology* **53**, 606–616.
- Tanou G., Ziogas V., Belghazi M., Christou A., Filippou P., Job D., Fotopoulos V., & Molassiotis A. (2014). Polyamines reprogram oxidative and nitrosative status and the proteome of citrus plants exposed to salinity stress. *Plant Cell & Environment* **37**, 864–885.
- Tassoni A., Franceschetti M., & Bagni N. (2008) Polyamines and salt stress response and tolerance in *Arabidopsis thaliana* flowers. *Plant Physiology and Biochemistry* **46**, 607–613.
- Tavladoraki P., Cona A., Federico R., Tempera G., Viceconte N., Saccoccio S., ... Agostinelli E. (2012) Polyamine catabolism: target for antiproliferative therapies in animals and stress tolerance strategies in plants. *Amino acids* **42**, 411–426.
- Tavladoraki P., Rossi M.N., Saccuti G., Perez-Amador M.A., Polticelli F., Angelini R., & Federico R. (2006) Heterologous expression and biochemical characterization of a polyamine oxidase from *Arabidopsis* involved in polyamine back conversion. *Plant Physiology* **141**, 1519–1532.
- Tiburcio A.F., Altabella T., Bitrián M., & Alcázar R. (2014) The roles of polyamines during the lifespan of plants: from development to stress. *Planta* **240**, 1–18.
- Tong W., Yoshimoto K., Kakehi J.-I., Motose H., Niitsu M., & Takahashi T. (2014) Thermospermine modulates expression of auxin-related genes in *Arabidopsis*. *Frontiers in Plant Science* **5**, 94.
- Toumi I., Moschou P.N., Paschalidis K.A., Bouamama B., Ben Salem-Fnayou A., Ghorbel A.W., ... Roubelakis-Angelakis K.A. (2010) Abscisic acid signals reorientation of polyamine metabolism to orchestrate stress responses via the polyamine exodus pathway in grapevine. *Journal of Plant Physiology* **167**, 519–525.
- Uno Y., Furihata T., Abe H., Yoshida R., Shinozaki K., Yamaguchi-Shinozaki K. (2000) *Arabidopsis* basic leucine zipper transcription factors involved in an abscisic acid-dependent signal transduction pathway under drought and high-salinity conditions. *Proceedings of the National Academy of Sciences of the United States of America* **97**, 11632–11637.
- Urano K., Yoshiba Y., Nanjo T., Igarashi Y., Seki M., Sekiguchi F., ... Shinozaki K. (2003) Characterization of *Arabidopsis* genes involved in biosynthesis of polyamines in abiotic stress responses and developmental stages. *Plant, Cell & Environment* **26**, 1917–1926.

- Vera-Sirera F., Minguet E.G., Singh S.K., Ljung K., Tuominen H., Blázquez M.A., & Carbonell J. (2010) Role of polyamines in plant vascular development. *Plant Physiology and Biochemistry* **48**, 534–539.
- Wagner C., Sefkow M., & Kopka J. (2003) Construction and application of a mass spectral and retention time index database generated from plant GC/EI-TOF-MS metabolite profiles. *Phytochemistry* **62**, 887–900.
- Yamaguchi K., Takahashi Y., Berberich T., Imai A., Miyazaki A., Takahashi T., ... Kusano T. (2006) The polyamine spermine protects against high salt stress in *Arabidopsis thaliana*. *FEBS Letters* **580**, 6783–6788.
- Zhao X., Wang W., Zhang F., Deng J., Li Z., & Fu B. (2014a) Comparative metabolite profiling of two rice genotypes with contrasting salt stress tolerance at the seedling stage. *PLoS One* **9**, e108020.
- Zhao Y., Dong W., Zhang N., Ai X., Wang M., Huang Z., ... Xia G. (2014b) A wheat allene oxide cyclase gene enhances salinity tolerance via jasmonate signaling. *Plant Physiology* **164**, 1068–1076.

SUPPLEMENTAL MATERIAL

Supplemental Tables S1-S20 may be found in the online version of this article.
<http://onlinelibrary.wiley.com/doi/10.1111/pce.12714/supinfo>

Table S1. List of oligonucleotides used in this work.

Table S2. Genes differentially expressed in the comparison between *atpao5-3* 3 h and *atpao5-3* 0 h of 50 mM NaCl treatment (p3p0).

Table S3. Genes differentially expressed in the comparison between wild-type 3 h and wild-type 0 h of 50 mM NaCl treatment (w3w0).

Table S4. Genes differentially expressed in the comparison between *atpao5-3* 0 h and wild-type 0 h (p0w0).

Table S5. Genes differentially expressed in the comparison between *atpao5-3* 3 h and wild-type 3 h of 50 mM NaCl treatment (p3w3).

Table S6. Genes differentially expressed only in the comparison between *atpao5-3* 3 h and *atpao5-3* 0 h of 50 mM NaCl treatment (p3p0 only).

Table S7. Genes differentially expressed only in the comparison between wild-type 3 h and wild-type 0 h of 50 mM NaCl treatment (w3w0 only).

Table S8. Genes differentially expressed only in the comparison between *atpao5-3* 0 h and wild-type 0 h of 50 mM NaCl treatment (p0w0 only).

Table S9. Genes differentially expressed only in the comparison between *atpao5-3* 3 h and wild-type 3 h of 50 mM NaCl treatment (p3w3 only).

Table S10. Genes differentially expressed and common between p3p0 and w3w0.

Table S11. Genes differentially expressed and common between p3p0 and p0w0.

Table S12. Genes differentially expressed and common between p3p0 and p3w3.

Table S13. Genes differentially expressed and common between w3w0 and p0w0.

Table S14. Genes differentially expressed and common between w3w0 and p3w3.

Table S15. Genes differentially expressed and common between p0w0 and p3w3.

Table S16. Microarray analysis of JA biosynthesis and signalling genes in the comparisons p0w0, p3w3, w3w0 and p3p0.

Table S17. Metabolite analysis of *atpao5-3* and wild-type plants exposed to NaCl stress.

Table S18. PC1 and PC2 loadings of wild-type and *atpao5-3* metabolite profiles 24 h after treatment with 0, 50, 100 and 150 mM NaCl.

Table S19. Late salt-responsive metabolites in wild-type. Metabolites exhibiting significant differences between wild-type and *atpao5-3* are highlighted.

Table S20. List of metabolites differentially accumulated in *atpao5-3* in response to salt stress.


a

Figure S1. (a) Schematic representation of T-DNA insertion positions in *atpao5-2* and *atpao5-3* mutants. The *AtPAO5* gene carries a single exon (highlighted in black). 5' and 3' uORFs are shown in grey. (b) RT-PCR expression analysis of *AtPAO5* transcript levels in wild-type, *atpao5-2* and *atpao5-3* homozygous mutants. *ACTIN2* was used as house-keeping gene.



b

Figure S2. Quantitation of sodium levels in leaves of *atpao5-3* and wild-type plants exposed to gradual increases of NaCl from 50 to 150 mM. Values are the mean \pm SD of five technical replicates. The experiment was repeated three times with similar results.




Figure S3. Hydrogen peroxide levels in *atpao5-3* and wild-type plants exposed to gradual increases of NaCl between 50 and 150 mM. Values are the mean \pm SD of five technical replicates. The analysis was repeated in two additional independent experiments showing similar results.





Figure S4. Quantitative real-time gene expression analyses of *BASS2* and *NHD1* genes from 7-day-old wild-type seedlings treated with 100 μ M tSpm or 100 μ M Spm during 0-24 h. Values are the mean from three biological replicates \pm SD, each with three technical replicates. Letters indicate values that are significantly different according to Student-Newman-Keuls test at *P* value <0.05.






Figure S5. Quantitative real-time gene expression analyses of JA biosynthesis (*LOX4* and *OPCL1*) and signalling (*JAZ5*) genes from 12-days-old wild-type seedlings treated for 15 min with different concentrations (0, 5, 10, 20, and 100 μM) of tSpm or Spm. Values are the mean from three biological replicates \pm SD, each with three technical replicates. Letters indicate values that are significantly different according to Student-Newman-Keuls test at P value < 0.05 .




Figure S6. Sum of putrescine and agmatine levels in *atpao5-3* and wild-type plants exposed to salt stress. Log₂ ratios ± SE are relative to wild-type 0 h.