



UvA-DARE (Digital Academic Repository)

Elliptic Genera and 3d Gravity

Benjamin, N.; Cheng, M.C.N.; Kachru, S.; Moore, G.W.; Paquette, N.M.

DOI

[10.1007/s00023-016-0469-6](https://doi.org/10.1007/s00023-016-0469-6)

Publication date

2016

Document Version

Final published version

Published in

Annales Henri Poincaré

License

CC BY

[Link to publication](#)

Citation for published version (APA):

Benjamin, N., Cheng, M. C. N., Kachru, S., Moore, G. W., & Paquette, N. M. (2016). Elliptic Genera and 3d Gravity. *Annales Henri Poincaré*, 17(10), 2623-2662. <https://doi.org/10.1007/s00023-016-0469-6>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, P.O. Box 19185, 1000 GD Amsterdam, The Netherlands. You will be contacted as soon as possible.



Elliptic Genera and 3d Gravity

Nathan Benjamin, Miranda C. N. Cheng, Shamit Kachru,
Gregory W. Moore and Natalie M. Paquette

Abstract. We describe general constraints on the elliptic genus of a 2d supersymmetric conformal field theory which has a gravity dual with large radius in Planck units. We give examples of theories which do and do not satisfy the bounds we derive, by describing the elliptic genera of symmetric product orbifolds of $K3$, product manifolds, certain simple families of Calabi–Yau hypersurfaces, and symmetric products of the “Monster CFT”. We discuss the distinction between theories with supergravity duals and those whose duals have strings at the scale set by the AdS curvature. Under natural assumptions, we attempt to quantify the fraction of (2,2) supersymmetric conformal theories which admit a weakly curved gravity description, at large central charge.

1. Introduction

The AdS/CFT correspondence [1] provides a concrete framework for holography, where very particular d dimensional quantum field theories can capture the dynamics of quantum gravity in $d + 1$ spacetime dimensions. A natural question from the outset has been: “which class of quantum field theories is dual to (large radius, weakly coupled) Einstein gravity theories?”

In a recent paper [2], interesting progress was made on this issue in the particular case of two-dimensional CFTs. The authors of [2] make the plausible assumption that a weakly coupled gravitational theory should reproduce the most basic aspects of the phase structure known in all of the simple examples of AdS/CFT. In particular, as one raises the temperature, there should be a phase transition at a critical temperature (usually taken to be $\beta^* = \frac{1}{kT^*} = 2\pi$) between a “gas of particles” and a black hole geometry [3]—the Hawking–Page transition [4]. By requiring that outside a small neighborhood of the critical temperature the thermal partition function should be dominated by BTZ black

M. C. N. Cheng: On leave from CNRS, France.

holes at high T , or the ground state at low T , one finds interesting constraints on the spectrum of any putative dual conformal field theory.

A significant consequence of this constraint is the derivation of the Bekenstein–Hawking black hole entropy (expressed here in the ensemble where one keeps track only of the total energy $E = h + \bar{h} - \frac{c}{12}$)

$$S(E) \sim 2\pi \sqrt{\frac{cE}{3}}, \quad E = h + \bar{h} - \frac{c}{12} \quad (1.1)$$

for $E > \frac{c}{12}$ and $c \gg 1$. Notice that this is the regime where we expect the Bekenstein–Hawking formula to give a good approximation of the black hole entropy on gravitational grounds. It is different from the regime of applicability of the usual Cardy formula based on familiar modular form arguments ($\frac{E}{c} \gg 1$).

Here, we turn our attention to 2d supersymmetric theories. In two-dimensional theories with at least $(0, 1)$ supersymmetry and left and right-moving \mathbb{Z}_2 fermion number symmetries, one can define the elliptic genus [5–8]. We will focus on the special case of $(2, 2)$ supersymmetry in this paper, but we expect that many of our considerations could be suitably generalized. In the $(2, 2)$ case, the elliptic genus associates to a 2d SCFT a weak Jacobi form; detailed knowledge of the space of such forms (see e.g., [9]) will allow us to make some strong statements about CFT/gravity duality in this case. Prominent cases of such 2d supersymmetric theories in the AdS/CFT correspondence include those arising in D-brane constructions of supersymmetric black strings [10], where the near-horizon geometry has a dual given by a σ -model with target M^N/S_N for $M = K3$ or T^4 . By requiring the Bekenstein–Hawking formula for these black objects to apply in the black hole regime, we derive a constraint on the coefficients of the elliptic genus.

Intuitively, the condition that the CFT elliptic genus exhibits an enlarged regime of applicability of the Bekenstein–Hawking entropy (which turns out to warrant a Hawking–Page transition) hints that there is indeed a weakly coupled gravity dual. In the simplest perturbative string theory constructions of AdS, there are at least three scales of interest—the Planck scale M_{Planck} , the string scale M_{string} , and the inverse AdS radius $\frac{1}{\ell_{\text{AdS}}}$. (There are also in general one or more Kaluza–Klein scales—for simplicity we are imagining constructions like the Freund–Rubin construction where the KK scale coincides with the AdS radius.) The most conventional regime of understanding string models is when $M_{\text{Planck}} \gg M_{\text{string}} \gg \frac{1}{\ell_{\text{AdS}}}$. However, the conditions we impose are also satisfied in some theories where there is no separation of scales between M_{string} and $\frac{1}{\ell_{\text{AdS}}}$ apparent in the elliptic genus. We therefore also discuss further criteria on the coefficients of the elliptic genus which may distinguish between theories with a separation of scales between supergravity and string modes, and theories without such separation.

It is important to keep in mind that our necessary condition serves only as an indicator of whether there might be a weakly coupled gravity dual to some region in the moduli space of the superconformal field theory. In simple examples, the moduli space will have other generic phases characterized by

duals with no simple geometric description, and the large radius gravity dual would characterize only a small region of the SCFT moduli space. However, as the elliptic genus is an invariant calculable (in principle) in this small region, it will have the properties expected of a theory with a weakly coupled gravity description if the SCFT admits such a description anywhere in its moduli space.

This paper is organized as follows. In Sect. 2, we review some basic facts about Jacobi forms. In Sect. 3, we describe the constraint we wish to place on the Fourier coefficients of these forms, following a similar philosophy to [2]. In Sect. 4, we check the bound on various simple constructions: $K3$ symmetric product orbifolds (which provide some of the simplest examples of $\text{AdS}_3/\text{CFT}_2$ and do satisfy the bound), product manifolds, a family of Calabi–Yau spaces going off to large dimension, and a symmetric product of the “Monster CFT”. As some of the examples will fail, we see that the bound does have teeth—there are simple examples of (2,2) superconformal field theories at large central charge that violate it. In Sect. 5, focusing on the distinctions between the $K3$ symmetric product and the “Monster” symmetric product, we discuss the distinction between low-energy supergravity theories and low-energy string theories. In Sect. 6, we attempt to quantify “the fraction of supersymmetric theories at large central charge which admit a gravity dual”, using a natural metric on a relevant (suitably projectivized) space of weak Jacobi forms. Detailed arguments supporting some of the assertions in the main body of the paper are provided in two appendices.

2. Modularity Properties

We can define the following elliptic genera for any 2d SCFT with at least (1, 1) supersymmetry and left and right-moving fermion quantum numbers. Denote by L_n, \bar{L}_n the left and right Virasoro generators, and F, \bar{F} the left and right-moving fermion number. The NS sector elliptic genus can be defined via:

$$Z_{\text{NS},+}(\tau) = \text{Tr}_{\text{NS},R}(-1)^{\bar{F}} q^{L_0-c/24} \bar{q}^{\bar{L}_0-c/24} . \tag{2.1}$$

It is a (weakly holomorphic) modular form under the congruence subgroup Γ_θ , defined in (3.34). Similar definitions apply in other sectors:

$$\chi = \text{Tr}_{R,R}(-1)^{F+\bar{F}} q^{L_0-c/24} \bar{q}^{\bar{L}_0-c/24} \tag{2.2}$$

$$Z_{R,+}(\tau) = \text{Tr}_{R,R}(-1)^{\bar{F}} q^{L_0-c/24} \bar{q}^{\bar{L}_0-c/24} \tag{2.3}$$

$$Z_{\text{NS},-}(\tau) = \text{Tr}_{\text{NS},R}(-1)^{F+\bar{F}} q^{L_0-c/24} \bar{q}^{\bar{L}_0-c/24} . \tag{2.4}$$

Here, $q = e^{2\pi i\tau}$ where τ takes values in the upper-half plane, and we have assumed equal left and right-moving central charges, $c_L = c_R = c$.

For the most part, we will consider theories with additional structure, e.g., (2, 2) superconformal theories. In fact for any (0, 2) theory with a left-moving $U(1)$ symmetry, and so in particular for any (2, 2) SCFT, one can define a refined elliptic genus as

$$Z_{R,R}(\tau, z) = \text{Tr}_{R,R}(-1)^{F+\bar{F}} q^{L_0-c/24} \bar{q}^{\bar{L}_0-c/24} y^{J_0} . \tag{2.5}$$

Here, $y = e^{2\pi iz}$. The additional symmetry promotes the two-variable elliptic genus into a weak Jacobi form [11]. We will also consider

$$\begin{aligned} Z_{\text{NS},R}(\tau, z) &= \text{Tr}_{\text{NS},R}(-1)^{\bar{F}} q^{L_0 - c/24} \bar{q}^{\bar{L}_0 - c/24} y^{J_0} \\ &= Z_{R,R} \left(\tau, z + \frac{\tau + 1}{2} \right) q^{\frac{c}{24}} y^{\frac{c}{6}}. \end{aligned} \tag{2.6}$$

Note that we could define $Z_{\text{NS},\text{NS}}$ as a quantity which localizes on right-moving chiral primaries, but with suitable definition it would give the same function as $Z_{\text{NS},R}$ above. So, while the AdS vacuum appears in the (NS, NS) sector, we will focus on $Z_{\text{NS},R}$ when stating our bounds in Sect. 3.

In the cases of interest to us, there is no anti-holomorphic dependence on $\bar{\tau}$ due to the $(-1)^{\bar{F}}$ insertion, and the elliptic genus is a purely holomorphic function of τ . In fact, much more is true. Using standard arguments one can show that the elliptic genus of an SCFT defined above in (2.5) transforms nicely under the group $\mathbb{Z}^2 \rtimes \text{SL}(2, \mathbb{Z})$. In particular, it is a so-called weak Jacobi form of weight 0 and index $c/6$, defined below. For instance, supersymmetric sigma models for Calabi–Yau target spaces of complex dimension $2m$ have elliptic genera that are weight 0 weak Jacobi form of index m . For the rest of this paper, we will be considering SCFTs with $m \in \mathbb{Z}$, or equivalently c divisible by 6.

Consider a holomorphic function $\phi(\tau, z)$ on $\mathbb{H} \times \mathbb{C}$ which satisfies the conditions

$$\phi \left(\frac{a\tau + b}{c\tau + d}, \frac{z}{c\tau + d} \right) = (c\tau + d)^w e^{2\pi i m \frac{cz^2}{c\tau + d}} \phi(\tau, z), \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z}) \tag{2.7}$$

$$\phi(\tau, z + \ell\tau + \ell') = e^{-2\pi i m(\ell^2\tau + 2\ell z)} \phi(\tau, z), \quad \ell, \ell' \in \mathbb{Z}. \tag{2.8}$$

In the present context, (2.8) can be understood in terms of the spectral flow symmetry in the presence of an $\mathcal{N} \geq 2$ superconformal symmetry.

The invariance $\phi(\tau, z) = \phi(\tau + 1, z) = \phi(\tau, z + 1)$ implies a Fourier expansion

$$\phi(\tau, z) = \sum_{n, \ell \in \mathbb{Z}} c(n, \ell) q^n y^\ell, \tag{2.9}$$

and the transformation under $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \in \text{SL}_2(\mathbb{Z})$ shows

$$c(n, \ell) = (-1)^w c(n, -\ell). \tag{2.10}$$

The function $\phi(\tau, z)$ is called a weak Jacobi form of index $m \in \mathbb{Z}$ and weight w if its Fourier coefficients $c(n, \ell)$ vanish for $n < 0$. Moreover, the elliptic transformation (2.8) can be used to show that the coefficients

$$c(n, \ell) = C_r(D(n, \ell)) \tag{2.11}$$

depend only on the so-called discriminant

$$D(n, \ell) := \ell^2 - 4mn \tag{2.12}$$

and $r = \ell \pmod{2m}$. Note that $D(n, \ell)$ is the negative of the polarity, defined in [12] as $4mn - \ell^2$.

Combining the above, we see that a Jacobi form admits the expansion

$$\phi(\tau, z) = \sum_{r \in \mathbb{Z}/2m\mathbb{Z}} h_{m,r}(\tau) \theta_{m,r}(\tau, z) \tag{2.13}$$

in terms of the index m theta functions,

$$\theta_{m,r}(\tau, z) = \sum_{k=r \pmod{2m}} q^{k^2/4m} y^k. \tag{2.14}$$

Both $h_{m,r}$ and $\theta_{m,r}$ only depend on the value of r modulo $2m$. However, for some later manipulations, we should note that it is sometimes useful to choose the explicit fundamental domain $-m < r \leq m$ for the shift symmetry in r . When $|r| \leq m$ we can write:

$$h_{m,r}(\tau) = (-1)^w h_{m,-r}(\tau) = \sum_{n \geq 0} c(n, r) q^{-D(n,r)/4m}. \tag{2.15}$$

The vector-valued functions $\theta_{m,r}(\tau, z)$ transform as

$$\theta_m \left(-\frac{1}{\tau}, -\frac{z}{\tau} \right) = \sqrt{-i\tau} e^{\frac{2\pi i m z^2}{\tau}} \mathcal{S} \theta_m(\tau, z), \tag{2.16}$$

$$\theta_m(\tau + 1, z) = \mathcal{T} \theta_m(\tau, z), \tag{2.17}$$

where \mathcal{S}, \mathcal{T} are the $2m \times 2m$ unitary matrices with entries

$$\mathcal{S}_{rr'} = \frac{1}{\sqrt{2m}} e^{\frac{\pi i r r'}{m}}, \tag{2.18}$$

$$\mathcal{T}_{rr'} = e^{\frac{\pi i r^2}{2m}} \delta_{r,r'}. \tag{2.19}$$

From this we see that $h = (h_{m,r})$ is a $2m$ -component vector transforming as a weight $w - 1/2$ modular form for $SL_2(\mathbb{Z})$.

In particular, an elliptic genus (with $w = 0$) of a theory with central charge $c = 6m$ can be written as

$$Z_{R,R}(\tau, z) = \sum_{r \in \mathbb{Z}/2m\mathbb{Z}} Z_r(\tau) \theta_{m,r}(\tau, z). \tag{2.20}$$

We have written $Z_r(\tau)$ for $h_{m,r}(\tau)$ in this expression. Thus $Z_r(\tau)$ only depends on r modulo $2m$, but again, when $|r| \leq m$ it is useful to expand:

$$Z_r(\tau) = Z_{-r}(\tau) = \sum_{n \geq 0} c(n, r) q^{n - \frac{r^2}{4m}} \tag{2.21}$$

The function $Z_r(\tau)$ can be thought of as the elliptic genus of the r th superselection sector corresponding to the eigenvalue of $J_0 = r \pmod{2m}$. From the CFT point of view, the $r \sim r + 2m$ identification can be understood in terms of the spectral flow symmetry of the superconformal algebra. When there is a gravity dual the $r \rightarrow r + 2m$ transformation corresponds, from the bulk viewpoint, to a large gauge transformation of a gauge field holographically dual to the $U(1)_R$.

Since the Fourier coefficients of a weak Jacobi form have to satisfy

$$c(n, \ell) = 0 \quad \text{for all } n < 0, \tag{2.22}$$

(which can be thought of as unitarity of the CFT), this leaves open the possibility to have “polar terms” $c(n, \ell)q^n y^\ell$ with

$$m^2 \geq D(n, \ell) > 0, \quad n \geq 0$$

in an index m weak Jacobi form. These are called polar terms because they are precisely the terms in the q -series of $Z_r(\tau)$ that have exponential growth when approaching the cusp $\tau \rightarrow i\infty$. The finite set of independent coefficients of the polar terms in the elliptic genus will play a crucial role in what follows. In what follows, we will denote by ϕ_P the sum of all the polar terms in the elliptic genus.

Importantly, the full set of Fourier coefficients of a weak Jacobi form can be reconstructed from just the polar part, ϕ_P . This can be understood through the fact that there are no non-vanishing negative weight modular forms at any level. For discussions of this in related contexts, see [12–14]. Let us denote by V_m the space of possible polar polynomials (without requiring that they correspond to the polar part of a bona fide weak Jacobi form). Given the symmetries of the $c(n, \ell)$, V_m is spanned by $q^n y^\ell$ in the region $\mathcal{P}^{(m)}$:

$$\mathcal{P}^{(m)} = \{(\ell, n) : 1 \leq \ell \leq m, 0 \leq n, D(n, \ell) > 0\}. \tag{2.23}$$

By a standard counting of the number of lattice points underneath the parabola $4mn - \ell^2 = 0$ in the ℓ, n plane [12], one can give a formula for the dimension of the vector space of polar parts $P(m) = \dim(V_m)$:

$$P(m) = \sum_{\ell=1}^m \left\lceil \frac{\ell^2}{4m} \right\rceil. \tag{2.24}$$

In this note, where we work at leading order in large m , we will only need the leading behavior of the sum (2.24); this is determined by the elementary formula $\sum_{\ell=1}^m \ell^2 = \frac{1}{3}m^3 + \frac{1}{2}m^2 + \frac{1}{6}m$ to be

$$P(m) = \frac{1}{12}m^2 + O(m), \quad m \gg 1. \tag{2.25}$$

Because we are working at leading order at large m (large central charge), we will not need to use the subleading corrections to (2.25) (see for instance [9] and [12]). Neither will we need to deal with the important subtlety that not all vectors in V_m actually correspond to a weak Jacobi form. Denoting the space of weak Jacobi forms of weight 0 and index m as $\tilde{J}_{0,m}$, in fact one has $\dim(\tilde{J}_{0,m}) - P(m) = O(m)$. These facts would become important if one were to extend our results to the next order in a $1/m$ expansion.

3. Gravity Constraints and Phase Structure

We will now derive a constraint on the polar coefficients of an SCFT as follows. The polar coefficients determine the elliptic genus, and we will require that the genus matches the expected Bekenstein–Hawking entropy of black holes in the high-energy regime. Happily, we will find that a second (a priori independent)

requirement of the existence of a sharp Hawking–Page transition at the critical temperature $\beta = 2\pi$ gives the same constraint on the coefficients.

More precisely, we will be considering infinite sequences of CFTs going off to large central charge, and we will bound the asymptotic behavior of physical observables in such sequences as $m \rightarrow \infty$. (One familiar example that can be taken as representative of what we have in mind is the sequence of σ -models with targets $\text{Sym}^m(K3)$.) Simple physical considerations will lead us to propose certain constraints on the growth of the polar coefficients at large m in the related families of elliptic genera.

Now, there are precise mathematical statements on the behaviors of coefficients of large powers of q in modular forms. For instance, there are theorems proving that for a generic holomorphic modular form $f = \sum_n c_n q^n$ of fixed weight k , c_n grows as $O(n^{k-1})$ at large n , while for a cusp form, the coefficients are of $O(n^{k/2})$.

Note that our growth estimates are rather different in nature from those of the previous paragraph. Our estimates will be *physically* motivated by known facts about corrections to Einstein gravity in the expansion in energy divided by M_{Planck} . We are proposing a mathematical criterion, motivated by physics, that would allow one to check whether a given sequence of CFTs can possibly have a weakly coupled gravity dual. This could equally well be viewed as a mathematical conjecture about the families of modular forms arising in sequences with gravity duals.

Our eventual criterion will be derived by considering the free energies F_m of the CFTs in this family. The free energy in these theories, as $m \rightarrow \infty$, gives a function with a sharp first-order phase transition at $\beta = 2\pi$. This is the physical phenomenon of the Hawking–Page transition [4]. (Sharp roughly because, in microscopic examples of $\text{AdS}_3/\text{CFT}_2$, semi-classical configurations of winding strings can condense and lower the free energy precisely at 2π , yielding the transition—see [§5.3.2, [15]]). Similarly, when we state physically motivated criteria about the free energies of our sequences of theories, we will be making statements about the sequence F_m and assuming that the limit as $m \rightarrow \infty$ of $\frac{1}{m}F_m$ exists as a piece-wise differentiable function with discontinuous first-derivative at $\beta = 2\pi$.

3.1. A Bekenstein–Hawking Bound on the Elliptic Genus

Suppose that ϕ is the elliptic genus of a superconformal field theory with a large radius gravitational dual. Define the “reduced mass” of a particle state in the dual gravity picture to be the eigenvalue of

$$L_0^{\text{red}} = L_0 - \frac{1}{4m}J_0^2 - \frac{m}{4}, \quad (3.1)$$

namely the quantity $-D(n, \ell)/4m$ for the term $q^n y^\ell$ in the elliptic genus. Define E^{red} to be the eigenvalue under L_0^{red} . Then:

- Classically, the states with $E^{\text{red}} > 0$ are black holes in AdS_3 . We will discuss their contribution to the supergravity computation of the elliptic genus in detail below.

- In contrast, in the gravitational computation of the elliptic genus, it is the states with $E^{\text{red}} < 0$ which contribute to the polar part of the supergravity partition function [13]. These are precisely the modes which are too light to form black holes in the bulk. These are the states which appear in ϕ_P .

We now present an argument that constrains the coefficients in ϕ_P using the supergravity estimate of the black hole contribution to the elliptic genus. We treat the elliptic genus as the grand canonical partition function

$$Z(\beta, \mu) = \sum_{\text{microstates}} e^{\beta(\mu Q - E)} = e^{-\beta F(\beta, \mu)}, \quad (3.2)$$

where $\tau = i\frac{\beta}{2\pi}$ and $z = -i\frac{\beta\mu}{2\pi}$ are the corresponding variables in the elliptic genus. In other words, we define

$$Z(\beta, \mu) = Z_{NS,R} \left(\tau = i\frac{\beta}{2\pi}, z = -i\frac{\beta\mu}{2\pi} \right). \quad (3.3)$$

To make contact with the usual thermodynamical analysis, we will require β and μ to be real numbers. Let us discuss the supergravity estimate for this in simple steps. See also, for instance, the nice discussions in [16, 17].

3.1.1. Uncharged BTZ. In calculating the elliptic genus for a 2d SCFT, we restrict to states that are ground states on the right-moving side, but with arbitrary L_0 . These correspond to extremal spinning black holes in the 3d bulk, with vanishing temperature $T = 0$.

We can calculate the entropy of these black holes using the standard properties of black hole thermodynamics [18]. We will work in units where $\ell_{\text{AdS}} = 1$. The inner and outer horizons coincide for the extremal geometries, and are located at

$$r_+ = r_- = 2\sqrt{GM}. \quad (3.4)$$

The entropy is given by

$$S = \frac{\pi r_+}{2G}. \quad (3.5)$$

Finally, the central charge of the Brown–Henneaux Virasoro algebra is related to G by

$$c = \frac{3}{2G}. \quad (3.6)$$

Combining, we get

$$S = 2\pi\sqrt{\frac{cM}{6}}. \quad (3.7)$$

If we were to include Planck-suppressed corrections to the black hole entropy, we expect no fractional powers of M_{Planck} to appear in the corrected formula, but corrections which involve $\log(M_{\text{Planck}})$ can appear. This translates into $O(\log c)$ corrections, but no power-law in c corrections, to the entropy.

The black hole mass M is identified with the eigenvalue of $L_0 - \frac{c}{24}$, which we will denote as n . This means that the degeneracy of states of the elliptic genus c_n goes as

$$c_n = e^{2\pi\sqrt{\frac{cn}{6}} + O(\log n)}. \tag{3.8}$$

This is the familiar Cardy-like growth. As we are interested in studying families of CFTs asymptoting to the large central charge limit, we would like to know about the behavior at fixed n as $c \rightarrow \infty$. For this purpose, the more informative expression would be

$$c_n = e^{2\pi\sqrt{\frac{cn}{6}} + O(\log c)}. \tag{3.9}$$

As an aside, let us discuss the validity of the above equation. The above derivation of the black hole contribution to the partition function is valid whenever the radius of the black hole is large in Planck units. The first BTZ black hole appears at a mass $\sim M_{\text{Planck}}$, and we see from (3.4) that its radius will already be quite large—of order ℓ_{AdS} , or $O(c)$ in Planck units. We then expect the semi-classical entropy formula to be valid for even very light black holes at large c . This is one way to understand the characteristic Cardy-like growth of the number of states of CFTs with gravity duals, even outside the usual range of validity of the Cardy formula that is guaranteed by modular invariance alone.

Writing the elliptic genus now as

$$Z(\tau) = \int dn e^{2\pi\sqrt{\frac{cn}{6}}} e^{2\pi i\tau n} \tag{3.10}$$

we can ask the question: at fixed τ (where $\frac{i}{2\pi\tau} = \frac{1}{\beta}$ is the formal “temperature” variable; not to be confused with the temperature of the black hole, which is zero), what value of n dominates the sum? This is solved using standard saddle point approximation methods. The derivative of $e^{2\pi\sqrt{\frac{cn}{6}}} e^{2\pi i\tau n}$ vanishes when

$$2\pi i\tau = -\pi\sqrt{\frac{c}{6n}} \tag{3.11}$$

or equivalently

$$\beta = \pi\sqrt{\frac{c}{6n}}. \tag{3.12}$$

Thus we get

$$n = \pi^2 \frac{m}{\beta^2}. \tag{3.13}$$

Using the famous relation $F = E - S/\beta$ we therefore get

$$F = -\pi^2 \frac{m}{\beta^2} + O(\log m). \tag{3.14}$$

We were careful to write β here to distinguish from the physical temperature of the extremal black holes contributing to the genus. (While the torus partition function at a given τ would correspond to a thermal ensemble, the elliptic genus is only counting extremal states and the temperature represented by $\text{Im}(\tau)$ is fictitious.)

3.1.2. Adding Wilson Lines. Now we turn to the elliptic genus, a refinement of the above discussion which keeps track of $U(1)$ charge.

In the bulk, the existence of the $U(1)_R$ symmetry of the dual (2,2) SCFT is manifested in the presence of Chern–Simons gauge fields. First, let us discuss the expected effect heuristically. By adding a $U(1)$ Chern–Simons gauge interaction at level k , we add to the action the following boundary term

$$S_{\text{gauge}}^{\text{bdry}} = -\frac{k}{16\pi} \int_{\partial\text{AdS}} d^2x \sqrt{g} g^{\alpha\beta} A_\alpha A_\beta. \tag{3.15}$$

For a BTZ black hole, the angular direction in the 2d spatial manifold (which we shall call the ϕ direction) is non-contractible, so we allow A_ϕ to be nonzero.

We thus shift the action by a term proportional to A^2 . This will add a term that goes as μ^2 to the free energy so we will get something like

$$F \sim \frac{m}{\beta^2} + k\mu^2. \tag{3.16}$$

Finally, for a (2,2) SCFT with k determined by the central charge and hence the index m , we will have

$$F \sim \frac{m}{\beta^2} + m\mu^2. \tag{3.17}$$

Now, let us be more explicit. The entropy of the black holes we are considering is given, in general, by [19]

$$\begin{aligned} S &= 2\pi\sqrt{m}\sqrt{n - \frac{\ell^2}{4m}} \\ &= \pi\sqrt{-D(n, \ell)}, \end{aligned} \tag{3.18}$$

where n is the eigenvalue under $L_0 - \frac{c}{24}$, and ℓ is the J_0 eigenvalue.

Now, again, we write the degeneracy

$$c(n, \ell) = e^{2\pi\sqrt{m}\sqrt{n - \frac{\ell^2}{4m}} + O(\log(n - \frac{\ell^2}{4m}))}, \tag{3.19}$$

or following the analogous discussion above

$$c(n, \ell) = e^{2\pi\sqrt{m}\sqrt{n - \frac{\ell^2}{4m}} + O(\log m)}, \tag{3.20}$$

and the elliptic genus can be approximated as

$$Z_{\text{NS},R}(\tau, z) = \int dn \int d\ell e^{2\pi i\tau n} e^{2\pi iz\ell} e^{2\pi\sqrt{m}\sqrt{n - \frac{\ell^2}{4m}}}. \tag{3.21}$$

This has a saddle when

$$\begin{aligned} \tau &= \frac{im}{\sqrt{4mn - \ell^2}} \\ z &= \frac{-i\ell}{2\sqrt{4mn - \ell^2}}. \end{aligned} \tag{3.22}$$

Rewriting, the dominant saddle occurs at

$$\begin{aligned} n &= m \left(\frac{\pi^2}{\beta^2} + \mu^2 \right) \\ \ell &= 2m\mu. \end{aligned} \tag{3.23}$$

Thus, we get the free energy as

$$F = -m \frac{\pi^2}{\beta^2} - m\mu^2 + O(\log m). \tag{3.24}$$

Identifying this free energy with $-\frac{1}{\beta} \log Z$ gives us the behavior of the elliptic genus. However, we need to be sure that the supergravity derivation is valid—i.e., that the configurations we included correspond to reliable and dominant saddle points. Reliability follows if the black hole is large in Planck units, which works for any $E^{\text{red}} > 0$ at large c . We also require that the black hole saddle be the dominant one. This will be true for any $\beta < 2\pi$ at very large m . For $\beta > 2\pi$, instead the “gas of gravitons” dominates, and (3.24) is not the appropriate expression for the free energy. Finally, in a tiny neighborhood of $\beta = 2\pi$, the free energy crosses from the value for the gas of gravitons to the value characteristic of black holes above; this is a regime where “enigma black holes” play an important role, and cannot be characterized in a universal way. In known microscopic examples of $\text{AdS}_3/\text{CFT}_2$, these are small black holes (localized on the transverse sphere) of negative specific heat (see e.g., [20, 21] for discussions).

Next we will derive constraints on the low-temperature expansion—and in particular the polar coefficients—from these results of black hole thermodynamics.

3.1.3. Bounds on Polar Coefficients. After these physical preliminaries, we are ready to derive the main result of this paper. This result will follow (given appropriate physical assumptions) by combining modular invariance with the physical requirement that $Z(\beta, \mu)$ has large m asymptotics given by

$$\log Z(\beta, \mu) = m \left(\frac{\pi^2}{\beta} + \beta\mu^2 \right) + O(\log m), \tag{3.25}$$

for all real (β, μ) such that $0 < \beta < 2\pi$. Recall from Eq. (3.3) that $Z(\beta, \mu)$ is just the elliptic genus $Z_{\text{NS},R}(\tau, z)$ evaluated for $\tau = i\beta/2\pi$ and $z = -i\beta\mu/2\pi$.

Now we write out the modular property:

$$Z_{\text{NS},R}(\tau, z) = (-1)^m e^{-\frac{2\pi i m z^2}{\tau}} Z_{\text{NS},R} \left(-\frac{1}{\tau}, -\frac{z}{\tau} \right). \tag{3.26}$$

We make a few elementary manipulations:

$$\begin{aligned} Z_{\text{NS},R}(\tau, z) &= e^{\frac{2\pi i \tau m}{4}} e^{2\pi i z m} Z_{R,R} \left(\tau, z + \frac{\tau + 1}{2} \right) \\ &= e^{\frac{\pi i \tau m}{2}} e^{2\pi i z m} \sum_{r \in \mathbb{Z}/2m\mathbb{Z}} Z_r(\tau) \theta_{m,r} \left(\tau, z + \frac{\tau + 1}{2} \right) \\ &= e^{\frac{\pi i \tau m}{2}} e^{2\pi i z m} \sum_{r \in \mathbb{Z}/2m\mathbb{Z}} \sum_{\substack{D \leq r^2 \\ D \equiv r^2 \pmod{4m}}} C_r(D) e^{-\frac{2\pi i \tau D}{4m}} \\ &\quad \times \sum_{k=r \pmod{2m}} e^{\frac{2\pi i k^2 \tau}{4m}} e^{2\pi i z k} (-1)^k e^{i\pi \tau k} \end{aligned}$$

$$\begin{aligned}
 &= e^{-\frac{\pi i \tau m}{2}} e^{2\pi i z m} \sum_{r \in \mathbb{Z}/2m\mathbb{Z}} \sum_{\substack{D \leq r^2 \\ D=r^2 \pmod{4m}}} \\
 &\quad \times \sum_{k=r \pmod{2m}} C_r(D) e^{\frac{2\pi i \tau}{4m}(-D+m^2+(k+m)^2)} e^{2\pi i z k} (-1)^k. \tag{3.27}
 \end{aligned}$$

Combining (3.26) and (3.27), we get

$$\begin{aligned}
 Z_{NS,R}(\tau, z) &= e^{-\frac{2\pi i m z^2}{\tau}} e^{\frac{i\pi m}{2\tau}} e^{-\frac{2\pi i z m}{\tau}} \sum_{r \in \mathbb{Z}/2m\mathbb{Z}} \sum_{\substack{D \leq r^2 \\ D=r^2 \pmod{4m}}} \\
 &\quad \times \sum_{k=r \pmod{2m}} C_r(D) e^{\frac{2\pi i}{4m\tau}(D-m^2-(k+m)^2)} e^{-\frac{2\pi i z k}{\tau}} (-1)^{k+m} \\
 &= e^{m\beta\mu^2 + \frac{m\pi^2}{\beta}} \sum_{r \in \mathbb{Z}/2m\mathbb{Z}} \sum_{\substack{D \leq r^2 \\ D=r^2 \pmod{4m}}} \\
 &\quad \times \sum_{k=r \pmod{2m}} C_r(D) e^{\frac{\pi^2}{m\beta}(D-m^2-(k+m)^2)} e^{2\pi i(k+m)(\mu+\frac{1}{2})} \tag{3.28}
 \end{aligned}$$

where in the last line we have used the substitutions $\tau = \frac{i\beta}{2\pi}$ and $z = -\frac{i\beta\mu}{2\pi}$.

Note that the prefactor in front of the sum in Eq. (3.28) gives the right-hand side of Eq. (3.25). Therefore

$$\begin{aligned}
 &\log \left(\sum_{r \in \mathbb{Z}/2m\mathbb{Z}} \sum_{\substack{D \leq r^2 \\ D=r^2 \pmod{4m}}} \sum_{k=r \pmod{2m}} C_r(D) e^{\frac{\pi^2}{m\beta}(D-m^2-(k+m)^2)} e^{2\pi i(k+m)(\mu+\frac{1}{2})} \right) \\
 &\quad \sim O(\log(m)). \tag{3.29}
 \end{aligned}$$

In order to turn this into a more useful statement we next introduce another physically motivated hypothesis—the “non-cancellation hypothesis”. This hypothesis states that the leading order large m asymptotics is not affected if we replace the terms in the expansion of $Z_{NS,R}$ above by their *absolute values*.¹ Given the noncancellation hypothesis none of the terms in the sum can get large, and hence we arrive at the necessary condition:

$$\log \left(|C_r(D)| e^{\frac{\pi^2}{m\beta}(D-m^2-(k+m)^2)} \right) = O(\log m) \quad \text{for all } \beta < 2\pi \text{ and } k=r \pmod{2m}. \tag{3.30}$$

The strongest bound is obtained by taking the limit as β increases to 2π from below, yielding:

$$|C_r(D)| \leq e^{\frac{2\pi}{4m}(m^2 - D + \min\{(k+m)^2 | k=r \pmod{2m}\})} + O(\log m). \tag{3.31}$$

¹ Since $Z_{NS,R}$ is modular this again can only be valid in a distinguished set of expansions around cusps, and we take it to apply to the expansion in (3.28).

We can write the bound simply in terms of coefficients $c(n, \ell)$ where $0 \leq \ell \leq m$; the rest of the coefficients will be determined from this subset via spectral flow and reflection of ℓ . We then get the bound

$$|c(n, \ell)| \leq e^{2\pi(n + \frac{m}{2} - \frac{|\ell|}{2}) + O(\log m)}. \quad (3.32)$$

Put differently, $|e^{-2\pi(n + \frac{m}{2} - \frac{|\ell|}{2})} c(n, \ell)|$ can grow at most as a power of m for $m \rightarrow \infty$. In addition to these conditions, the bound should not be saturated by an exponentially large number of states. Note that in the special case $\ell = 0$, our bound (3.32) coincides with the result of [2].

We conclude with a few remarks.

1. To be fastidious, the bound (3.32) applies to any family $\mathcal{C}^{(m)}$ of CFTs with a weakly coupled gravity dual, together with a sequence $(n(m), \ell(m))$ of lattice points such that the sequence of elliptic genus coefficients $c(n(m), \ell(m); \mathcal{C}^{(m)})$ has well-defined large m asymptotics.
2. The $O(\log m)$ error term in the exponent can be understood in various ways. Perhaps the most enlightening physically is that it can be directly connected (via modularity) to the M_{Planck} suppressed corrections to the black hole entropy in the $\beta < 2\pi$ regime.
3. Note that the bound is already nontrivial for the coefficient $c(0, m)$ of the extreme polar term with $(n, \ell) = (0, \pm m)$. Under spectral flow, the states contributing to this degeneracy correspond to the unique NS-sector vacuum on the left tensored with one of the Ramond sector ground states on the right. We will see that already the bound on the extreme polar states is useful.
4. Notice that in (3.18), we have only written a formula for the entropy in the stable black hole region $E^{\text{red}} > \frac{c}{24}$. This follows because our saddle point approximation is only self-consistent when $\beta < 2\pi$ in this range of energies. While it may seem naively that the large c behavior of the free energy would guarantee this formula also for $0 < E^{\text{red}} < \frac{c}{24}$, this is not the case. Because there is a jump of $O(c)$ in the energy in a small neighborhood of $\beta = 2\pi$, in this window $O(1)$ contributions to the free energy (which we've neglected in the large c limit) could lead to significant changes in E^{red} ; our formula for $S(E^{\text{red}})$ is then unreliable. It becomes reliable once one reaches the stable range of energies $E^{\text{red}} > \frac{c}{24}$. For further discussion of this issue, see [2] as well as [20, 21].

3.2. On the Hawking–Page Transition

In what follows, we will present an alternate derivation of (3.32) by insisting on a sharp Hawking–Page phase transition near $\beta = 2\pi$ (in the limit of large central charge) in the NS-R sector. The sharp transition is not a surprise. It is expected from general properties of the $\text{AdS}_3/\text{CFT}_2$ duality (and in particular, from the existence of light multiply-wound strings which can lower the free energy once $\beta < 2\pi$, in known microscopic examples [§5.3.2, [15]]).

Recall that the NS sector elliptic genus has a q -expansion of the form

$$Z_{\text{NS},+}(\tau) = q^{\frac{m}{4}} Z_{\text{RR}}\left(\tau, \frac{\tau+1}{2}\right) = \sum_{n,\ell} (-1)^\ell c(n, \ell) q^{\frac{m}{4}+n+\frac{\ell}{2}}. \tag{3.33}$$

From the modular properties of $Z_{\text{RR}}(\tau, z)$ we see that $Z_{\text{NS},+}(\tau)$ is invariant under the group

$$\Gamma_\theta = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z}) \mid c-d \equiv a-b \equiv 1 \pmod{2} \right\} \tag{3.34}$$

which is conjugate to the Hecke congruence group $\Gamma_0(2)$.

Clearly, it satisfies at the lowest temperatures

$$\log Z_{\text{NS},+}\left(\tau = i\frac{\beta}{2\pi}\right) = \frac{c}{24}\beta, \quad \beta \gg 2\pi. \tag{3.35}$$

To have a phase dominated by the ground state until temperatures parametrically close to $\beta = 2\pi$ at large central charge $c = 6m$, one requires:

$$\log Z_{\text{NS},+}\left(\tau = i\frac{\beta}{2\pi}\right) = \frac{c}{24}\beta + O(\log c), \quad \beta > 2\pi. \tag{3.36}$$

Again, this can be viewed as an asymptotic condition on a family of CFTs which has a weakly curved gravity dual at large m : the limit as $m \rightarrow \infty$ of $\frac{1}{m} \log Z_{\text{NS},+}$ for any $\beta > 2\pi$ exists and asymptotes to $\frac{1}{4}\beta$.

The size of the sub-leading terms in (3.36) requires some discussion. In fact, just for the purpose of having a phase transition at $\beta = 2\pi$ in the large c limit, it is possible to relax the condition of strict ground state dominance and to allow $\log Z_{\text{NS},+}(\beta) = \frac{c}{24}\beta + O(c^{1-\delta})$ for some $\delta > 0$, instead of restricting to $O(\log c)$. As noted before, however, in the large temperature regime this would imply corrections to the Bekenstein–Hawking entropy suppressed by fractional powers of M_{Planck} , which are not expected. On the other hand, logarithmic corrections are expected. This suggests one should set $\delta = 1$. In any case, we shall not pursue the slight generalization to $\delta \neq 0$ in the present paper—the requisite modification of the analysis can be implemented in a relatively straightforward way.

A sufficient condition for (3.36) to be true is that $|c(n, \ell)q^{\frac{m}{4}+n+\frac{\ell}{2}}| \leq e^{m\beta/4}$ for a number of terms which grows at most polynomially in m . If we invoke the noncancellation hypothesis, we can also say that a necessary condition is:

$$|c(n, \ell)| \leq e^{2\pi(n - \frac{|\ell|}{2} + \frac{m}{2}) + O(\log m)}, \tag{3.37}$$

If we combine this statement with the spectral flow property $c(n, \ell) = c(n + s\ell + ms^2, \ell + 2sm)$ for all integers s we can get the best bound by minimizing with respect to s , subject to the condition that s is integer. Combining with reflection invariance on ℓ it is not difficult to show then that the best bound is

$$|c(n, \ell)| \leq e^{2\pi(n_0 - \frac{|\ell_0|}{2} + \frac{m}{2}) + O(\log m)}, \tag{3.38}$$

where (n, ℓ) is related to (n_0, ℓ_0) by spectral flow and reflection and $0 \leq \ell_0 \leq m$. This is the same condition we have derived to reproduce Bekenstein–Hawking entropy (3.32).




FIGURE 1. The tessellation by Γ_θ and its sub-tessellation by $\Gamma_\infty \setminus \Gamma_\theta$. The thick lines are where phase transitions in supergravity can occur

The above phase transition corresponds to moving between $\text{Im}(\tau) = 1 - \epsilon$ and $\text{Im}(\tau) = 1 + \epsilon$ with $\text{Re}(\tau) = 0$ between two specific copies of the fundamental domain of Γ_θ ; see Fig. 1. In the Euclidean signature, other saddle points corresponding to analytic continuation of the BTZ black holes are also believed to be relevant [13, 22], and one is led to a stronger prediction for a phase diagram requiring an infinite number of different phases corresponding to pairs (c, d) of co-prime integers with $c \geq 0$, $c - d \equiv 1 \pmod{2}$ (see [13] and [23], §7.3).²

One should then obtain a phase structure which divides the upper-half plane into regions dominated by the various saddle points labeled by different values of (c, d) . This corresponds to a tessellation of the upper-half plane by $\Gamma_\infty \setminus \Gamma_\theta$ where Γ_∞ is the group generated by T^2 , coinciding with the intersection of Γ_θ and $\langle T \rangle$. In the above sentence, we have used the definition $T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \in PSL_2(\mathbb{Z})$ and $\langle T \rangle = \{T^n, n \in \mathbb{Z}\}$. This tessellation is drawn in Fig. 1 with the thick lines. We discuss the derivation of this phase diagram in detail in Appendix A, and show that in each region, one has a phase transition at the thick line in Fig. 1 which is similar in nature to our transition between thermal AdS dominance and the black hole regime.

4. Examples

In this section, we discuss how the elliptic genera of various simple CFTs— σ -models with targets $\text{Sym}^N(K3)$, product manifolds $(K3)^N$, or Calabi–Yau hypersurfaces up to relatively high dimension d —fare against the bound. Somewhat unsurprisingly, the first class of theories passes the bound while the others

² Reference [13] erroneously claimed the phase diagram would be invariant under $PSL(2, \mathbb{Z})$. However the argument given there is easily corrected, and it predicts a phase diagram invariant under $\Gamma_0(2)$ for the NS-sector genus considered there. For further discussion see Appendix A.

fail dramatically, exhibiting far too rapid a growth in polar coefficients [24]. We close with a discussion of $\text{Sym}^N(\mathcal{M})$, with \mathcal{M} the Monster CFT of Frenkel–Lepowsky–Meurman. This example proves a useful foil in contrasting theories with low energy supergravity vs low energy string duals.

4.1. $\text{Sym}^N(K3)$

The first example is one which we expect to satisfy the bound, and serves as a test of the bound. A system which historically played an important role in the development of the AdS/CFT correspondence was the D1–D5 system on $K3$ [10], and the duality between the σ -model with target space $(K3)^N/S_N$ and supergravity in AdS_3 was one of the first examples of $\text{AdS}_3/\text{CFT}_2$ duality [1]. See also [25] for a more detailed analysis.

The elliptic genus of the symmetric product CFT was discussed extensively in [26]. One can define a generating function for elliptic genera

$$\mathcal{Z}_X(\sigma, \tau, z) = \sum_{N \geq 0} p^N Z_{R,R}(\text{Sym}^N(X); \tau, z), \quad p = e^{2\pi i \sigma}, \quad (4.1)$$

which is given by [26] as

$$\mathcal{Z}_X(\sigma, \tau, z) = \prod_{n > 0, n' \geq 0, l} \frac{1}{(1 - p^n q^{n'} y^l)^{c_X(nn', l)}}. \quad (4.2)$$

The coefficients $c_X(n, l)$ are defined as the Fourier coefficients of the original CFT X ,

$$Z_{R,R}(X; \tau, z) = \sum_{n \geq 0, l} c_X(n, l) q^n y^l. \quad (4.3)$$

If we are interested in calculating the $O(q^0)$ piece of the elliptic genus of $\text{Sym}^N(X)$, we can set $n' = 0$ in (4.2), giving

$$\lim_{\tau \rightarrow i\infty} \mathcal{Z}_X(\sigma, \tau, z) = \prod_{n > 0, l} \frac{1}{(1 - p^n y^l)^{c_X(0, l)}}. \quad (4.4)$$

When X is the sigma model with Calabi–Yau target space (which we also call X), the above is, up to simple factors, the generating function for the χ_y -genus of $\text{Sym}^N(X)$.

The most polar term of $\text{Sym}^N(X)$ is given by y^{mN} where $m = \dim_{\mathbb{C}} X/2$ is the index of the elliptic genus of X . This is the coefficient of $y^{mN} p^N$ in (4.2), which only receives contributions from

$$\frac{1}{(1 - py^m)^{c_X(0, m)}}. \quad (4.5)$$

By calculating the coefficient of $p^N y^{Nm}$ in (4.5) we get

$$c_{\text{Sym}^N X}(0, Nm) = \binom{c_X(0, m) + N - 1}{c_X(0, m) - 1}, \quad (4.6)$$

a polynomial of degree $c_X(0, m) - 1$ in N and therefore allowed by the bound (3.32).

In order to find the subleading polar piece for $\text{Sym}^N(X)$, we calculate the coefficient of the term $p^N y^{Nm-1}$ in (4.2). This has contributions from

$$\frac{1}{(1 - py^m)^{c_X(0,m)}} \frac{1}{(1 - py^{m-1})^{c_X(0,m-1)}} \frac{1}{(1 - p^2 y^m)^{c_X(0,m)}}. \tag{4.7}$$

The $p^N y^{Nm-1}$ term generically comes from multiplying a $p^{N-1} y^{m(N-1)}$ in the first term in (4.7) with a py^{m-1} from the second term. For the special case of $m = 1$, it can also come from multiplying a $p^{N-2} y^{m(N-2)}$ from the first term with a $p^2 y^m$ from the third term.

The coefficient of $p^{N-1} y^{m(N-1)}$ in the first term is $\binom{c_X(0,m)+N-2}{c_X(0,m)-1}$, and the coefficient of py^{m-1} in the second term is $c_X(0, m - 1)$. The coefficient of $p^{N-2} y^{m(N-2)}$ in the first term is $\binom{c_X(0,m)+N-3}{c_X(0,m)-1}$ and the coefficient of $p^2 y^m$ in the third term is $c_X(0, m)$. Thus the coefficient of the penultimate polar piece is given by

$$c_{\text{Sym}^N X}(0, Nm - 1) = \begin{cases} \binom{c_X(0,m)+N-2}{c_X(0,m)-1} c_X(0, m - 1), & \text{if } m > 1 \\ \binom{c_X(0,1)+N-2}{c_X(0,1)-1} c_X(0, 0) + \binom{c_X(0,1)+N-3}{c_X(0,1)-1} c_X(0, 1), & \text{if } m = 1. \end{cases} \tag{4.8}$$

Again, this exhibits polynomial growth in N and is allowed by (3.32). Any term a finite distance away from the most polar term (e.g., $y^{Nm-x} q^0$ for constant x) will grow as a polynomial in N of degree $c_X(0, m) - 1$.

For Calabi–Yau manifolds X with $\chi_0 = 2$, we have $c_X(0, m) = 2$ so the two most polar terms simplify to

$$c_{\text{Sym}^N X}(0, Nm) = N + 1$$

$$c_{\text{Sym}^N X}(0, Nm - 1) = \begin{cases} N c_X(0, m - 1), & \text{if } m > 1 \\ N c_X(0, 0) + 2(N - 1), & \text{if } m = 1. \end{cases} \tag{4.9}$$

For the special case of $X = K3$, we have $m = 1$ and $c_X(0, 0) = 20$, so the penultimate polar piece grows as $22N - 2$.

We can do a similar calculation to find the coefficient in front of y^{N-x} for $\text{Sym}^N(K3)$ with $x > 1$. We find the asymptotic large N value for the coefficient, presented in Table 1. In Fig. 2, we plot the polar coefficients of $\text{Sym}^{20}(K3)$ against the values allowed by the bound. Although some very polar terms exceed $e^{2\pi(n - \frac{|k|}{2} + \frac{m}{2})}$ in (3.32), the deviation is of the order $O(\log N)$ in the exponent, which is allowed in our analysis. See [27] for more information on the order $O(\log N)$ corrections. For terms with polarity close to zero, the $O(\log N)$ corrections are less important, and we see that the bound is subsaturated as expected.

The fact that $\text{Sym}^N(K3)$ satisfies our bounds is part of a more general story—in fact all symmetric products will satisfy this bound, regardless of the “seed” SCFT. This follows from the general class of arguments presented in [2, 24].

TABLE 1. Coefficient of y^{N-x} in $\text{Sym}^N(K3)$ elliptic genus at large N

x	Coefficient
0	$N + 1$
1	$22N - 2$
2	$277N - 323$
3	$2576N - 5752$
4	$19574N - 64474$
5	$128156N - 557524$
6	$746858N - 4035502$
7	$3959312N - 25550800$
8	$19391303N - 145452673$
9	$88757346N - 758554926$
10	$383059875N - 3673549725$
11	$1569800280N - 16690133400$
12	$6143337474N - 71708443374$
13	$23066290212N - 293213888652$
14	$83418524934N - 1146991810674$
15	$291538891984N - 4310932524176$
16	$987440609467N - 15624074962373$
17	$3249156243514N - 54773846935526$
18	$10408875430635N - 186236541847125$
19	$32525691116400N - 615565850482800$
20	$99302600734650N - 1981904206578750$

We later plot these values in Fig. 8




FIGURE 2. Here, we plot the polar coefficients of $\text{Sym}^{20}(K3)$ versus polarity, and also the coefficients allowed by the bounds. We see that at this value of c ($=120$), the bounds are satisfied by the symmetric product conformal field theory, after allowing minor shifts due to the $O(\log m)$ correction

4.2. Products of $K3$ (or, X^N)

The most obvious families of CFTs that “should” fail any reasonable test for having a (weakly coupled) gravity dual are given by tensor products of many small c CFTs. Here, as a foil to $\text{Sym}^N(K3)$, we describe the results for the product $(K3)^N$. Not surprisingly, it fails to satisfy the bounds. We will use the fact that




FIGURE 3. Here, we plot the polar coefficients of the product conformal field theory with target $K3^{20}$

$$Z_{R,R}^{(X^{\otimes N})}(\tau, z) = (Z_{R,R}^X(\tau, z))^N = \left(\sum_{n,\ell} c_X(n, \ell) q^n y^\ell \right)^N. \tag{4.10}$$

For concreteness, we look at the χ_y genus of $K3^N$. Since

$$Z_{R,R}^{(K3)}(\tau, z) = 2y^{-1} + 20 + 2y + O(q), \tag{4.11}$$

the $q^0 y^N$ term in the elliptic genus of $K3^N$ is given by

$$c_{K3^N}(0, N) = 2^N, \tag{4.12}$$

which violates the bound (3.32) of only polynomial growth for the most polar term.

To visualize the violation we plot the polar coefficients of $K3^{20}$ against the bound in Fig. 3. Note that the violations are not of the order $O(\log N)$, and (3.32) is clearly not satisfied.

We conclude with a few remarks about examples similar to the above:

1. We cannot rule out *all* product manifolds using this method. For instance, the elliptic genus of T^4 is zero, which means that products of T^4 will surely satisfy the bound, having a vanishing elliptic genus. The vanishing is due to cancellations arising from the $U(1)^4$ translation symmetry acting on $\text{Sym}^N(T^4)$. One could instead work with $\text{Sym}^N(T^4)/T^4$. In worldsheet terms, there are fermion zero modes due to the extra translation symmetry which must be saturated by the insertion of a suitable number of fermion currents. The relevant modification of the genus is worked out in [28]. It should be fairly straightforward to generalize our considerations to situations such as this where extra insertions are required to define a proper index.
2. Another simple example that violates the bound is the iterated symmetric product $\text{Sym}^{N_1}(\text{Sym}^{N_2}(K3))$. Taking, for simplicity, $N_1 = N_2 = N$, so $m = N^2$, the coefficient of the most polar term is $\binom{2N}{N} \sim \frac{1}{\sqrt{\pi N}} 4^N = \frac{1}{\pi^{1/2} m^{1/4}} 4\sqrt{m}$ for large m . Indeed, the iterated symmetric product is an example of the more general class of permutation orbifolds. It would

be interesting to explore the relation of our bound to the oligomorphic criterion of [29, 30].

4.3. Calabi–Yau Spaces of High Dimension

To provide a slightly more nontrivial test, we discuss the elliptic genera of Calabi–Yau sigma models with target spaces $X^{(d)}$ given by the hypersurfaces of degree $d + 2$ in $\mathbb{C}\mathbb{P}^{d+1}$, e.g.,

$$\sum_{i=0}^{d+1} z_i^{d+2} = 0. \tag{4.13}$$

We have chosen these as the simplest representatives among Calabi–Yau manifolds of dimension d ; as they are not expected to have any particularly special property uniformly with dimension, we suspect this choice is more or less representative of the results we could obtain by surveying a richer class of Calabi–Yau manifolds at each d . In any case we will settle with one Calabi–Yau per complex dimension. Since $m = d/2$, and we have been assuming m is integral, we restrict to even d .

The elliptic genus for these spaces is independent of moduli, and can be conveniently computed in the Landau–Ginzburg orbifold phase. This yields the formula [11]

$$Z_{R,R}^d(\tau, z) = \frac{1}{d+2} \sum_{k,\ell=0}^{d+1} y^{-\ell} \left(\frac{\theta_1\left(\tau, -\frac{d+1}{d+2}z + \frac{\ell}{d+2}\tau + \frac{k}{d+2}\right)}{\theta_1\left(\tau, \frac{1}{d+2}z + \frac{\ell}{d+2}\tau + \frac{k}{d+2}\right)} \right)^{d+2} \tag{4.14}$$

Many further facts about elliptic genera of Calabi–Yau spaces can be found in [31].

First, we discuss the explicit data. To facilitate this we computed all polar coefficients numerically for $d = 2, 4, \dots, 36$. Then, we provide a simple analytical proof of bound violation valid for all values of d (just following from the behavior of the subleading polar term).

Using (4.14) we can extract the polar coefficients explicitly for any given d . In Figs. 4, 5, and 6 we plot the coefficients of the polar pieces against polarity for Calabi–Yau 10-, 20-, and 36-fold, respectively. In Fig. 7, we plot the subleading polar coefficients of these Calabi–Yau spaces as a function of their dimension. In all cases, we see that the bounds are badly violated.

Numerics aside, it is easy to give a simple analytical argument proving that these Calabi–Yaus will violate the bound. Consider the subleading y^{m-1} polar piece of $Z_{RR}^{d=2m}$.

The coefficients $c_{X^{(d)}}(0, p)$ of the elliptic genera of Calabi–Yau spaces are determined simply by topological invariants:

$$c_{X^{(d)}}(0, m - i) = \sum_k (-1)^{i+k} h^{k,i}, \tag{4.15}$$

so the coefficient in front of y^{m-1} is

$$-\chi_1 = \sum_p -(-1)^p h^{1,p}. \tag{4.16}$$




FIGURE 4. Here, we plot the polar coefficients of $Z_{RR}^{d=10}$




FIGURE 5. Here, we plot the polar coefficients of $Z_{RR}^{d=20}$




FIGURE 6. Here, we plot the polar coefficients of $Z_{RR}^{d=36}$

We know $h^{1,d-1}$ is given by the number of complex structure parameters of the hypersurface, or

$$\begin{aligned}
 h^{1,d-1} &= \frac{(d+2) \times (d+3) \times \cdots \times (2d+3)}{1 \times 2 \times \cdots \times (d+2)} - (d+2)^2 \\
 &= \binom{2d+3}{d+2} - (d+2)^2.
 \end{aligned}
 \tag{4.17}$$




FIGURE 7. Here, we plot the subleading polar coefficients of the Calabi–Yau elliptic genera against the dimension

By a standard application of the Lefschetz hyperplane theorem, the remaining $h^{1,p}$ vanish except for $h^{1,1} = 1$. Thus we get (recall $d = 2m$ is even)

$$c_{X^{(d)}}(0, m - 1) = \binom{2d + 3}{d + 2} - (d + 2)^2 + 1. \tag{4.18}$$

And just as a check, for $d = 36$, we numerically get

$$\begin{aligned} c_{X^{(36)}}(0, 17) &= 3446310324346630675857 \\ &= \binom{75}{38} - 38^2 + 1, \end{aligned} \tag{4.19}$$

which matches the expectation on the nose.

Asymptotically, (4.18) goes as:

$$\begin{aligned} \log c_{X^{(d)}}(0, m - 1) &\sim \log(2d)! - 2 \log(d)! \\ &\sim 2d \log(2d) - 2d \log(d) \\ &= 2d \log 2 \end{aligned} \tag{4.20}$$

so

$$c_{X^{(d)}}(0, m - 1) \sim 2^{2d} = 2^{4m}. \tag{4.21}$$

To satisfy the bound, we need $c_X^{(d)}(0, m - 1)$ to grow at most polynomially with m when it in fact grows exponentially with m .

4.4. Enter the Monster

We now discuss a theory which passes our bounds but seemingly exhibits no supergravity regime—instead exhibiting a Hagedorn degeneracy of states already at low energies. We have benefited immensely in thinking about this theory from the unpublished work of Yin.

A $c = 24$ CFT with Monster symmetry was constructed many years ago by Frenkel, Lepowsky, and Meurman [32]. Let us call the non-chiral CFT with Monster symmetry \mathcal{M} . In this section, we wish to consider the symmetric products $\text{Sym}^N(\mathcal{M})$. As \mathcal{M} has no moduli, there is a unique partition function

canonically associated with this theory, and we will consider the chiral partition function instead of the elliptic genus in this section.

This requires a word of explanation. While the elliptic genera we've considered are related to non-chiral CFTs with conventional AdS gravity duals (in favorable cases), a chiral CFT can never have a conventional Einstein gravity dual. However, as explained in [33, 34], there are candidates for chiral gravity duals to holomorphic CFTs. See also [35] and references therein for a more detailed discussion on these theories. In this sense, we can consider the partition functions which follow as (candidate) duals to (a suitably defined theory of) chiral gravity (coupled to suitable matter).

Using the formula for the second-quantized partition function [26], along with the famous denominator identity due to Borcherds [36]:

$$\prod_{n>0, m \in \mathbb{Z}} (1 - p^n q^m)^{c(nm)} = p(J(\sigma) - J(\tau)) \tag{4.22}$$

where $p = e^{2\pi i \sigma}$ and $q = e^{2\pi i \tau}$ and $J(\tau) = q^{-1} + \sum_{n=1}^{\infty} c(n)q^n$, one can write the generating function:

$$\sum_{N=0}^{\infty} e^{2\pi i N \sigma} Z(\text{Sym}^N(\mathcal{M}); \tau) = \frac{e^{-2\pi i \sigma}}{J(\sigma) - J(\tau)}. \tag{4.23}$$

For large $\text{Im}(\tau)$ the infinite sum only converges for $\text{Im}(\sigma) > \text{Im}(\tau)$, while for small $\text{Im}(\tau)$ the infinite sum only converges for $\text{Im}(\sigma + \frac{1}{\tau}) > 1$. Choosing large $\text{Im}(\tau)$ we can say that

$$Z(\text{Sym}^N(\mathcal{M}); \tau) = \oint d\sigma \frac{e^{-2\pi i(N+1)\sigma}}{J(\sigma) - J(\tau)}, \tag{4.24}$$

where the contour is a circle at constant $\text{Im}(\sigma)$ on the cylinder given by the quotient of the σ -plane by $\sigma \sim \sigma + 1$ and we must assume $\text{Im}(\sigma) > \text{Im}(\tau)$. The contour integral can—at least naively—be evaluated by deforming the contour to smaller values of $\text{Im}(\sigma)$ approaching $\text{Im}(\sigma) = 0$. (We certainly cannot deform to large $\text{Im}(\sigma)$ because of the exponential growth from the term $e^{-2\pi i(N+1)\sigma}$.) This deformation leads to residues from an infinite set of simple poles at $\sigma = \tau$ together with σ equal to all the modular images of τ within the strip $|\text{Re}(\tau)| \leq \frac{1}{2}$. Using

$$\frac{1}{2\pi i} \frac{\partial}{\partial \tau} J(\tau) = -\frac{E_4^2(\tau)E_6(\tau)}{\eta(\tau)^{24}}. \tag{4.25}$$

This naive contour deformation yields:

$$Z(\text{Sym}^N(\mathcal{M}); \tau) = \mathbf{P}_2(q^{-N-1}) \frac{\eta(\tau)^{24}}{E_4(\tau)^2 E_6(\tau)}. \tag{4.26}$$

Here, $\mathbf{P}_2(q^{-N-1})$ is the weight 2 Poincaré series of q^{-N-1} .³

³ This Poincaré series requires regularization, indicating the above contour deformation argument is subtle. A standard procedure for obtaining a well-defined Poincaré series is described in detail in many places. See, for examples, Sect. 4 of [14] or Sect. 2 of [46]. As explained in those references, the modular anomaly of the series $\mathbf{P}_2(q^{-N-1})$ is expressed in terms of a

Because

$$\mathbf{P}_2(q^{-N-1}) = q^{-N-1} + \mathcal{O}(1) , \tag{4.27}$$

all of the modes which provide the low-energy spectrum (i.e., the states which are not black holes) are visible in the expansion of

$$F(\tau) = \frac{\eta(\tau)^{24}}{E_4(\tau)^2 E_6(\tau)} . \tag{4.28}$$

It now follows from the fact that $c/24 = N$ and the structure of \mathbf{P}_2 that we can find the modes at energies below the black hole bound just from expanding F . Writing

$$F(\tau) = \sum_{n=1}^{\infty} a_n q^n , \tag{4.29}$$

a_1 is the ground-state contribution and the higher a_k count the excited states visible in the partition function (until one reaches the threshold to form black holes).

One can extract the k th coefficient via the contour integral

$$a_k = \frac{1}{2\pi i} \oint d\tau \frac{1}{q^{k+1}} F(\tau) . \tag{4.30}$$

As $\eta(\tau)$ has no poles, E_4 has a simple zero at $\tau = e^{\frac{2\pi i}{3}}$ with no other zeroes, and E_6 has a simple zero at $\tau = i$ with no other zeros, we can now evaluate (4.30) explicitly.

The pole at $\tau = i$ provides the dominant behavior of the integral for $k \gg 1$. One finds

$$a_k \sim e^{2\pi k} \frac{\eta(i)^{24}}{E_4(i)^2 E_6'(i)} , \tag{4.31}$$

and hence in the regime $1 \ll n \ll N = \frac{c}{24}$, the $\text{Sym}^N(\mathcal{M})$ theory has a degeneracy of polar states governed by

$$a_n \sim e^{2\pi n} . \tag{4.32}$$

One can view this as satisfying an analog of the bound (3.32) for chiral gravity. In harmony with this, the singularity of (4.23) at $\sigma = \tau$ and at $\sigma = -1/\tau$ should come from the $N \rightarrow \infty$ limit of the partition functions, and this strongly suggests that the partition functions $Z(\text{Sym}^N(\mathcal{M}); \tau)$ exhibit the expected Hawking–Page first-order transition (as indeed follows from the general results of [24]), that is, the large N asymptotics at fixed pure imaginary τ is given by:

$$Z(\text{Sym}^N(\mathcal{M}); \tau) \sim \begin{cases} \kappa_1 N^{\kappa_2} q^{-N} (1 + \mathcal{O}(N^{-1})) & \text{Im}(\tau) \geq 1 \\ \kappa_1 N^{\kappa_2} \tilde{q}^{-N} (1 + \mathcal{O}(N^{-1})) & \text{Im}(\tau) \leq 1 \end{cases} \tag{4.33}$$

where $\tilde{q} := \exp(-2\pi i/\tau)$. Here κ_1, κ_2 are constants we have not attempted to determine.

Footnote 3 continued

period of a weight zero cusp form. Since no such nonzero cusp form exists we conclude that $\mathbf{P}_2(q^{-N-1})$ is in fact modular, as is required by modularity of $Z(\text{Sym}^N(\mathcal{M}); \tau)$.

The growth (4.32) exhibits a Hagedorn spectrum, hinting that if there is a holographically dual theory it must be a string theory with string scale comparable to the AdS radius.

5. String Versus Supergravity Duals

We have just seen that some theories with a low-energy Hagedorn degeneracy

$$\# \text{ of states at energy } E \sim e^{2\pi n}, \quad 1 \ll E \ll \frac{c}{24} \quad (5.1)$$

still satisfy our bounds. This might indicate that such theories are low-energy string theories—there is no parametric separation of scales evident between the emergence of a Hagedorn degeneracy and some other set of low-energy modes with well-defined asymptotics (which could serve as a proxy for supergravity KK modes).⁴

This is to be contrasted with the growth of states exhibited by a supergravity theory in d spatial dimensions, in the regime where the supergravity modes have wavelengths shorter than any scale set by the curvature. The gravity modes then behave, to leading approximation, like a gas of free particles in d dimensions. The energy per unit volume scales as

$$\mathcal{E} \sim T^{d+1}, \quad (5.2)$$

while the entropy per unit volume scales as

$$s \sim T^d. \quad (5.3)$$

Hence, in such a theory, one expects (simply from dimensional analysis) that

$$c_E \sim e^{\text{const} \times E^\alpha}, \quad \alpha \equiv \frac{d}{d+1} \quad (5.4)$$

in the regime dominated by supergravity modes. For instance, in the canonical $\text{AdS}_5 \times S^5$ solution of IIB supergravity, there is a supergravity regime with $E^{\frac{9}{10}}$ growth of the entropy as a function of energy [15].

For $\text{AdS}_3 \times S^3 \times K3$ compactifications where the $K3$ is much smaller than the S^3 , one would expect a 6d supergravity regime to occur at low energies. We now provide some simple analytical and numerical arguments demonstrating that the growth is indeed sub-Hagedorn. Related discussions appear in [37, 38]. The naive “gas of particles” analogy discussed above, for polar terms, would suggest a growth of $e^{\text{const} \times E^{5/6}}$. One can get slower growth, however, due to cancellations in the supergravity modes which contribute to the elliptic genus. We also note that at $g_{\text{string}} \ll 1$, there would be a regime of energies in the full physical theory exhibiting a Hagedorn degeneracy of string states. These do not, however, contribute in the elliptic genus.

⁴ Two subtleties could invalidate the considerations of this section. In one direction, cancellations between terms in a partition function could lead to subexponential growth of coefficients when in fact the entropy grows exponentially. In the other direction, when considering the entropy at finite volume it can happen that the entropy grows exponentially with energy, even though the theory is not a string theory. For an example, see Sect. 7 of [47].

First, we provide an analytical argument demonstrating that there is a range in which the polar terms of the elliptic genus of $\text{Sym}^N(K3)$ clearly has subexponential growth (though we do not quantify beyond this). Taking (4.4) at $y = 1$, we get that the sum of all $O(q^0)$ coefficients of the EG of $\text{Sym}^N(K3)$ is the N th coefficient of

$$\frac{q}{\eta(\tau)^{24}} \tag{5.5}$$

which goes as

$$e^{4\pi\sqrt{N}+O(\log N)}. \tag{5.6}$$

Since all of the $O(q^0)$ pieces of the EG of $\text{Sym}^N(K3)$ are positive (which can be shown from (4.4) for instance), each individual term must be smaller than (5.6). If we label the $O(q^0)$ states by E as above (we are interested in the growth in the NS sector, and the different powers of y at $O(q^0)$ in the R sector genus give states of different NS energy), we must have

$$a_E < e^{4\pi\sqrt{N}} \tag{5.7}$$

Thus

$$a_{N^\alpha} < e^{4\pi\sqrt{N}} \tag{5.8}$$

for $\alpha < 1$ which correspond to states parametrically below the Planck mass in the NS sector as $N \rightarrow \infty$. Relabeling gives us

$$a_E < e^{4\pi E^{\frac{1}{2\alpha}}}. \tag{5.9}$$

We therefore find states parametrically lighter than the Planck mass with a subexponential growth of states. Note that there may be other states at the same energy level that we neglect due to only considering $O(q^0)$ terms in the elliptic genus. However, as we expect the entropy to be a function of polarity up to small corrections, taking terms with positive powers of q into account would only multiply our expression in (5.9) by some polynomial factor without changing the leading order.

Because we expect the only relevant scales (other than supergravity KK scales) to be the string scale and Planck scale, and we do not get stringy growth in this regime, we expect subexponential growth throughout the polar terms. We now provide further (weak) numerical evidence in favor of this hypothesis. We include a plot of the normalized coefficients of y^{N-x} for $x = 1, \dots, 40$ in the large N limit in Fig. 8 (these numbers do not change past some N since they only involve twisted sectors of permutations of some fixed length).

These examples suggest a criterion that distinguishes between theories with low-energy Einstein gravity duals as opposed to low-energy string duals, with the usual qualifier that cancellation is possible in an index computation. Writing

$$c_E \sim e^{\text{const} \times E^\alpha}, \quad 1 \ll E \ll \frac{c}{24}, \tag{5.10}$$

theories with $\alpha < 1$ are likely to have a range of scales at low energy where supergravity applies, while theories with $\alpha = 1$ are evidently string theories already at the scale set by the curvature. We note that similar issues have been




FIGURE 8. Here, we plot the normalized coefficients of y^{N-x} terms in elliptic genus of $\text{Sym}^N(K3)$ for $x = 1, \dots, 40$ in the large N limit. Note the subexponential growth in the plot. Numerical values for the first twenty terms are given in Table 1

discussed, in the context of the duality between AdS_4 gravity and CFT_3 , in the interesting paper [39].

6. Estimating the Volume of an Interesting Set of Modular Forms

In this section, we use (3.32) to try and quantify a lower bound on the “fraction of large m superconformal field theories which may admit a gravity dual”. Our approach will be to ask: “How special is the class of weight zero, index m Jacobi forms corresponding to such superconformal theories?” As we have seen, thermodynamic arguments constrain the growth of the polar coefficients provided there is a physically reasonable gravitational dual, so the problem reduces to quantifying “what fraction” of all possible polar coefficients corresponds to the theories with gravitational duals.

Since the Jacobi form is completely determined by its polar coefficients, the map from CFTs to elliptic genera can be viewed as a map from the space of $(0, 2)$ field theories to a subset $\mathcal{E} \subset \mathbb{Z}^{j(m)}$. Now, there is a natural metric on the moduli spaces of conformal field theories, namely, the Zamolodchikov metric [40]. The moduli space of such theories, with a fixed central charge c , is a union of connected components $\Pi_\alpha \mathcal{M}_\alpha^{(c)}$. It was suggested some time ago that, at least for the space of $(2, 2)$ superconformal theories, the total Zamolodchikov volume of $V^{(c)} := \sum_\alpha \text{vol}(\mathcal{M}_\alpha^{(c)})$ should be finite. This was based on physical arguments [41, 42]. For the case of components arising from Calabi–Yau manifolds it has been shown that indeed $\text{vol}(\mathcal{M}_\alpha^{(c)})$ is finite. (See [43] and references therein for the mathematical work on this subject.) The

finiteness of $V^{(c)}$ would allow us to define a measure on the space of $(2, 2)$ theories of a fixed central charge and thereby to quantify statements of “how often” a property is exhibited in a natural way. We will assume that $V^{(c)}$ is in fact finite.⁵

Using the push-forward measure under the map to the polar coefficients of elliptic genera we obtain a natural measure on the space \mathcal{E} of polar coefficients. Unfortunately, our present state of knowledge of conformal field theory is too primitive to evaluate this measure in great detail, but to illustrate the idea, and some of the issues which will arise, we will sketch two toy computations.

For our first toy computation we consider the pushforward to a measure on \mathbb{Z}_+ for the absolute value of the extreme polar coefficient of the elliptic genus. We denote this by

$$\epsilon(\mathcal{C}) = |c(0, m; \mathcal{C})| \tag{6.1}$$

for a $(2, 2)$ CFT \mathcal{C} with $c = 6m$.

Now ϵ is multiplicative on CFTs,

$$\epsilon(\mathcal{C}_1 \times \mathcal{C}_2) = \epsilon(\mathcal{C}_1)\epsilon(\mathcal{C}_2). \tag{6.2}$$

We would also like to say the same for the volumes:

$$\text{vol}(\mathcal{C}_1 \times \mathcal{C}_2) \stackrel{?}{=} \text{vol}(\mathcal{C}_1)\text{vol}(\mathcal{C}_2) \tag{6.3}$$

but this is in general not the case. Here $\text{vol}(\mathcal{C})$ denotes the volume of the connected component of $V^{(c)}$ in which \mathcal{C} lies. A simple counterexample is provided by conformal field theories with toroidal target spaces. Nevertheless, for ensembles such as theories based on *generic* Calabi–Yau manifolds the volume is multiplicative, because the relevant Hodge numbers are additive. We will refer to an ensemble of CFT’s for which (6.3) holds as a *multiplicative ensemble* and here we restrict attention to such ensembles. Extending our discussion beyond multiplicative ensembles is an interesting, but potentially difficult, problem.

Given a multiplicative ensemble, let us say an $N = (2, 2)$ CFT \mathcal{C} is *prime* if it is not the product of two such theories \mathcal{C}_1 and \mathcal{C}_2 each with positive central charge. Let $\mathcal{C}(m, \alpha)$ denote the distinct prime CFT’s of central charge $c = 6m$, with $\alpha = 1, \dots, f_m$. We expect f_m to be finite, but this is not necessary for our construction, so long as the relevant products below converge. Denote the absolute value of the extreme polar coefficient, and the Zamolodchikov volume of $\mathcal{C}(m, \alpha)$ by $\epsilon(m, \alpha), v(m, \alpha)$, respectively. Then the Zamolodchikov volume $\text{vol}(M)$ of theories of central charge $c = 6M$ is determined from:

$$\prod_{m=1}^{\infty} \prod_{\alpha=1}^{f_m} \frac{1}{1 - v(m, \alpha)q^m} = 1 + \sum_{M=1}^{\infty} \text{vol}(M)q^M. \tag{6.4}$$

Similarly, we can write a generating function for the volume of the theories with a fixed extreme polar coefficient. We assume that $\epsilon(m, \alpha) \neq 0$ in our

⁵ Friedan has proposed a mechanism by which such a probability distribution might in fact be dynamically generated from more fundamental principles [48, 49].

ensemble (thus excluding, for example, Calabi–Yau models with odd complex dimension) and form the generating function:

$$\prod_{m=1}^{\infty} \prod_{\alpha=1}^{f_m} \frac{1}{1 - v(m, \alpha) \epsilon(m, \alpha)^{-s} q^m} = 1 + \sum_{M=1}^{\infty} \xi(s; M) q^M \tag{6.5}$$

Then

$$\xi(s; M) = \sum_{\mathbf{e}=1}^{\infty} \frac{\text{vol}(\mathbf{e}; M)}{\mathbf{e}^s} \tag{6.6}$$

and the measure for the extreme polar coefficient is

$$\mu(\mathbf{e}; M) := \frac{\text{vol}(\mathbf{e}; M)}{\text{vol}(M)}. \tag{6.7}$$

In order to make this slightly more concrete, let us restrict even further to the ensemble of (4, 4) theories generated by taking products of the symmetric products of $K3$ sigma models, such as

$$(\text{Sym}^1(K3))^{n_1} \times (\text{Sym}^2(K3))^{n_2} \times \dots \times (\text{Sym}^{\ell}(K3))^{n_{\ell}}. \tag{6.8}$$

We will call this the $K3$ -ensemble and it is a multiplicative ensemble of CFT’s. In this ensemble the prime CFTs are simply the symmetric products $\text{Sym}^n(K3)$. For $\text{Sym}^1(K3)$ the moduli space \mathcal{M}_1 is the famous double quotient

$$\mathcal{M}_1 = O(\Gamma) \backslash O(4, 20; \mathbb{R}) / O(4) \times O(20) \tag{6.9}$$

with $\Gamma \cong \Pi^4 \oplus E_8 \oplus E_8$, while for $N > 1$ the moduli space is [25, 44, 45]

$$\mathcal{M}_N = O(\Gamma') \backslash O(4, 21; \mathbb{R}) / O(4) \times O(21) \tag{6.10}$$

with Γ' a lattice of signature 4, 21 determined in [45]. The four “extra moduli” in (6.10) compared to (6.9) are due to the blowup multiplet at the locus of A_1 singularities in $\text{Sym}^N(K3)$ where two points meet. All higher twist fields are irrelevant. Denote the Zamolodchikov volume of these moduli spaces by v_N . The Zamolodchikov volume $\text{vol}(M)$ of the ensemble of models (6.8) is then simply given by

$$\prod_{n=1}^{\infty} \frac{1}{1 - v_n q^n} = 1 + \sum_{M=1}^{\infty} \text{vol}(M) q^M \tag{6.11}$$

Now, to get the measure for a fixed extreme polar term we noted above that

$$\epsilon(\text{Sym}^n(K3)) = n + 1, \tag{6.12}$$

so the extreme polar term of the elliptic genus of (6.8) is just the product:

$$2^{n_1} 3^{n_2} \dots (\ell + 1)^{n_{\ell}}. \tag{6.13}$$

Therefore, our general formula specializes to

$$\prod_{m=1}^{\infty} \frac{1}{1 - v_m(m + 1)^{-s} q^m} = 1 + \sum_{M=1}^{\infty} \xi(s; M) q^M, \tag{6.14}$$

where $\xi(s; M)$ defines the conditional volume as in (6.6) and the measure for the extreme polar term is given by (6.7), above.

Determining the numerical values of the constants v_N used above is a very interesting problem in number theory. This will be discussed in a separate paper, along with some applications of the function $\xi(s; M)$ to the central issue of this paper.⁶ It would also be very interesting to extend the above discussion to the ensemble of all $(4, 4)$ theories, but this looks quite challenging. We would need to include products with $\text{Sym}^N(T^4)/T^4$. Moreover, we have omitted products with other $(4, 4)$ models constructable from permutation orbifolds, or from other compact hyperkahler manifolds arising from moduli spaces of hyperholomorphic bundles on $K3$ and T^4 . And we have omitted the unknown unknowns since we do not know that every $(4, 4)$ model can be realized geometrically. Nevertheless, we expect some of the basic features of the above discussion to survive better knowledge of the moduli space.

The above discussion is our first toy computation. Given our poor knowledge of the moduli space of conformal field theories we will resort to a second toy computation. We hope it proves instructive. We enumerate the polar coefficients $c(a)$ by decreasing discriminant $D(a) = \ell(a)^2 - 4mn(a)$, $a = 0, \dots, j(m) - 1$ where $j(m) = \dim \tilde{J}_{0,m}$. Thus, $D(0) = m^2$. The idea of the second toy computation is to find a natural probability measure on the vector space of polar coefficients $(c(0), \dots, c(N))$. Of course, a vector space has infinite measure in its Euclidean norm so we map these coefficients to an affine coordinate patch of $\mathbb{R}P^N$, with $N = j(m)$. That is, we consider the points $[1 : c(0) : \dots : c(N)]$ in $\mathbb{R}P^N$. We then consider the Fubini-Study measure on this patch. Whether this measure bears any relation to the a priori Zamolodchikov measure (in the large N limit) remains to be seen. (Since we do not like the answer, we suspect the answer is that it does not.)

The volume element for the unit radius $\mathbb{R}P^N$ in affine coordinates $[1 : \xi^1 : \dots : \xi^N]$ is:

$$d\text{vol} = \frac{d\xi^1 \wedge \dots \wedge d\xi^N}{(1 + \sum_a (\xi^a)^2)^{(N+1)/2}}. \tag{6.15}$$

Now we consider the subspace of the affine coordinate patch with

$$|c(a)| \leq R(a). \tag{6.16}$$

$R(a)$ is a bound which is supposed to come from physics. One reasonable guess is

$$R(a) = e^{2\pi(n(a) - \frac{|\ell(a)|}{2} + \frac{m}{2})}. \tag{6.17}$$

Note that this is imposing (3.32) without allowing an $O(\log m)$ correction. Concretely, we are interested in the fraction

$$f_N = \frac{\Gamma(\frac{N+1}{2})}{\pi^{\frac{N+1}{2}}} \left(\prod_{i=1}^N \int_{-e^{2\pi(n_i - \frac{|\ell_i|}{2} + \frac{m}{2})}}^{e^{2\pi(n_i - \frac{|\ell_i|}{2} + \frac{m}{2})}} d\xi_i \right) \frac{1}{(1 + \sum_i (\xi^i)^2)^{\frac{N+1}{2}}} \tag{6.18}$$

in the limit $N \rightarrow \infty$.

⁶ For further details, see <https://www.perimeterinstitute.ca/video-library/collection/mock-modularity-moonshine-and-string-theory>.

In Appendix B, we show that in the limit of large N ,

$$0.9699 < f_N < 0.9725. \quad (6.19)$$

We actually view this as a good indication that the Fubini-Study measure is *not* a good surrogate for the Zamolodchikov measure. On general grounds, one actually expects theories with weakly coupled gravity duals (even characterizing some small region of their moduli space) to be rare creatures.

In general CFTs, the number of excited states at large energies n grows like $e^{2\pi\sqrt{\frac{c}{6}n}}$ by the Cardy formula. Hence a measure which was based on “expecting” there to be a small number of states in that regime would clearly be incorrect. While one cannot use Cardy’s result in the energy range characterizing polar coefficients, it seems suspicious that our measure “expects” the fewer polar coefficients—related to states with high energy, though below the black hole bound—to be close to 0. In fact, one might expect that in a random SCFT, the polar coefficients typically grow fairly rapidly with decreasing polarity. In such a case, it would be more difficult for them to lie within the polydisc specified by our bounds. Finding a modified volume estimate (or attaching a plausible physical meaning to our present estimate) will have to remain a problem for the future.

Acknowledgements

We thank J. de Boer, A. Castro, E. Dyer, A.L. Fitzpatrick, D. Friedan, S. Harrison, K. Jensen, C. Keller, S. Minwalla, D. Tong, S.P. Trivedi, E. Verlinde, R. Volpato, and X. Yin for interesting discussions. We thank D. Ramirez for help with sophisticated mathematical diagrams and C. Keller for useful comments on the draft. This work was initiated at the Aspen Center for Physics and we thank the ACP for its hospitality and excellent working conditions. The ACP is supported by NSF Grant No. PHY-1066293. N.B. acknowledges the support of the Stanford Graduate Fellowship. N.M.P. acknowledges support from an NSF Graduate Research Fellowship. S.K. is grateful to the 2014 Indian Strings Conference, IISER-Pune, the Tata Institute for Fundamental Research, and the Kavli Institute for Theoretical Physics at UC Santa Barbara for hospitality while thinking about the physics of this note. His work was supported in part by National Science Foundation Grant PHY-0756174 and DOE Office of Basic Energy Sciences contract DE-AC02-76SF00515. M.C. is grateful to Cambridge, Case Western Reserve, and Stanford Universities and Max Planck Institut für Mathematik for hospitality. The work of G.M. is supported by the DOE under Grant DOE-SC0010008 to Rutgers, and NSF Focused Research Group award DMS-1160461.

Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the

source, provide a link to the Creative Commons license, and indicate if changes were made.

Appendix A: Extended Phase Diagram

Here, we derive in detail the extended phase diagram depicted in Fig. 1. The logic of the argument can be summarized as follows. The expression of the elliptic genus as a regularized Poincaré sum involves a sum over all co-prime pairs of integers (c, d) . For each such pair arising from the invariant group Γ_θ of $Z_{NS,+}$, we will find that there is an (n, ℓ) labeling a polar term in the elliptic genus which can serve as the analogue of our ground state in the ground-state dominance condition. As a consequence, in each region in the tessellated upper-half plane there is a single pair (c, d) labeling the saddle point which dominates the gravitational path integral (CFT elliptic genus). Each phase transition across the bold lines in Fig. 1 is then a modular copy of the one we studied in this paper.

The elliptic genus can be written in terms of its polar part as [14]

$$\begin{aligned}
 Z_{R,R}(\tau, z) &= \frac{1}{2} \sum_{r \in \mathbb{Z}/2m\mathbb{Z}} C_r(0) \theta_{m,r}(\tau, z) + \frac{1}{2} \sum_{\substack{\ell \in \mathbb{Z}, n \geq 0 \\ D(n,\ell) > 0}} \lim_{K \rightarrow \infty} \sum_{(\Gamma_\infty \backslash \Gamma)_K} \\
 C_\ell(D(n, \ell)) \exp \left(2\pi i \left(n \frac{a\tau + b}{c\tau + d} + \ell \frac{z}{c\tau + d} - m \frac{cz^2}{c\tau + d} \right) \right) &R \left(\frac{2\pi i D(n, \ell)}{4m c(c\tau + d)} \right)
 \end{aligned}
 \tag{A.1}$$

where the limit coset is given by

$$\lim_{K \rightarrow \infty} \sum_{(\Gamma_\infty \backslash \Gamma)_K} := \lim_{K \rightarrow \infty} \sum_{\substack{0 < c < K \\ -K^2 < d < K^2 \\ (c,d)=1}}
 \tag{A.2}$$

and R is the regularization factor

$$R(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-z} z^{1/2} dz = \operatorname{erf}(\sqrt{x}) - 2\sqrt{\frac{x}{\pi}} e^{-x},
 \tag{A.3}$$

where $\operatorname{erf}(z) = \frac{2}{\sqrt{\pi}} \int_0^z e^{-t^2} dt$ denotes the error function.

As discussed in [13], §6, using the classic identities

$$\begin{aligned}
 \frac{a\tau + b}{c\tau + d} &= \frac{a}{c} - \frac{1}{c(c\tau + d)} \\
 \frac{1}{2} \frac{\tau + 1}{c\tau + d} &= \frac{1}{2c} - \frac{d}{2} \frac{1}{c(c\tau + d)} + \frac{1}{2} \frac{1}{c\tau + d} \\
 \frac{c(\tau/2 + 1/2)^2}{c\tau + d} &= \frac{\tau}{4} + \frac{2c - d}{4c} + \frac{1}{4} \frac{c^2 - 2cd + d^2}{c(c\tau + d)}
 \end{aligned}$$

and $\text{Im}\left(-\frac{1}{c(c\tau+d)}\right) = \frac{\text{Im}(\tau)}{|c\tau+d|^2} = \text{Im}\left(\frac{a\tau+b}{c\tau+d}\right)$, we see that

$$Z_{\text{NS},+}(\tau) = (-1)^m q^{\frac{m}{4}} \frac{1}{2} \sum_{r \in \mathbb{Z}/2m\mathbb{Z}} C_r(0) \theta_{m,r} \left(\tau, \frac{\tau+1}{2} \right) + \frac{1}{2} \sum_{\substack{\ell \in \mathbb{Z}, n \geq 0 \\ D(n,\ell) > 0}} \lim_{K \rightarrow \infty} \sum_{(\Gamma_\infty \setminus \Gamma)_K} X(n, \ell; c, d) R \left(\frac{2\pi i D(n, \ell)}{4mc(c\tau + d)} \right) \quad (\text{A.4})$$

with

$$\left| X(n, \ell; c, d) \right| = |C_\ell(D(n, \ell))| \exp \left(-2\pi \text{Im} \left(\frac{a\tau + b}{c\tau + d} \right) \left(m \frac{(d-c)^2}{4} + n + \ell \frac{d-c}{2} \right) \right). \quad (\text{A.5})$$

We would like to know which term in the elliptic genus, i.e., which pair (n, ℓ) , contributes the most to the sum in (A.1) with a given pair (c, d) . First, focusing on the exponential factor in (A.5), using that $\text{Im}\left(\frac{a\tau+b}{c\tau+d}\right) > 0$ and $0 < D(n, \ell) \leq m^2$ we conclude that the maximum of $\exp \left(-2\pi \text{Im}\left(\frac{a\tau+b}{c\tau+d}\right) \left(m \frac{(d-c)^2}{4} + n + \ell \frac{d-c}{2} \right) \right)$ occurs at $(n, \ell) = (n_{c,d}, \ell_{c,d})$,

$$(n_{c,d}, \ell_{c,d}) := \left(\frac{m}{4}((d-c)^2 - 1), -m(d-c) \right)$$

when $d - c$ is odd. Ignoring the other factors for the moment, we expect that $\left| X(n, \ell; c, d) \right|$ has its maximum

$$\left| X(n_{c,d}, \ell_{c,d}; c, d) \right| = C_{-m}(m^2) \exp \left(2\pi \frac{m}{4} \text{Im}\left(\frac{a\tau + b}{c\tau + d}\right) \right) \quad (\text{A.6})$$

when $(n, \ell) = (n_{c,d}, \ell_{c,d})$. In the above we have used the fact that $c(n_{c,d}, \ell_{c,d}) = c(0, m) = C_{-m}(m^2)$ is equal to the number of NS ground states [see (2.11)].

The situation is different for the pair of co-primes integers (c, d) with even $d - c$. Using the more refined condition for the discriminants of the polar terms

$$0 < D(n, \ell) \leq r^2 \text{ where } -m < r \leq m, \ell = r \pmod{2m}$$

that holds for all weak Jacobi forms as a straightforward consequence of (2.13), we see that the maximum of the exponential term in (A.5) is of order 1 which is achieved whenever $\ell = -(d-c)m + r, n = m(d-c)^2 - \frac{(d-c)r}{2}$ for any $-m < r \leq m$. In other words, the contribution of the part of the sum given by a pair (c, d) with $c - d \equiv 0 \pmod{2}$ in (A.1) is exponentially suppressed.

As a result, assuming that the exponential factor in (A.6) is the dominating factor and ignoring for the moment the regularization factor, one concludes that in each region in the upper-half plane given by the tessellation by $\Gamma_\infty \setminus \Gamma_\theta$ there is a unique pair (c, d) that dominates and this corresponds to the infinitely many phases of 3d quantum gravity. To see this, notice that

$$\text{Im} \left(\frac{a\tau + b}{c\tau + d} \right) = \frac{\text{Im}(\tau)}{|c\tau + d|^2} \leq \text{Im}(\tau) \quad \forall \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_\theta$$

whenever $\tau \in \Gamma_\infty \mathcal{F}$ is in the (interior of the) fundamental domain

$$\mathcal{F} = \{\tau \in \mathbb{H} \mid |\tau| > 1, -1 < \operatorname{Re}(\tau) < 1\}$$

of Γ_θ or any of its images under the translation $\tau \rightarrow \tau + 2n, n \in \mathbb{Z}$. See Fig. 1.

Next we would like to discuss the conditions under which that the term with $(n, \ell) = (n_{c,d}, \ell_{c,d})$ indeed dominates the sum over all polar terms for a given pair (c, d) . First we show that the effect of the regularization factor can be ignored at the large central charge limit where $D(n_{c,d}, \ell_{c,d})/4m = m/4 \gg 1$. To see this, note that $R(x) \rightarrow 0$ as $x \rightarrow 0$ and

$$R(x) - 1 = O(\sqrt{x} e^{-x})$$

as $x \rightarrow \infty$, and

$$\operatorname{Re} \left(\frac{2\pi i D(n, \ell)}{4mc(c\tau + d)} \right) = \frac{2\pi D(n, \ell)}{4m} \operatorname{Im} \left(\frac{a\tau + b}{c\tau + d} \right).$$

Second, for there to be no term over dominating the term coming from $(n, \ell) = (n_{c,d}, \ell_{c,d})$ in the sum in the region where $\frac{a\tau + b}{c\tau + d} \in \Gamma_\infty \mathcal{F}$ as predicted by analyzing the exponential factor alone as in (A.5), focussing on the line $\operatorname{Re}(\frac{a\tau + b}{c\tau + d}) = 0$ we see that the coefficients of the polar terms have to satisfy

$$\log |c(n_{c,d}, \ell_{c,d})| \leq \left(2\pi \left(\frac{m((d-c)^2 + 1)}{4} + n + \frac{\ell(d-c)}{2} \right) \right) + O(\log m) \quad (\text{A.7})$$

for all co-prime pairs (c, d) with $d - c$ odd. It is not hard to show that the seemingly stronger condition (A.7) is in fact implied by our bound (3.37) when taking the spectral flow symmetry into account. Recalling that $c(n_{c,d}, \ell_{c,d}) = C_{-m}(m^2)$ and

$$c(n, \ell) = c(n(k), \ell(k)), \quad n(k) = n + k^2 m + k\ell, \quad \ell(k) = \ell + 2km$$

for all $k \in \mathbb{Z}$, we can write (A.7) as

$$\log |c(n(k), \ell(k))| \leq 2\pi \left(n(k) - \frac{|\ell(k)|}{2} + \frac{m}{2} \right) + O(\log m)$$

where $k = \frac{d-c-1}{2}$.

In summary, we have proved the following. The condition (A.7) is required for $Z_{NS,+}(\tau)$ to be consistent with the phase structure given by the group $\Gamma_\infty \setminus \Gamma_\theta$ (corresponding to distinct Euclidean BTZ black holes which dominate in different regions of parameter space [23], §7.3). We have seen that the necessary condition (3.37) that we derived earlier in the paper, governing the Hawking–Page transition, is sufficient to guarantee (A.7), and hence the full expected phase diagram.

Appendix B: Estimating the Volumes of Regions in \mathbb{RP}^N

Recall the problem we have. We would like to estimate

$$f_N = \frac{\Gamma(\frac{N+1}{2})}{\pi^{\frac{N+1}{2}}} \left(\prod_{i=1}^N \int_{-e^{2\pi(n_i - \frac{|\ell_i|}{2} + \frac{m}{2})}}^{e^{2\pi(n_i - \frac{|\ell_i|}{2} + \frac{m}{2})}} d\xi_i \right) \frac{1}{(1 + \sum_i (\xi_i)^2)^{\frac{N+1}{2}}} \quad (\text{B.1})$$

TABLE 2. Most polar terms at index m (excluding y^m)

Term	$e^{2\pi(n - \frac{ \ell }{2} + \frac{m}{2})}$
y^{m-1}	e^π
y^{m-2}	$e^{2\pi}$
qy^m	$e^{2\pi}$
y^{m-3}	$e^{3\pi}$
qy^{m-1}	$e^{3\pi}$
y^{m-4}	$e^{4\pi}$
qy^{m-2}	$e^{4\pi}$
q^2y^m	$e^{4\pi}$
y^{m-5}	$e^{5\pi}$
qy^{m-3}	$e^{5\pi}$
q^2y^{m-1}	$e^{5\pi}$

in the large N limit where $N + 1$ is the number of polar terms of a Jacobi form of index m .

As an example, let us consider $m = 2$. We will later switch to the large m limit. There are only two polar terms: y^2 and y^1 so $N = 1$. We normalize the y^2 coefficient to 1, and the coefficient for y^1 parametrizes $\mathbb{R}P^1$ (the “point at infinity” corresponds to a y^2 coefficient of 0, but this has measure zero).

For y^1 , $\ell = 1$ and $n = 0$, so

$$e^{2\pi(n - \frac{|\ell|}{2} + \frac{m}{2})} = e^\pi. \tag{B.2}$$

Thus the integral is

$$f_2 = \frac{\Gamma(1)}{\pi} \int_{-e^\pi}^{e^\pi} d\xi_1 \frac{1}{(1 + \xi_1^2)^1} = 0.9725. \tag{B.3}$$

In the large m limit, there are $\lceil \frac{k+1}{2} \rceil$ integrals with limits $-e^{k\pi}$ to $e^{k\pi}$ (see Table 2).

Appendix B.1: An Upper Bound

In this section, we will derive an upper bound on f_N of 0.9725. Recall the famous fact of life that

$$\begin{aligned} & \frac{\Gamma(\frac{N+2}{2})}{\pi^{\frac{N+2}{2}}} \int_{-\infty}^{\infty} d\xi_{N+1} \frac{1}{(1 + \xi_1^2 + \dots + \xi_{N+1}^2)^{\frac{N+2}{2}}} \\ &= \frac{\Gamma(\frac{N+1}{2})}{\pi^{\frac{N+1}{2}}} \frac{1}{(1 + \xi_1^2 + \dots + \xi_N^2)^{\frac{N+1}{2}}}. \end{aligned} \tag{B.4}$$

Thus, we can always take extra integrals to ∞ and we will get a strictly bigger value. No matter how big N is, we will always have an integral with limits

$-e^\pi$ to e^π (coming from the y^{m-1} term). In particular

$$\begin{aligned}
 f_N &= \frac{\Gamma(\frac{N+1}{2})}{\pi^{\frac{N+1}{2}}} \int_{-e^\pi}^{e^\pi} d\xi_1 \int d\xi_2 \dots \int d\xi_N \frac{1}{(1 + \xi_1^2 + \dots + \xi_N^2)^{\frac{N+1}{2}}} \\
 &< \frac{\Gamma(\frac{N+1}{2})}{\pi^{\frac{N+1}{2}}} \int_{-e^\pi}^{e^\pi} d\xi_1 \int_{-\infty}^{\infty} d\xi_2 \dots \int_{-\infty}^{\infty} d\xi_N \frac{1}{(1 + \xi_1^2 + \dots + \xi_N^2)^{\frac{N+1}{2}}} \\
 &= \frac{1}{\pi} \int_{-e^\pi}^{e^\pi} d\xi_1 \frac{1}{1 + \xi_1^2} \\
 &= 0.9725,
 \end{aligned} \tag{B.5}$$

where we define an integral $d\xi_i$ with unlabeled limits as from $-e^{2\pi(n_i - \frac{|\ell_i|}{2} + \frac{m}{2})}$ to $e^{2\pi(n_i - \frac{|\ell_i|}{2} + \frac{m}{2})}$.

Appendix B.2: A Lower Bound

Now we will show a lower bound of 0.9699 through a series of inequalities. Again we use the same convention of unlabeled limits of integration $d\xi_i$ being from $-e^{2\pi(n_i - \frac{\ell_i^2}{4m} + \frac{m}{4})}$ to $e^{2\pi(n_i - \frac{\ell_i^2}{4m} + \frac{m}{4})}$.

First we will show

$$\begin{aligned}
 f_N &= \frac{\Gamma(\frac{N+1}{2})}{\pi^{\frac{N+1}{2}}} \int d\xi_1 \int d\xi_2 \dots \int d\xi_N \frac{1}{(1 + \xi_1^2 + \dots + \xi_N^2)^{\frac{N+1}{2}}} \\
 &> 1 - \sum_{i=1}^N \left(1 - \frac{1}{\pi} \int d\xi_i \frac{1}{1 + \xi_i^2} \right).
 \end{aligned} \tag{B.6}$$

To see this, first rewrite (B.6) as

$$\begin{aligned}
 &1 - \frac{\Gamma(\frac{N+1}{2})}{\pi^{\frac{N+1}{2}}} \int d\xi_1 \int d\xi_2 \dots \int d\xi_N \frac{1}{(1 + \xi_1^2 + \dots + \xi_N^2)^{\frac{N+1}{2}}} \\
 &< \sum_{i=1}^N \left(1 - \frac{1}{\pi} \int d\xi_i \frac{1}{1 + \xi_i^2} \right).
 \end{aligned} \tag{B.7}$$

Now note that

$$\frac{1}{\pi} \int d\xi_i \frac{1}{1 + \xi_i^2} \tag{B.8}$$

represents the fraction of \mathbb{RP}^N where ξ_i is between the appropriate limits of $-e^{2\pi(n_i - \frac{|\ell_i|}{2} + \frac{m}{2})}$ to $e^{2\pi(n_i - \frac{|\ell_i|}{2} + \frac{m}{2})}$. We write this as a fraction of \mathbb{RP}^N instead of \mathbb{RP}^1 by using (B.4) to add the remaining $N - 1$ integrals from $-\infty$ to ∞ and change the prefactor.

In more detail, let us take the first term ($i = 1$) in the sum in the right-hand side of (B.7). That term is

$$\begin{aligned}
 &1 - \frac{1}{\pi} \int_{-e^\pi}^{e^\pi} d\xi_1 \frac{1}{1 + \xi_1^2} \\
 &= 1 - \frac{\Gamma(\frac{N+1}{2})}{\pi^{\frac{N+1}{2}}} \int_{-e^\pi}^{e^\pi} d\xi_1 \int_{-\infty}^{\infty} d\xi_2 \dots \int_{-\infty}^{\infty} d\xi_N \frac{1}{(1 + \xi_1^2 + \dots + \xi_N^2)^{\frac{N+1}{2}}}
 \end{aligned} \tag{B.9}$$

which is exactly the region *outside* $-e^\pi < \xi_1 < e^\pi$ in \mathbb{RP}^N . However, the left-hand side of (B.7) is \mathbb{RP}^N with the region

$$|\xi_i| \leq R(i), \quad \forall i$$

excluded.

Thus, (B.7) is satisfied by using the fact that the complement of the intersection is less than the sum of complements.

Now we are in business. It is another classic fact of life that

$$\begin{aligned} \frac{1}{\pi} \int_{-R(i)}^{R(i)} d\xi_i \frac{1}{1 + \xi_i^2} &= \frac{2}{\pi} \arctan R(i) \\ &= 1 - \frac{2}{\pi R(i)} + \dots \end{aligned} \quad (\text{B.10})$$

Plug into (B.6), to get:

$$1 - \sum_{i=1}^N \left(1 - \frac{1}{\pi} \int d\xi_i \frac{1}{1 + \xi_i^2} \right) > 1 - \sum_{i=1}^N \frac{2}{\pi R(i)} \quad (\text{B.11})$$

In the large m limit, the first terms look like

$$\begin{aligned} 1 - \sum_{i=1}^N \frac{2}{\pi a_i} &= 1 - \frac{2}{\pi} \left(\frac{1}{e^\pi} + \frac{2}{e^{2\pi}} + \frac{2}{e^{3\pi}} + \frac{3}{e^{4\pi}} + \frac{3}{e^{5\pi}} + \dots \right) \\ &> 1 - \frac{2}{\pi} \left(\frac{1}{e^\pi} + \frac{2}{e^{2\pi}} + \frac{3}{e^{3\pi}} + \frac{4}{e^{4\pi}} + \frac{5}{e^{5\pi}} + \dots \right) \\ &= 1 - \frac{2}{\pi} \left(\frac{1}{e^\pi (1 - \frac{1}{e^\pi})^2} \right) \\ &= 0.9699. \end{aligned} \quad (\text{B.12})$$

Thus, putting everything together, we get

$$f_N > 0.9699. \quad (\text{B.13})$$

References

- [1] Maldacena, J.M.: The large N limit of superconformal field theories and supergravity. *Adv. Theor. Math. Phys.* **2**, 231 (1998). [arXiv:hep-th/9711200](#)
- [2] Hartman, T., Keller, C. A., Stoica, B.: Universal spectrum of 2d conformal field theory in the large c limit. [arXiv:1405.5137](#) [hep-th]
- [3] Witten, E.: Anti-de Sitter space, thermal phase transition, and confinement in gauge theories. *Adv. Theor. Math. Phys.* **2**, 505 (1998). [arXiv:hep-th/9803131](#)
- [4] Hawking, S.W., Page, D.N.: Thermodynamics of black holes in anti-De Sitter space. *Commun. Math. Phys.* **87**, 577 (1983)
- [5] Schellekens, A.N., Warner, N.P.: Anomalies and modular invariance in string theory. *Phys. Lett. B* **177**, 317 (1986)
- [6] Schellekens, A.N., Warner, N.P.: Anomaly cancellation and selfdual lattices. *Phys. Lett. B* **181**, 339 (1986)

- [7] Pilch, K., Schellekens, A.N., Warner, N.P.: Path integral calculation of string anomalies. *Nucl. Phys. B* **287**, 362 (1987)
- [8] Witten, E.: Elliptic genera and quantum field theory. *Commun. Math. Phys.* **109**, 525 (1987)
- [9] Eichler, M., Zagier, D.: *The theory of Jacobi forms*. Birkhäuser, Boston (1985)
- [10] Strominger, A., Vafa, C.: Microscopic origin of the Bekenstein-Hawking entropy. *Phys. Lett. B* **379**, 99 (1996). [arXiv:hep-th/9601029](#)
- [11] Kawai, T., Yamada, Y., Yang, S. K.: Elliptic genera and $N = 2$ superconformal field theory. *Nucl. Phys. B* **414**, 191 (1994). [arXiv:hep-th/9306096](#)
- [12] Gaberdiel, M.R., Gukov, S., Keller, C.A., Moore, G.W., Ooguri, H.: Extremal $N = (2,2)$ 2D conformal field theories and constraints of modularity. *Commun. Number Theor. Phys.* **2**, 743 (2008), [arXiv:0805.4216](#) [hep-th]
- [13] Dijkgraaf, R., Maldacena, J.M., Moore, G.W., Verlinde, E.P.: A black hole farey tail. [arXiv:hep-th/0005003](#)
- [14] Manschot, J., Moore, G.W.: A modern farey tail. *Commun. Number Theor. Phys.* **4**, 103 (2010), [arXiv:0712.0573](#) [hep-th]
- [15] Aharony, O., Gubser, S. S., Maldacena, J.M., Ooguri, H., Oz, Y.: Large N field theories, string theory and gravity. *Phys. Rep.* **323**, 183 (2000). [arXiv:hep-th/9905111](#)
- [16] Kraus, P., Larsen, F.: Partition functions and elliptic genera from supergravity. *JHEP* **0701**, 002 (2007). [arXiv:hep-th/0607138](#)
- [17] Kraus, P.: Lectures on black holes and the AdS(3)/CFT(2) correspondence. *Lect. Notes Phys.* **755**, 193 (2008). [arXiv:hep-th/0609074](#)
- [18] Banados, M., Teitelboim, C., Zanelli, J.: The black hole in three-dimensional space-time. *Phys. Rev. Lett.* **69**, 1849 (1992). [arXiv:hep-th/9204099](#)
- [19] Cvetic, M., Larsen, F.: Near horizon geometry of rotating black holes in five-dimensions. *Nucl. Phys. B* **531**, 239 (1998). [arXiv:hep-th/9805097](#)
- [20] de Boer, J., Denef, F., El-Showk, S., Messamah, I., Van den Bleeken, D.: Black hole bound states in AdS(3) \times S**2. *JHEP* **0811**, 050 (2008). [arXiv:0802.2257](#) [hep-th]
- [21] Bena, I., Chowdhury, B.D., de Boer, J., El-Showk, S., Shigemori, M.: moulting black holes. *JHEP* **1203**, 094 (2012). [arXiv:1108.0411](#) [hep-th]
- [22] Maldacena, J.M., Strominger, A.: AdS(3) black holes and a stringy exclusion principle. *JHEP* **9812**, 005 (1998). [arXiv:hep-th/9804085](#)
- [23] Maloney, A., Witten, E.: Quantum gravity partition functions in three dimensions. *JHEP* **1002**, 029 (2010). [arXiv:0712.0155](#) [hep-th]
- [24] Keller, C.A.: Phase transitions in symmetric orbifold CFTs and universality. *JHEP* **1103**, 114 (2011). [[arXiv:1101.4937](#) [hep-th]]
- [25] Dijkgraaf, R.: Instanton strings and hyperKähler geometry. *Nucl. Phys. B* **543**, 545 (1999). [arXiv:hep-th/9810210](#)
- [26] Dijkgraaf, R., Moore, G.W., Verlinde, E.P., Verlinde, H.L.: Elliptic genera of symmetric products and second quantized strings. *Commun. Math. Phys.* **185**, 197 (1997). [arXiv:hep-th/9608096](#)
- [27] Manschot, J., Zapata Rolon, J.M.: The asymptotic profile of χ_y -genera of Hilbert schemes of points on K3 surfaces. *Commun. Number Theor. Phys.* **09**, 413 (2015). [arXiv:1411.1093](#) [math.AG]

- [28] Maldacena, J.M., G.W., Strominger, A.: Counting BPS black holes in toroidal type II string theory. [arXiv:hep-th/9903163](#)
- [29] Haehl, F.M., Rangamani, M.: Permutation orbifolds and holography. [arXiv:1412.2759](#) [hep-th]
- [30] Belin, A., Keller, C.A., Maloney, A.: String universality for permutation orbifolds. [arXiv:1412.7159](#) [hep-th]
- [31] Gritsenko, V.: Elliptic genus of Calabi-Yau manifolds and Jacobi and Siegel modular forms. [arXiv:math/9906190](#)
- [32] Frenkel, I., Lepowsky, J., Meurman, A.: Vertex operator algebras and the monster, Pure and Applied Mathematics, vol. 134. Academic (1988), Boston, p. 508
- [33] Li, W., Song, W., Strominger, A.: Chiral gravity in three dimensions. JHEP **0804**, 082 (2008). [arXiv:0801.4566](#) [hep-th]
- [34] Maloney, A., Song, W., Strominger, A.: Chiral gravity, log gravity and extremal CFT. Phys. Rev. D **81**, 064007 (2010). [arXiv:0903.4573](#) [hep-th]
- [35] Skenderis, K., Taylor, M., Van Rees, B.C.: Topologically massive gravity and the AdS/CFT correspondence. JHEP **0909**, 045 (2009). [arXiv:0906.4926](#) [hep-th]
- [36] Borchers, R.E.: The monster Lie algebra. Adv. Math. **83**, 30 (1990)
- [37] de Boer, J.: Large N elliptic genus and AdS/CFT correspondence. JHEP **9905**, 017 (1999). [arXiv:hep-th/9812240](#)
- [38] de Boer, J., El-Showk, S., Messamah, I., Van den Bleeken, D.: A bound on the entropy of supergravity? JHEP **1002**, 062 (2010). [arXiv:0906.0011](#) [hep-th]
- [39] Minwalla, S., Narayan, P., Sharma, T., Umesh, V., Yin, X.: Supersymmetric states in large N Chern-Simons-Matter theories. JHEP **1202**, 022 (2012). [arXiv:1104.0680](#) [hep-th]
- [40] Zamolodchikov, A.B.: Irreversibility of the flux of the renormalization group in a 2D field theory. JETP Lett. **43**, 730 (1986)
- [41] Horne, J.H., Moore, G.W.: Chaotic coupling constants. Nucl. Phys. B **432**, 109 (1994). [arXiv:hep-th/9403058](#)
- [42] Douglas, M.R., Lu, Z.: Finiteness of volume of moduli spaces. [arXiv:hep-th/0509224](#)
- [43] Douglas, M., Lu, Z.: On the geometry of moduli space of polarized Calabi-Yau manifolds. [arXiv:math/0603414](#) [math-dg]
- [44] Giveon, A., Kutasov, D., Seiberg, N.: Comments on string theory on AdS(3). Adv. Theor. Math. Phys. **2**, 733 (1998). [arXiv:hep-th/9806194](#)
- [45] Seiberg, N., Witten, E.: The D1/D5 system and singular CFT. JHEP **9904**, 017 (1999). [arXiv:hep-th/9903224](#)
- [46] Cheng, M.C.N., Duncan, J.F.R.: Rademacher sums and rademacher series. [arXiv:1210.3066](#) [math.NT]
- [47] Galakhov, D., Longhi, P., Mainiero, T., Moore, G.W., Neitzke, A.: Wild wall crossing and BPS giants. JHEP **1311**, 046 (2013). [arXiv:1305.5454](#) [hep-th]
- [48] Friedan, D.: A Tentative theory of large distance physics. JHEP **0310**, 063 (2003). [arXiv:hep-th/0204131](#)
- [49] Friedan, D.: Two talks on a tentative theory of large distance physics. [arXiv:hep-th/0212268](#)

Nathan Benjamin, Shamit Kachru and Natalie M. Paquette

Stanford Institute for Theoretical Physics

Stanford University

Stanford, CA 94305, USA

e-mail: nathansb@stanford.edu; skachru@stanford.edu; npaquett@stanford.edu

and

SLAC National Accelerator Laboratory

2575 Sand Hill Road

Menlo Park, CA 94025, USA

Miranda C. N. Cheng

Institute of Physics and Korteweg-de Vries Institute for Mathematics

University of Amsterdam

Amsterdam, The Netherlands

e-mail: mcheng@uva.nl

Gregory W. Moore

NHETC and Department of Physics and Astronomy

Rutgers University

Piscataway, NJ 08855, USA

e-mail: gmoore@physics.rutgers.edu

Communicated by Boris Pioline.

Received: July 8, 2015.

Accepted: December 7, 2015.