



UvA-DARE (Digital Academic Repository)

Periodontitis in twins : smoking, microbiological and immunological aspects

Torres de Heens, G.L.

Publication date
2010

[Link to publication](#)

Citation for published version (APA):

Torres de Heens, G. L. (2010). *Periodontitis in twins : smoking, microbiological and immunological aspects*. [Thesis, fully internal, Universiteit van Amsterdam].

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Chapter 2

Effects of smoking on the *ex vivo* cytokine production in periodontitis

G. L. Torres de Heens¹, R. Kikkert², L. A. Aarden², U. van der Velden¹
and B. G. Loos¹

¹Department of Periodontology, Academic Center for Dentistry Amsterdam, ACTA, The Netherlands and ²Department of Immunopathology, Sanquin Research and Landsteiner Laboratory, Academic Medical Center, University of Amsterdam, The Netherlands

Journal of Periodontal Research 2009; **44**, 28-34

Abstract

Background and Objective: Smoking is associated with increased severity of periodontitis. The underlying mechanisms of this phenomenon are not well understood. The purpose of the present study was to compare the monocyte-derived T cell directing (Th1/Th2) response and pro-inflammatory cytokine production in *ex vivo* whole blood cell cultures (WBCC) of smoking and non-smoking chronic periodontitis patients.

Material and Methods: Venous blood was collected from 29 periodontitis patients (18 non-smokers and 11 smokers) receiving supportive periodontal treatment, and diluted 10-fold for WBCC. WBCC were stimulated for 18 hours with *Neisseria meningitidis* lipo-oligosaccharide (LOS) and *Porphyromonas gingivalis* sonic extract (Pg-SE). The production of the T cell directing cytokines interleukin (IL)-12 p40 and IL-10, as well as the pro-inflammatory cytokines IL-1 β , IL-6 and IL-8, was measured in the culture supernatants.

Results: After LOS stimulation of WBCC, smokers showed a lower IL-12 p40/IL-10 ratio than non-smokers ($p < 0.05$). IL-1 β production was significantly lower in smokers as compared to non-smokers after stimulation with both LOS and Pg-SE ($p < 0.05$). IL-6 and IL-8 production was similar between both groups, for both LOS and Pg-SE.

Conclusion: A more pronounced Th2 response in smoking periodontitis patients may be related to increased severity of the disease.

Introduction

Periodontitis is a chronic, multifactorial, infectious disease of the supporting tissue of the teeth. Periodontitis patients suffer from gradual loss of tooth attachment in the jaw bone leading to periodontal pockets, receding gums, loose teeth and eventually tooth exfoliation (Kinane & Lappin 2001). The onset and progression of periodontitis is due to an imbalance of the interaction between bacterial pathogens and host immunity. Host immunity is greatly influenced by both genetic susceptibility and environmental risk factors (Page et al. 1997). Although several specific periodontopathogens have been implicated in the disease, two Gram-negative bacteria, *Porphyromonas gingivalis* and *Actinobacillus actinomycetemcomitans* have the strongest association (van Winkelhoff et al. 2002). Cigarette smoking is considered as one of the most important environmental risk factors in periodontitis (Martinez-Canut et al. 1995) as more clinical attachment loss and bone loss have been observed in smoking than in non-smoking patients (Grossi et al. 1995, Grossi et al. 1994). Moreover, smoking may be responsible for a less favorable outcome after periodontal treatment and may more frequently cause disease progression despite a strict periodontal maintenance care program (Darby et al. 2005).

Monocytes play a crucial role in orchestrating the host immune system. When triggered by whole bacteria as well as bacterial components, monocytes produce cytokines (monokines) which direct both innate and adaptive immunity (Seymour & Gemmell 2001b). Monokines such as the pro-inflammatory interleukin (IL)-1 β , IL-6, IL-8, IL-12 and anti-inflammatory IL-10 have been shown to be part of the inflammatory response in periodontitis and most likely may determine the host susceptibility and thereby variation in periodontal destruction (Gemmell et al. 1997, Gemmell et al. 2002, Niho et al. 1998, Seymour & Gemmell 2001a).

Cytokines derived mainly from dendritic cells, monocytes and macrophages, play a pivotal role in directing lymphocytic differentiation of non-committed precursors CD4⁺ T cells into either T-helper (Th)1 or Th2 cells (Kidd 2003). In human infections, an imbalance of Th1 and Th2 cytokine production may be related to disease progression (Kidd 2003). The role of T-helper cells in amplifying immune responsiveness is well established. In periodontitis, the nature of the lymphocytic infiltrate seems to be crucial in disease progression (Sigusch et al. 1998). A Th1 cytokine profile is the major mediator in the early/stable lesion, while the dominance

of B-cells/plasma cells in the advanced/progressive lesion would suggest a role for Th2 cells. Therefore, it is likely that in those later stages, changes in cytokine profiles that modulate the Th1/Th2 balance may affect the susceptibility to or the course of the periodontal infection (Gemmell et al. 2002). Previous studies have shown that periodontitis patients display a monocytic-cytokine profile which may favor a Th2 immune response. Indeed, periodontitis is currently regarded as a Th2-type disease; therefore, the Th2-monocytic promoting phenotype may be an important risk factor (Fokkema et al. 2002, Gemmell & Seymour 1994a).

Differences in monocytic cytokine production between smoking and non-smoking periodontitis patients have been scarcely investigated. However, clinical data demonstrate that, compared to non-smokers, patients who smoke show a more severe disease and relapse during supportive periodontal treatment (SPT). In other inflammatory diseases it has been shown that tobacco smoke may exacerbate disease progression through “priming” of immune cells toward a Th2 phenotype (Byron et al. 1994, Cozen et al. 2004). In several inflammatory conditions, including periodontitis, *ex vivo* stimulation assays of peripheral blood cells with LPS stimulant have been used as a measure of the host immune capacity (Fokkema et al. 2003, Malave et al. 1989, Swaak et al. 1997, van der Pouw Kraan et al. 1997). Monocyte responsiveness, both in isolated monocyte cell cultures and whole blood cell cultures (WBCC), is regarded as a reliable measure for the *in vivo* situation (van den Heuvel et al. 1998).

The aim of the present study was to compare the *ex vivo* production of IL-1 β , IL-6, IL-8, IL-10, and IL-12 p40 in whole blood cell cultures of smoking and non-smoking periodontitis patients positive for *Porphyromonas gingivalis* (*P. gingivalis*), after stimulation with lipooligosaccharide from *Neisseria meningitidis* (LOS) and a sonic extract of *P. gingivalis* (SE-Pg).

Material and Methods

Patients

The study population consisted of patients referred to our clinic (Department of Periodontology at the Academic Center for Dentistry Amsterdam [ACTA]) for the treatment of periodontal disease. Patients were selected from a pool of 900 consecutive patients that were sampled for bacteriological investigation at the intake before periodontal treatment. The selection criteria included: 1) Western European

Caucasian descent, 2) diagnosis of chronic adult periodontitis at intake, 3) receiving SPT after non-surgical and surgical treatment had been completed, 4) age between 40 and 60 years, 5) presence of ≥ 20 permanent teeth, 6) periodontal bone loss of $\geq 1/3$ of the total root length at ≥ 2 teeth as visible on peri-apical radiographs, and 7) subgingival presence of *P. gingivalis*. Exclusion criteria were: 1) presence of any systemic condition that may affect the periodontal status, 2) pregnancy, 3) use of antibiotics within the last 6 months preceding the study, and 4) use of any medicine that may interfere with the periodontal health. This selection resulted in 48 potentially eligible patients. Of the 48 patients, 29 volunteered to participate in the present study. Subsequently, patients were classified into two groups according to the reported smoking status: a) non-smokers: those who had never smoked or had ceased smoking more than 10 years before entering the study, and b) smokers: current smokers who had been smoking for at least 10 years with a consumption of ≥ 10 cigarettes/day (c/d).

The final study population included 18 non-smokers and 11 smokers. For each subject, all teeth were radiographically examined for interproximal bone loss at the mesial and distal aspects, using cemento-enamel junction (CEJ) of the tooth and the bone crest as reference points. With the use of a translucent plastic ruler (Schei ruler technique), the percentage of bone loss at the deepest proximal site of each tooth was measured.

Approval for this study was obtained by the Medical Ethical Committee of the Academic Medical Center of the University of Amsterdam. Participants were informed both verbally and in writing about the purpose of the study, and provided signed informed consent.

Stimuli

Lipo-oligosaccharide (LOS) as used previously (van der Pouw Kraan et al. 1995) was purified from *Neisseria meningitidis*, strain H44/76 (a kind gift from Dr. J. Poolman, RIVM, Bilthoven, The Netherlands). *P. gingivalis* strain 381 was grown in brain heart infusion broth enriched with hemin (5 mg/l) and menadione (1 mg/l) in an anaerobic atmosphere (80% N₂, 10% H₂, 10% CO₂) for 48 h at 37 °C during 48 h at 37 °C. The *P. gingivalis* bacteria were harvested in the log phase, pelleted by centrifugation (8000 g), washed three times in phosphate-buffered saline (PBS), and resuspended in PBS at a concentration of optical density (OD)₆₉₀ = 1, corresponding to approximately 7×10^8 colony forming units per ml. Aliquots (500 μ l) of resuspended

bacteria were disrupted using a sonifier in a sonifying vessel on ice (Soniprep MSE 150, amplitude 18, 4 min, 5 s intervals).

The degree of disruption of the bacteria was assessed by phase-contrast microscopy and with Gram-staining by light microscopy. Sonicates were stored at 4 °C until use. Before use, *P. gingivalis* sonicates were centrifuged (8000 g, 1 min) and used in WBCC as described below.

A mouse monoclonal antibody raised against human CD3 (anti-CD3, CLB-T3/4.E) was from CLB, Amsterdam, The Netherlands and has been previously described (van Lier et al. 1987b).

Whole Blood Cell Cultures

Preliminary experiments were performed to determine the most optimal LOS and Pg-SE concentration for the WBCC. Whole blood samples of periodontally healthy donors were used for this purpose. From each subject, venous blood was collected by venipuncture from the antecubital fossae in a sterile pyrogen-free blood collection tube (Vacuette, Greiner, Alphen a/d Rijn, The Netherlands) containing sodium heparine and diluted 10-fold in pyrogen-free Iscove's modified Dulbecco's medium (IMDM, Bio Whittaker, Verviers, Belgium) supplemented with 0.1% fetal calf serum (FCS, Bodinco, Alkmaar, The Netherlands), 100 IU/ml penicillin, 100 µg/ml streptomycin (Gibco, Merelbeke, Belgium), and 15 IU/ml sodium-heparin (Leo Pharmaceutical Products B.V., Weesp, The Netherlands). Diluted whole blood in 200 µl flat-bottom microtitre culture plates (Nunc, Roskilde, Denmark) was stimulated during 18 hours with different LOS concentrations (1000, 250, 62, 16 pg/ml) (van der Pouw Kraan et al. 1995) or Pg-SE dilutions (1:100, 1:400, 1:1600, 1:64000), and cytokine production of IL-1 β, IL-6, IL-8, IL-10 and IL-12 p40 was measured. The most optimal LOS and Pg-SE concentrations for WBCC stimulation were 1000 pg/ml and 1:100 respectively (data not shown). Especially for IL-10, the capacity of both stimulants alone to elicit cytokine at the indicated concentrations was still weak. To increase IL-10 production we decided to combine monocyte stimulation with a T cell stimulus. In the 18 hr incubation period anti-CD3 alone did not lead to measurable cytokine production. However, when combined with monocyte stimuli such as LOS or Pg-SE, anti-CD3 led to strongly increased production of cytokines such as IL-10 and IL-12 (data not shown).

From each subject of the final study population (29 subjects) diluted whole blood as described previously was stimulated with LOS at a final concentration of 1000 pg/ml or with Pg-SE 1:100 dilution, in the presence of anti-CD3 at 1 µg/ml. Unstimulated diluted whole blood served as a negative control. Supernatants were harvested and stored at -20 °C until cytokine measurements.

Venous blood of each participant was also collected in an EDTA (K₃)-containing tube (Becton Dickinson Vacutainer System Europe, Meylan, France) for the determination of the total number of leukocytes, and leukocyte differentiation (neutrophils, eosinophils, basophils, lymphocytes and monocytes), with standard automated procedures (Cell-Dyn 4000, Hematology Analyzer, Abbott Laboratories, Park, Illinois, USA.) operated in the clinical chemistry laboratory of the Slotervaart Hospital, Amsterdam, The Netherlands.

Toll-like receptor-transfected human embryonic kidney 293 (HEK 293) cell cultures

Human Embryonic Kidney 293 (HEK 293) cells stably transfected with CD14, CD14-Toll-like receptor (TLR)2 or CD14-TLR4 were a kind gift from Drs. D. Golenbock and E. Latz, Worcester, University of Massachusetts Medical School, Division of Infectious Diseases, MA, USA. Transfected HEK cells were cultured in IMDM supplemented with 5% heat-inactivated FCS, 100 U/ml penicillin, 100 µl/ml streptomycin, 50 µM 2-mercaptoethanol (Sigma-Aldrich, Steinheim, Germany), and 5 µg/ml puromycin (Sigma-Aldrich). For stimulation experiments, cells were seeded at 5×10^5 cells per well in 96-well flat-bottomed microtitre plates (Nunc, Roskilde, Denmark), and stimulated the next day with the appropriate preparation. HEK 293-CD14-TLR4 cells were stimulated in the presence of 5% human serum as described elsewhere (25). After 16-20 h incubation, supernatants were harvested for determination of IL-8 production as a marker for nuclear factor kappa B (NF-κB) activation, i.e. cell activation.

Assays for Cytokines

Cytokine levels of IL-1β, IL-6, IL-8, IL-10, and IL-12 p40 were measured in the supernatants of WBCC using commercially available enzyme-linked immunosorbent assay (ELISA) kits (PeliKine Compact™ human ELISA kits, CLB,

Amsterdam, The Netherlands) as previously described (van der Pouw Kraan et al. 1997). The plates were read in an ELISA-reader (Labsystems Multiskan Multisoft, Helsinki, Finland) at 450 nm, with 540 nm as a reference. Cytokine production in WBCC supernatants was adjusted per 10^6 monocytes.

Statistical Analysis

The SPSS package version 11.0 for Windows (Chicago, IL, USA) was used for descriptive data, data analysis and box plot generation. Differences between smokers and non-smokers were analyzed by Mann Whitney *U*-test. Differences in number of subjects per group were tested by Fisher exact tests.

Results

Subject background characteristics age, gender and educational level are presented in Table 1. Cigarette smoking consumption reported by current smokers showed an average of 14 c/d during 35 years with a range of 10-20 cigarettes/day. The levels of total white blood cells were significantly higher in smokers than non-smokers (7.42 and $5.78 \times 10^9/l$ respectively, $p= 0.001$). This was mainly due to increased numbers of neutrophils in smokers compared to their non-smoking counterparts. Values for eosinophils, basophils, lymphocytes and monocytes did not differ between the two groups (Table 1). Dental radiographic analysis of subjects at the moment of intake is presented in Table 2. Smokers had, on average, 16.1 teeth with $\geq 30\%$ bone loss and 5.1 teeth with $\geq 50\%$, while non-smoking patients had 12.1 and 3.7 respectively (differences not significant).

Table 1. Characteristics of the study population (non-smoking and smoking periodontitis patients). Values are means \pm standard deviation or number (%) of subjects.

	Non-smokers (N = 18)	Smokers (N = 11)	<i>p</i> -value
Age (years)	54 \pm 5.3	51 \pm 7.2	0.310
Gender			
Female	11 (61%)	7 (64%)	1.000
Male	7 (39%)	4 (36%)	
Education level			
\leq High school	6 (33%)	2 (18%)	0.671
$>$ High school	12 (67%)	9 (82%)	
Smoking habits			
Number of cigarettes/day	0	14.2 \pm 4.4	
Number of years smoking	0	35 \pm 6.9	
Total WBC ($10^9/l$)	5.78 \pm 1.36	7.42 \pm 1.11	0.001
Neutrophils ($10^9/l$)	3.22 \pm 1.11	4.31 \pm 0.81	0.005
Eosinophils ($10^9/l$)	0.14 \pm 0.12	0.19 \pm 0.12	0.280
Basophils ($10^9/l$)	0.02 \pm 0.07	0 \pm 0	0.261
Lymphocytes ($10^9/l$)	1.90 \pm 0.46	2.21 \pm 0.77	0.079
Monocytes ($10^9/l$)	0.46 \pm 0.13	0.50 \pm 0.13	0.485

Table 2. Dental radiographic characteristics of the study population at intake (non-smoking and smoking periodontitis patients). Values are means \pm standard deviation.

	Non-smokers (N = 18)	Smokers (N = 11)	<i>p</i> -value
Number of teeth	26.4 \pm 2.7	26.2 \pm 3.2	0.964
Teeth \geq 30% bone loss	12.1 \pm 7.5	16.1 \pm 5.6	0.120
Teeth \geq 50% bone loss	3.7 \pm 3.3	5.1 \pm 4.3	0.420

After stimulation of WBCC with LOS, supernatants were analyzed for the cytokines IL-12 p40 and IL-10 (Figure 1, panels A and B). Regarding IL-12 p40 no

differences could be assessed between smokers and non-smokers. The median of IL-10 for non-smokers was 14300 ng/10⁶ monocytes, whereas in smokers the median was 21300 ng/10⁶ monocytes. However the values were not significantly different ($p=0.217$). To explore the balance of Th1/Th2, we calculated the IL-12 p40/IL-10 ratio (Fig. 1, panel C). Results showed a significant lower ratio in smokers when compared to non-smokers ($p=0.022$), which may indicate a more pronounced Th2 response in this group. Production of IL-1 β , IL-6 and IL-8 after LOS stimulation is presented in Figure 1, panels D, E and F, respectively. The IL-1 β production was significantly higher in non-smokers than smokers ($p=0.012$), whereas no differences were found for IL-6 and IL-8 cytokine production between the two groups.

In addition to the LOS stimulation experiments, Pg-SE was similarly used to explore the cytokine production of WBCC when challenged with a periodontal pathogen. Results are displayed next to the LOS box plots in Figure 1. No difference in IL-12 p40 production was found between smokers and non-smokers (panel A). The levels of IL-10 and the calculated IL-12 p40/IL10 ratio showed no difference between smokers and non-smokers (Fig. 1, panels B and C). Pro-inflammatory cytokine IL-1 β , as it occurred after LOS stimulation, was significantly higher in non-smokers than smokers ($p=0.015$, Fig. 1, panel D), while IL-6 and IL-8 did not differ between the groups (Fig. 1, panels E and F).

Smokers and non-smokers produced higher amounts of IL-8 after stimulation with Pg-SE when compared to LOS ($p<0.001$, Fig. 1, panel F) showing a clear difference in the levels of cytokine production dependent on the stimulation used. It is well known that Toll-like receptors (TLR) have the ability of transmitting LPS signaling across the cell membrane. It has been previously shown that the activation of either TLR2 or TLR4 may influence the T_H1/T_H2 balance. Therefore, in an effort to explain the observed differences between LOS and Pg-SE, it was decided to test the specificity of LOS and Pg-SE used in our experiments for both TLR2 and TLR4 on transfected HEK 293 cells. After stimulation of the cells with either LOS or Pg-SE, IL-8 production was measured in harvested supernatants. We observed that LOS stimulated HEK-CD14-TLR4 but not HEK-CD14-TLR2 to produce IL-8. Inversely, when Pg-SE was used, only HEK-CD14-TLR2 cells were able to produce IL-8 (one graph representative of three experiments is shown, Figure 2). So in line with

expectations we observed that LOS signaled through TLR4, whereas Pg-SE stimulated TLR2.




Figure 1. Boxplots for T cell directing monocytic cytokines IL-12 p40, IL-10, IL-12 p40/ IL-10 ratio (panels A, B, C, respectively), and pro-inflammatory cytokines IL-1 β , IL-6, IL-8 (panels D, E, F) in ng/10⁶ monocytes of whole blood cell cultures from non-smoking () and smoking (■) periodontitis patients after 18 hrs stimulation with *Neisseria meningitidis* LOS (LOS) and *Porphyromonas gingivalis* sonicate extract (Pg-SE), in the presence of anti-CD3 1 μ g/ml. *: $p < 0.05$ and ***: $p < 0.001$




Figure 2. IL-8 cytokine production of TLR2 and TLR4 in transfected HEK cell cultures induced by LOS and Pg-SE. HEK 293 cell lines were stimulated with *Neisseria meningitidis* LOS (LOS) at concentrations 0.01, 0.1, 1 and 10 pg/ml (panel A) and *Porphyromonas gingivalis* sonicate extract (Pg-SE) at serial dilutions (panel B).

Discussion

A body of evidence indicates that both prevalence and severity of periodontal disease is increased in smokers compared to non-smokers, and that smokers in SPT show faster relapse and/or progression of disease (Bergstrom 2003, Bergstrom 2004, Zambon et al. 1996b). However, to date, no comparative study of stimulated cytokine profile in smoking and non-smoking periodontitis patients exists in the literature. The present study constitutes a first attempt to explore the effects of smoking on the monocytic cytokine profile in chronic adult periodontitis patients. In previous studies a common and highly purified LPS from *Escherichia coli* (*E. coli*) has been mostly used for WBCC stimulation purposes. However, the use of *E. coli* may be criticised. Firstly, in the *in vivo* situation and during infection, LPS is surely not the only bacterial component interacting with immune cells. Secondly, *E. coli* is not a periodontal pathogen. In order to overcome these problems to some extent, we selected periodontitis patients who before treatment were subgingival positive for *P. gingivalis*, a major periodontal pathogen, and used next to *N. meningitidis* LOS, a sonic extract of *P. gingivalis* for WBCC stimulation. The addition of anti-CD3 in the WBCC enhanced the monocytic cytokine production elicited by the single use of either LOS or Pg-SE. Via stimulation of the T cell, anti-CD3 leads to co-stimulation

of TLR-triggered monocytes, possibly by upregulation of CD40-ligand (Kennedy et al. 1996). It could be argued that a potential mechanism of co-stimulation would be multivalent immunoglobulin (Ig) exposure to low-affinity Fc(γ) receptors (Fc γ R) or complement activation. However, to avoid such complications we made use of an anti-CD3 monoclonal antibody of the IgE isotype which is incapable of interacting with Fc γ R and with complement (Van Lier et al. 1987a).

In our experiments the decision of measuring IL-12 p40 and not IL-12 p70 is supported by previous observations. Firstly, IL-12 p40 subunit can be produced in large excess over the heterodimer IL-12 p70, favouring IL-12 p40 detection in the supernatants as IL-12 p70 could be practically undetectable (Bucht et al. 1996, Venier et al. 2007). Secondly, the production of IL-12 p40 and IL-12 p70 by LPS-IFN- γ -stimulated macrophages is affected to the same degree by smokeless tobacco suggesting that both cytokines may be similarly influenced by tobacco (Petro et al. 2002). Moreover, others have used the measurement of either p40 mRNA or p40 itself as indicators of total IL-12 production in humans (Chougnnet et al. 1996, Fulton et al. 1996). The Th1-promoting activity of IL-12 p40 has been shown and this cytokine is considered as an indicator of Th1 differentiation (Giambartolomei et al. 2004, Nakahara et al. 2005, Prebeck et al. 2001, Zaitseva et al. 1996).

Cytokine production has been usually studied in cultures of peripheral blood mononuclear cells (PBMC). However the study of cytokine production in WBCC has some advantages over separated cultures. Firstly, the whole blood culture system reduces the likelihood of endotoxin contamination due to minimal handling of the cells. Secondly, the risk of cellular activation due to the isolation procedures is reduced. Thirdly, the WBCC system may represent more closely the natural environment with the presence of various immunomodulating and pro- and anti-inflammatory mediators in whole blood. Fourth, disturbances in the ratios of different cell types due to purification procedures are avoided (Muller et al. 1998, van der Pouw Kraan et al. 1997). Therefore, the integrity of the cellular interactions is maintained as best possible, although the whole blood is diluted 1/10. It has been extensively studied and shown in parallel cultures of whole blood and freshly isolated monocytes as well as in kinetics that WBCC stimulated with LPS specifically reflect the behaviour of the monocytes (Snijders et al. 1996, Snijders et al. 1998, van der Pouw Kraan et al. 1995).

The present data show that smokers suffering from periodontitis have a lower IL-12 p40/IL-10 ratio after LOS stimulation and lower IL-1 β production after LOS and Pg-SE stimulation than their non-smoking counterparts. Recently it was reported that tobacco smoke leads to increase in cAMP levels (Du et al. 2005). In T cells cAMP elevations lead to a strongly decreased IL-12 p40/IL-10 ratio (van der Pouw Kraan et al. 1995). The lower IL-1 β production is in agreement with *in vitro* experiments that demonstrated that tobacco smoke can inhibit cytokine production by peripheral blood monocytes, including production of IL-1 β (Pabst et al. 1995). Furthermore, it has been shown that IL-1 is involved in the up-regulation of interferon gamma (IFN- γ) production by Th1 cells and down-regulation of IL-4 production by Th2 cells (O'Garra & Murphy 1994, Sandborg et al. 1995, Schmitz et al. 1993). Since we found in smokers lower levels of IL-1 β production and a lower IL-12 p40/IL-10 ratio it may be suggested that in periodontitis smoking influences the Th1/Th2 balance into a more pronounced Th2 profile. This may result in continuous polyclonal B cell activation with less protective antibody production (Gemmell & Seymour 2004) which could explain partly the clinical finding of more severe periodontitis in smokers.

The current study also showed that more IL-8 was produced in response to Pg-SE compared to LOS stimulation, both in smoking and non-smoking periodontitis patients. The finding that Pg-SE stimulates via TLR2, whereas LOS is recognised by TLR4, might provide an explanation for the difference in IL-8 production. This is in line with a body of evidence demonstrating striking differences in cytokine gene transcription in TLR2 and TLR4 activation (Pulendran et al. 2001, Re & Strominger 2001). Alternatively, the difference in IL-8 levels might be caused by differences in the cell types involved in IL-8 production. Since TLR2, but not TLR4, is substantially expressed on the surface of human neutrophils, it is likely that Pg-SE activates both monocytes and neutrophils, whereas after LOS stimulation only monocytes are activated. (Kurt-Jones et al. 2002). It is clear that the role of TLRs in periodontal pathogen recognition and subsequent cytokine production is not yet completely understood, and that the use of different microbial stimuli may be an important consideration for the interpretation of past and future data.

In summary, the results showed differences between smoking and non-smoking periodontitis patients in *ex vivo* cell culture cytokine production. Our

findings suggest that the Th2 pattern in periodontitis may be accentuated by smoking. This hypothesis is consistent with the contention that a strong innate immune response is associated with a protective Th1 pattern, favoring inflammatory responses that contain the infection with periodontal pathogens. Therefore, a more pronounced Th2 phenotype may accelerate disease progression and account for the relapse during SPT as frequently observed in smoking periodontitis patients. Our findings may serve as another step in revealing generalized inflammatory patterns in smoking periodontitis patients.

References

- Bergstrom, J. (2003) Tobacco smoking and risk for periodontal disease. *Journal of Clinical Periodontology* **30**, 107-113.
- Bergstrom, J. (2004) Tobacco smoking and chronic destructive periodontal disease. *Odontology* **92**, 1-8.
- Bucht, A., Larsson, P., Weisbrot, L., Thorne, C., Pisa, P., Smedegard, G., Keystone, E. C. & Gronberg, A. (1996) Expression of interferon-gamma (IFN-gamma), IL-10, IL-12 and transforming growth factor-beta (TGF-beta) mRNA in synovial fluid cells from patients in the early and late phases of rheumatoid arthritis (RA). *Clinical & Experimental Immunology* **103**, 357-367.
- Byron, K. A., Varigos, G. A. & Wootton, A. M. (1994) IL-4 production is increased in cigarette smokers. *Clinical & Experimental Immunology* **95**, 333-336.
- Chougnet, C., Wynn, T. A., Clerici, M., Landay, A. L., Kessler, H. A., Rusnak, J., Melcher, G. P., Sher, A. & Shearer, G. M. (1996) Molecular analysis of decreased interleukin-12 production in persons infected with human immunodeficiency virus. *The Journal of Infectious Diseases* **174**, 46-53.
- Cozen, W., Diaz-Sanchez, D., James Gauderman, W., Zadnick, J., Cockburn, M. G., Gill, P. S., Masood, R., Hamilton, A. S., Jyrala, M. & Mack, T. M. (2004) Th1 and Th2 cytokines and IgE levels in identical twins with varying levels of cigarette consumption. *Journal of Clinical Immunology* **24**, 617-622.
- Darby, I. B., Hodge, P. J., Riggio, M. P. & Kinane, D. F. (2005) Clinical and microbiological effect of scaling and root planing in smoker and non-smoker chronic and aggressive periodontitis patients. *Journal of Clinical Periodontology* **32**, 200-206.

- Du, B., Altorki, N. K., Kopelovich, L., Subbaramaiah, K. & Dannenberg, A. J. (2005) Tobacco smoke stimulates the transcription of amphiregulin in human oral epithelial cells: evidence of a cyclic AMP-responsive element binding protein-dependent mechanism. *Cancer Research* **65**, 5982-5988.
- Fokkema, S. J., Loos, B. G., de Slegte, C., Burger, W., Piscoer, M., Y, I. J. & Van der Velden, U. (2003) Increased release of IL-12p70 by monocytes after periodontal therapy. *Journal of Clinical Periodontology* **30**, 1091-1096.
- Fokkema, S. J., Loos, B. G., Slegte, C. & van der Velden, U. (2002) A type 2 response in lipopolysaccharide (LPS)-stimulated whole blood cell cultures from periodontitis patients. *Clinical & Experimental Immunology* **127**, 374-378.
- Fulton, S. A., Johnsen, J. M., Wolf, S. F., Sieburth, D. S. & Boom, W. H. (1996) Interleukin-12 production by human monocytes infected with Mycobacterium tuberculosis: role of phagocytosis. *Infection and Immunity* **64**, 2523-2531.
- Gemmell, E., Marshall, R. I. & Seymour, G. J. (1997) Cytokines and prostaglandins in immune homeostasis and tissue destruction in periodontal disease. *Periodontology 2000* **14**, 112-143.
- Gemmell, E. & Seymour, G. J. (1994) Modulation of immune responses to periodontal bacteria. *Current Opinion in Periodontology*, 28-38.
- Gemmell, E. & Seymour, G. J. (2004) Immunoregulatory control of Th1/Th2 cytokine profiles in periodontal disease. *Periodontology 2000* **35**, 21-41.
- Gemmell, E., Yamazaki, K. & Seymour, G. J. (2002) Destructive periodontitis lesions are determined by the nature of the lymphocytic response. *Critical Reviews in Oral Biology & Medicine* **13**, 17-34.
- Giambartolomei, G. H., Zwerdling, A., Cassataro, J., Bruno, L., Fossati, C. A. & Philipp, M. T. (2004) Lipoproteins, not lipopolysaccharide, are the key mediators of the proinflammatory response elicited by heat-killed Brucella abortus. *The Journal of Immunology* **173**, 4635-4642.
- Grossi, S. G., Genco, R. J., Machtei, E. E., Ho, A. W., Koch, G., Dunford, R., Zambon, J. J. & Hausmann, E. (1995) Assessment of risk for periodontal disease. II. Risk indicators for alveolar bone loss. *Journal of Periodontology* **66**, 23-29.
- Grossi, S. G., Zambon, J. J., Ho, A. W., Koch, G., Dunford, R. G., Machtei, E. E., Norderyd, O. M. & Genco, R. J. (1994) Assessment of risk for periodontal

- disease. I. Risk indicators for attachment loss. *Journal of Periodontology* **65**, 260-267.
- Kennedy, M. K., Picha, K. S., Fanslow, W. C., Grabstein, K. H., Alderson, M. R., Clifford, K. N., Chin, W. A. & Mohler, K. M. (1996) CD40/CD40 ligand interactions are required for T cell-dependent production of interleukin-12 by mouse macrophages. *European Journal of Immunology* **26**, 370-378.
- Kidd, P. (2003) Th1/Th2 balance: the hypothesis, its limitations, and implications for health and disease. *Alternative Medicine Reviews* **8**, 223-246.
- Kinane, D. F. & Lappin, D. F. (2001) Clinical, pathological and immunological aspects of periodontal disease. *Acta Odontologica Scandinavica* **59**, 154-160.
- Kurt-Jones, E. A., Mandell, L., Whitney, C., Padgett, A., Gosselin, K., Newburger, P. E. & Finberg, R. W. (2002) Role of toll-like receptor 2 (TLR2) in neutrophil activation: GM-CSF enhances TLR2 expression and TLR2-mediated interleukin 8 responses in neutrophils. *Blood* **100**, 1860-1868.
- Malave, I., Searles, R. P., Montano, J. & Williams, R. C., Jr. (1989) Production of tumor necrosis factor/cachectin by peripheral blood mononuclear cells in patients with systemic lupus erythematosus. *International Archives of Allergy & Applied Immunology* **89**, 355-361.
- Martinez-Canut, P., Lorca, A. & Magan, R. (1995) Smoking and periodontal disease severity. *Journal of Clinical Periodontology* **22**, 743-749.
- Muller, K., Herner, E. B., Stagg, A., Bendtzen, K. & Woo, P. (1998) Inflammatory cytokines and cytokine antagonists in whole blood cultures of patients with systemic juvenile chronic arthritis. *British Journal of Rheumatology* **37**, 562-569.
- Nakahara, T., Urabe, K., Fukagawa, S., Uchi, H., Inaba, K., Furue, M. & Moroi, Y. (2005) Engagement of human monocyte-derived dendritic cells into interleukin (IL)-12 producers by IL-1beta + interferon (IFN)-gamma. *Clinical & Experimental Immunology* **139**, 476-482.
- Niho, Y., Niiro, H., Tanaka, Y., Nakashima, H. & Otsuka, T. (1998) Role of IL-10 in the crossregulation of prostaglandins and cytokines in monocytes. *Acta Haematologica* **99**, 165-170.
- O'Garra, A. & Murphy, K. (1994) Role of cytokines in determining T-lymphocyte function. *Current Opinion in Immunology* **6**, 458-466.

- Pabst, M. J., Pabst, K. M., Collier, J. A., Coleman, T. C., Lemons-Prince, M. L., Godat, M. S., Waring, M. B. & Babu, J. P. (1995) Inhibition of neutrophil and monocyte defensive functions by nicotine. *Journal of Periodontology* **66**, 1047-1055.
- Page, R. C., Offenbacher, S., Schroeder, H. E., Seymour, G. J. & Kornman, K. S. (1997) Advances in the pathogenesis of periodontitis: summary of developments, clinical implications and future directions. *Periodontology 2000* **14**, 216-248.
- Petro, T. M., Anderson, L. L., Gowler, J. S., Liu, X. J. & Schwartzbach, S. D. (2002) Smokeless tobacco extract decreases IL-12 production from LPS-stimulated but increases IL-12 from IFN-gamma-stimulated macrophages. *International Immunopharmacology* **2**, 345-355.
- Prebeck, S., Kirschning, C., Durr, S., da Costa, C., Donath, B., Brand, K., Redecke, V., Wagner, H. & Miethke, T. (2001) Predominant role of toll-like receptor 2 versus 4 in Chlamydia pneumoniae-induced activation of dendritic cells. *The Journal of Immunology* **167**, 3316-3323.
- Pulendran, B., Kumar, P., Cutler, C. W., Mohamadzadeh, M., Van Dyke, T. & Banchereau, J. (2001) Lipopolysaccharides from distinct pathogens induce different classes of immune responses in vivo. *The Journal of Immunology* **167**, 5067-5076.
- Re, F. & Strominger, J. L. (2001) Toll-like receptor 2 (TLR2) and TLR4 differentially activate human dendritic cells. *J Biol Chem* **276**, 37692-37699.
- Sandborg, C. I., Imfeld, K. L., Zaldivar, F., Jr., Wang, Z., Buckingham, B. A. & Berman, M. A. (1995) IL-4 expression in human T cells is selectively inhibited by IL-1 alpha and IL-1 beta. *The Journal of Immunology* **155**, 5206-5212.
- Schmitz, J., Assenmacher, M. & Radbruch, A. (1993) Regulation of T helper cell cytokine expression: functional dichotomy of antigen-presenting cells. *European Journal of Immunology* **23**, 191-199.
- Seymour, G. J. & Gemmell, E. (2001a) Cytokines in periodontal disease: where to from here? *Acta Odontologica Scandinavica* **59**, 167-173.
- Seymour, G. J. & Gemmell, E. (2001b) Cytokines in periodontal disease: where to from here? *Acta Odontol Scand* **59**, 167-173.

- Sigusch, B., Klinger, G., Glockmann, E. & Simon, H. U. (1998) Early-onset and adult periodontitis associated with abnormal cytokine production by activated T lymphocytes. *Journal of Periodontology* **69**, 1098-1104.
- Snijders, A., Hilkens, C. M., van der Pouw Kraan, T. C., Engel, M., Aarden, L. A. & Kapsenberg, M. L. (1996) Regulation of bioactive IL-12 production in lipopolysaccharide-stimulated human monocytes is determined by the expression of the p35 subunit. *The Journal of Immunology* **156**, 1207-1212.
- Snijders, A., Van der Pouw Kraan, T. C., Engel, M., Wormmeester, J., Widjaja, P., Zonneveld, I. M., Bos, J. D. & Kapsenberg, M. L. (1998) Enhanced prostaglandin E2 production by monocytes in atopic dermatitis (AD) is not accompanied by enhanced production of IL-6, IL-10 or IL-12. *Clinical & Experimental Immunology* **111**, 472-476.
- Swaak, A. J., van den Brink, H. G. & Aarden, L. A. (1997) Cytokine production in whole blood cell cultures of patients with rheumatoid arthritis. *Annals of the Rheumatic Diseases* **56**, 693-695.
- van den Heuvel, M. M., Vanhee, D. D., Postmus, P. E., Hoefsmit, E. C. & Beelen, R. H. (1998) Functional and phenotypic differences of monocyte-derived dendritic cells from allergic and nonallergic patients. *Journal of Allergy and Clinical Immunology* **101**, 90-95.
- van der Pouw Kraan, T. C., Boeije, L. C., de Groot, E. R., Stapel, S. O., Snijders, A., Kapsenberg, M. L., van der Zee, J. S. & Aarden, L. A. (1997) Reduced production of IL-12 and IL-12-dependent IFN-gamma release in patients with allergic asthma. *The Journal of Immunology* **158**, 5560-5565.
- van der Pouw Kraan, T. C., Boeije, L. C., Smeenk, R. J., Wijdenes, J. & Aarden, L. A. (1995) Prostaglandin-E2 is a potent inhibitor of human interleukin 12 production. *Journal of Experimental Medicine* **181**, 775-779.
- van Lier, R. A., Boot, J. H., De Groot, E. R. & Aarden, L. A. (1987a) Induction of T cell proliferation with anti-CD3 switch-variant monoclonal antibodies: effects of heavy chain isotype in monocyte-dependent systems. *European Journal of Immunology* **17**, 1599-1604.
- van Lier, R. A., Boot, J. H., Verhoeven, A. J., de Groot, E. R., Brouwer, M. & Aarden, L. A. (1987b) Functional studies with anti-CD3 heavy chain isotype switch-variant monoclonal antibodies. Accessory cell-independent induction

- of interleukin 2 responsiveness in T cells by epsilon-anti-CD3. *The Journal of Immunology* **139**, 2873-2879.
- van Winkelhoff, A. J., Loos, B. G., van der Reijden, W. A. & van der Velden, U. (2002) Porphyromonas gingivalis, Bacteroides forsythus and other putative periodontal pathogens in subjects with and without periodontal destruction. *Journal of Clinical Periodontology* **29**, 1023-1028.
- Venier, C., Guthmann, M. D., Fernandez, L. E. & Fainboim, L. (2007) Innate-immunity cytokines induced by very small size proteoliposomes, a Neisseria-derived immunological adjuvant. *Clinical & Experimental Immunology* **147**, 379-388.
- Zaitseva, M., Golding, H., Manischewitz, J., Webb, D. & Golding, B. (1996) Brucella abortus as a potential vaccine candidate: induction of interleukin-12 secretion and enhanced B7.1 and B7.2 and intercellular adhesion molecule 1 surface expression in elutriated human monocytes stimulated by heat-inactivated B. abortus. *Infection and Immunity* **64**, 3109-3117.
- Zambon, J. J., Grossi, S. G., Machtei, E. E., Ho, A. W., Dunford, R. & Genco, R. J. (1996) Cigarette smoking increases the risk for subgingival infection with periodontal pathogens. *Journal of Periodontology* **67**, 1050-1054.