UvA-DARE (Digital Academic Repository)

Conducting-angle-based percolation in the XY model

Wang, Y.; Guo, W.; Nienhuis, B.; Blöte, H.W.J.

DOI

[10.1103/PhysRevE.81.031117](https://doi.org/10.1103/PhysRevE.81.031117)

Publication date

2010

Document Version

Final published version

Published in

Physical Review E

[Link to publication](#)

Citation for published version (APA):

Wang, Y., Guo, W., Nienhuis, B., & Blöte, H. W. J. (2010). Conducting-angle-based percolation in the XY model. *Physical Review E*, 81(3), 031117. <https://doi.org/10.1103/PhysRevE.81.031117>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Conducting-angle-based percolation in the XY model

Yancheng Wang,¹ Wenan Guo,^{1,*} Bernard Nienhuis,² and Henk W. J. Blöte³

¹*Physics Department, Beijing Normal University, Beijing 100875, People's Republic of China*

²*Instituut voor Theoretische Fysica, Universiteit van Amsterdam, Valckenierstraat 65, The Netherlands*

³*Instituut Lorentz, Leiden University, P.O. Box 9506, 2300 RA Leiden, The Netherlands*

(Received 18 December 2009; published 17 March 2010)

We define a percolation problem on the basis of spin configurations of the two-dimensional XY model. Neighboring spins belong to the same percolation cluster if their orientations differ less than a certain threshold called the conducting angle. The percolation properties of this model are studied by means of Monte Carlo simulations and a finite-size scaling analysis. Our simulations show the existence of percolation transitions when the conducting angle is varied, and we determine the transition point for several values of the XY coupling. It appears that the critical behavior of this percolation model can be well described by the standard percolation theory. The critical exponents of the percolation transitions, as determined by finite-size scaling, agree with the universality class of the two-dimensional percolation model on a uniform substrate. This holds over the whole temperature range, even in the low-temperature phase where the XY substrate is critical in the sense that it displays algebraic decay of correlations.

DOI: [10.1103/PhysRevE.81.031117](https://doi.org/10.1103/PhysRevE.81.031117)

PACS number(s): 05.50.+q, 64.60.Cn, 75.10.Hk

I. INTRODUCTION

Consider the XY or planar model on the square lattice with periodic boundary conditions, described by the reduced Hamiltonian

$$\mathcal{H} = -\frac{J}{k_B T} \sum_{\langle ij \rangle} \vec{s}_i \cdot \vec{s}_j, \quad (1)$$

where the sum is over all nearest-neighbor pairs and the $\vec{s}_i$ are two-dimensional unit vectors labeled by the site number i . We restrict the nearest-neighbor interaction to be ferromagnetic, i.e., $J > 0$.

When the temperature of this model is lowered, it undergoes a phase transition of an interesting character, which was explained by Kosterlitz and Thouless [1]. More exact results for the exponents were obtained by Nienhuis [2] for an O(2) model in the same universality class. These results show that the renormalization exponent y_t of the temperature is equal to 0, which means that the temperature-driven transition is of infinite order, i.e., the specific-heat singularity is extremely weak. In contrast, the magnetic susceptibility displays a very strong divergence when the temperature is lowered to the transition point.

At temperatures $T > 0$ below the transition point there is no spontaneous long-range order in the sense that the magnetization is zero [3]. Instead, the low-temperature phase resembles a critical state; the correlations decay algebraically, with exponents that are still dependent on the temperature.

Recently, there has been ample attention to percolation problems defined on a critical substrate, see e.g., Refs. [4–8] and references therein. It thus appears that, for Potts and O(n) models, the universal properties of such percolation transitions do reflect the nature of the critical substrate.

These investigations are based on model representations with discrete degrees of freedom.

It is thus interesting to investigate related problems using a substrate with continuous degrees of freedom. For instance, the mechanical properties of static granular matter can be analyzed in terms of the so-called force networks [9,10]. By introducing a threshold force, such that forces exceeding the threshold form clusters, a percolation transition is seen at a critical force threshold. This approach was applied to different models for granular piles that are expected to belong to different universality classes. It was found that also the corresponding percolation behavior could discriminate between the different models [11]. This might suggest that such substrate dependence is a general phenomenon for critical models with continuous degrees of freedom, i.e., the critical behavior of the percolation clusters might generally reflect the long-range correlations of the original degrees of freedom.

In order to shed more light on this issue, the present work investigates a percolation problem using the substrate of the two-dimensional XY model. In particular we are interested how the universal properties of the percolation transition will depend on the temperature of the underlying XY model. We define the percolation problem such as to depend only on the XY configuration and not on any additional random variables. This is achieved by introducing a “conducting angle” θ such that neighboring spins whose orientations differ by less than θ are connected by a percolation bond. This name is based on the analogy with a conductance problem of conducting units on a lattice such that neighboring units are in electrical contact only if their orientations match to a sufficient degree.

It is clear that this conducting angle also defines a threshold pair energy, below which a pair of neighboring XY spins is connected by a percolation bond. These percolation bonds define a bond percolation configuration involving a complete decomposition of the lattice in percolation clusters. An example of an XY configuration with the corresponding percolation cluster decomposition is shown in Fig. 1. It is obvious that the resulting bond percolation configurations will not

*waguo@bnu.edu.cn

FIG. 1. (Color online) Construction of a bond percolation configuration (b) from an XY spin configuration (a). If the angle θ_{ij} between a pair (i, j) of neighboring spins is less than a given angle θ , the neighboring spins are connected by a bond, so that the lattice decomposes into a system of percolation clusters. The threshold was chosen as $\theta = \pi/6$ in this figure.

percolate for $\theta=0$ and that they will percolate for $\theta=\pi$ and larger. We are thus left with the task to find the percolation threshold θ_c and the critical exponents as a function of the temperature. To answer these questions, we perform simulations of the two-dimensional (2D) XY spin model, which are described in Sec. II. Section III presents the numerical results, including the critical parameters. We include a short discussion in Sec. IV.

II. ALGORITHM

For reasons of efficiency, we make use of a cluster algorithm [12] to simulate the XY model. We applied the single-cluster algorithm formulated by Wolff [13] to the model on the square lattice. We recall the steps involved in one Wolff cluster flip:

(1) choose an arbitrary direction as the y direction and denote by α_i the angle between the spin $\vec{s}_i$ and the y axis. Define an Ising spin $s_i = \pm 1$ with the same sign as the y component of $\vec{s}_i$. Thus, the nearest-neighbor interaction term in $\mathcal{H}_{ij}$ between spins i and j reads

$$\begin{aligned} \mathcal{H}_{ij} = & -K \vec{s}_i \cdot \vec{s}_j = -K s_i^x s_j^x - K s_i^y s_j^y = -K s_i^x s_j^x \\ & - K \cos(\alpha_i) \cos(\alpha_j) s_i s_j, \end{aligned} \quad (2)$$

where $K \equiv J/k_B T$. As far as the dependence of $\mathcal{H}_{ij}$ on the Ising variables s_i and s_j is concerned, this is an Ising coupling between the two spins with strength $K_{ij} = K \cos \alpha_i \cos \alpha_j$. One may update the Ising variables using a cluster algorithm that takes into account these position-dependent couplings. The following steps are used to update the y components of the spins.

(2) Choose a spin randomly, say on site i . For each nearest-neighbor site j of i , connect i and j by a bond with probability $p_{ij} = \max(0, 1 - e^{-2K_{ij}})$. Then do the same for each of the nearest-neighbor sites of each newly connected site, and so on. The process continues until no more new sites are connected. Then, the construction of the cluster, which contains all sites connected via some path of bonds to site i , is finished.

(3) Change the sign of the y components of all spins in that cluster.

III. NUMERICAL RESULTS AND ANALYSIS

During the simulations, we constructed percolation cluster decompositions on the basis of a chosen conducting angle θ . For each such decomposition we sampled several quantities in order to estimate the second moment of the cluster size distribution S_2 , the probability P that there exists a cluster that wraps the system, a dimensionless ratio Q that can be related to the Binder cumulant, and the cluster size distribution function n_s . We shall proceed to introduce these quantities in more detail and list their expected finite-size scaling properties.

Since we are using finite systems with periodic boundary conditions, we define a “wrapping cluster” [14] as a percolation cluster that connects to itself along at least one of the periodic directions. For each XY configuration S we thus define a quantity $p(S, \theta)$ that has the value 1 if there exists a wrapping cluster, and $p(S, \theta) = 0$ otherwise. Thus, for a system with finite-size L , the probability $P(K, \theta, L)$ that a wrapping cluster exists is given by

$$P(K, \theta, L) = \langle p(S, \theta) \rangle, \quad (3)$$

where the ensemble average is taken for an XY system with linear size L and coupling K . If there is a percolation transition at a conducting angle θ_c , one expects that θ plays the role of a temperaturelike variable and thus that the finite-size scaling behavior [15] of P is described by

$$P(K, \theta, L) = P^{(c)} + a(\theta - \theta_c)L^{y_p} + \cdots + b_1 L^{y_1} + b_2 L^{y_2} + \cdots, \quad (4)$$

where y_p controls the scaling of θ and acts as a temperaturelike exponent, corresponding with the bond dilution exponent in the language of the percolation model. The exponents y_1 , y_2 , etc. are correction-to-scaling exponents, which are unknown in principle. The constant $P^{(c)}$, which is defined as the value of P at θ_c in the limit $L \rightarrow \infty$, and the exponent y_p are universal, but the universality class of the present percolation problem remains to be determined.

The second moment S_2 of the percolation cluster size distribution can also be viewed as the mean size of the cluster containing an arbitrary point. It is defined as

$$S_2 = \frac{1}{N^2} \left\langle \sum_i^{N_c} s_i^2 \right\rangle, \quad (5)$$

where s_i is the size of the i -th cluster, N_c the total number of clusters for a configuration, and $N = L^2$ is the volume of the system. The quantity S_2 is also closely related with the generalization χ_{RC} of the q -state Potts magnetic susceptibility to the random-cluster model, applied to the special case $q=1$. This relation is expressed as $\chi_{RC} = N S_2$.

In analogy with the Potts susceptibility, we expect the following finite-size-scaling behavior for S_2 in the neighborhood of a percolation threshold at θ_c

$$S_2 = L^{2y_h - 2d} [a_0 + a_1(\theta - \theta_c)L^{y_p} + \cdots + b_1 L^{y_1} + b_2 L^{y_2} + \cdots], \quad (6)$$

where y_h is the fractal dimension of critical percolation clusters. At the percolation threshold θ_c , this equation reduces to

FIG. 2. (Color online). Wrapping probability versus θ at $K=0.5$. The curves are added as a guide for the eye.

$$S_2 = L^{2y_h - 2d} (a_0 + b_1 L^{y_1} + b_2 L^{y_2} + \dots). \quad (7)$$

We also define a dimensionless ratio related to the Binder cumulant [16] as

$$Q = \frac{S_2^2}{3S_2^{(2)} - 2S_4}, \quad (8)$$

where $S_2^{(2)}$ is defined as

$$S_2^{(2)} = \frac{1}{N^4} \left\langle \left(\sum_i^{N_c} s_i^2 \right)^2 \right\rangle, \quad (9)$$

and S_4 as

$$S_4 = \frac{1}{N^4} \left\langle \sum_i^{N_c} s_i^4 \right\rangle. \quad (10)$$

The relation with the Binder cumulant is based on the fact that Eq. (8) is obtained when the Binder ratio $\langle m^2 \rangle^2 / \langle m^4 \rangle$ of magnetization moments of the Ising model is expressed in the language of the $q=2$ random-cluster model. In analogy with the Binder cumulant, we expect that, in the neighborhood of the percolation threshold, it behaves like

$$Q(K, \theta, L) = Q^{(c)} + a_1(\theta - \theta_c)L^{y_p} + \dots + b_1 L^{y_1} + b_2 L^{y_2} + \dots. \quad (11)$$

A. Wrapping probability as a function of θ

We simulated XY systems with linear sizes $L=8, 16, 32, 64, 128$, and 256 at different inverse temperatures: $K=0, 0.001, 0.01, 0.10, \dots, 5.00$, and 10.00 . For $K=0.9$ and 1.0 , system sizes up to $L=1024$ and 2048 were simulated, respectively. Typically, 10^7 XY configurations were sampled. The samples were taken at intervals consisting of one Metropolis sweep and four Wolff clusters. For each of these XY configurations, percolation cluster decompositions were constructed with several different values of the conducting angle.

Some results for the wrapping probability $P(K, \theta, L)$ as a function of the conducting angle are shown in Figs. 2–4. These figures show clear intersections corresponding with percolation thresholds. Furthermore, the behavior appears to

FIG. 3. (Color online). Wrapping probability versus θ at $K=1$. The lower figure shows the details in the vicinity of θ_c , where the finite-size effect becomes relatively strong. The curves are added as a guide for the eye.

be remarkably similar for different values of the XY coupling K . We applied the least-squares method to fit the wrapping probability by the finite-size scaling Eq. (4). The results for the percolation threshold θ_c are shown in Fig. 5 as a function of the XY coupling K . They are also included in Table I for each value of K , together with the results for the exponent y_p and the universal probability $P^{(c)}$. The results for the latter quantity reproduce, within error bounds, the literature value $P^{(c)}=0.690473725$ which applies to the ordinary two-dimensional percolation model [14,17,18].

FIG. 4. (Color online). Wrapping probability versus θ at $K=2.0$. The curves are added as a guide for the eye.

FIG. 5. (Color online) Percolation threshold θ_c versus K . For $K=0$, the critical value of the conducting angle is 1.610 78 radians or about 92 degrees. This value applies to a system of randomly oriented conductors on the square lattice. The curve is added as a guide for the eyes and estimated error bars are smaller than the size of the symbols.

B. Dimensionless ratio Q as a function of θ

We also sampled the ratio Q for systems at different inverse temperatures. Some of the data for $K=1.0$ are shown in Fig. 6. These data display the same general behavior as the wrapping probability in Sec. III A. Furthermore, the data for other values of K (not shown) display a very similar behavior, as also seen in the preceding subsection.

A least-squares analysis of the data for $Q(K, \theta, L)$ on the basis of the finite-size scaling Eq. (11) results in the estimates of y_p , $Q^{(c)}$, and the percolation threshold θ_c . The estimates of y_p and θ_c are in agreement with those found by fitting $P(K, \theta, L)$. The estimates of the universal value $Q^{(c)}$

FIG. 6. (Color online). Dimensionless ratio Q versus conducting angle θ for $K=1.0$. The curves are added as a guide for the eye.

are also listed in Table I for each K . Their values are in agreement with the literature value [19] 0.870 53(2) for the ordinary two-dimensional percolation model.

C. Second moment S_2 at the percolation threshold

We simulated the model with system sizes $L=8, 16, 32, 64, 128, 256$, and 512 at the estimated percolation threshold θ_c for each K and sampled the second moment S_2 of the cluster size distribution. Samples were taken at intervals consisting of one Metropolis sweep and four Wolff clusters. The numbers of samples taken for system sizes $L=8$ to 32 , $L=64$ to 128 , and $L=256$ to 512 are 10^7 , 4×10^6 , and 2×10^6 , respectively. For $K=1.0$, which is in the vicinity of the Kosterlitz-Thouless (KT) transition, additional simulations took place for system sizes $L=1024$ and 2048 , involv-

TABLE I. Numerical results for the percolation threshold θ_c , the bond dilution exponent y_p , the fractal dimension y_h , the universal wrapping probability $P^{(c)}$, and the universal dimensionless ratio $Q^{(c)}$. The exact values for two-dimensional percolation exponents are $y_p=3/4$ and $y_h=91/48=1.895833 \dots$

K	θ_c	y_p	y_h	$P^{(c)}$	$Q^{(c)}$
0	1.61078(2)	0.750(2)	1.8961(3)	0.6906(2)	0.8708(3)
0.001	1.60994(2)	0.748(3)	1.8961(3)	0.6908(3)	0.8707(5)
0.010	1.60252(2)	0.750(2)	1.8959(3)	0.6905(3)	0.8706(3)
0.100	1.52866(2)	0.751(2)	1.8959(3)	0.6905(3)	0.8708(5)
0.200	1.44651(2)	0.752(2)	1.8958(3)	0.6903(4)	0.8708(5)
0.500	1.18179(2)	0.750(2)	1.8956(3)	0.6903(3)	0.8709(5)
0.750	0.91074(2)	0.751(2)	1.8957(3)	0.6901(5)	0.8704(5)
0.800	0.85002(2)	0.749(5)	1.8957(3)	0.6905(3)	0.8702(5)
0.900	0.72804(2)	0.751(4)	1.8958(3)	0.6902(4)	0.8705(5)
1.000	0.62209(2)	0.74 (2)	1.894 (2)	0.6907(4)	0.8705(5)
1.100	0.55695(1)	0.753(3)	1.8954(5)	0.6902(6)	0.8709(5)
1.120	0.54815(1)	0.752(3)	1.8958(2)	0.690 (1)	0.8706(5)
1.200	0.51869(1)	0.751(2)	1.8959(2)	0.6903(5)	0.8707(5)
1.500	0.44598(2)	0.753(3)	1.897 (2)	0.6907(3)	0.8708(5)
2.000	0.37560(2)	0.752(3)	1.895 (1)	0.6900(8)	0.8708(3)
5.000	0.22851(1)	0.751(2)	1.8958(3)	0.690 (1)	0.8708(5)
10.00	0.15981(2)	0.751(4)	1.895 (1)	0.689 (3)	0.8708(5)

FIG. 7. (Color online) Second moment S_2 of the cluster size distribution versus finite-size L , for XY coupling $K=1.0$. The curve is added as a guide for the eyes and estimated error bars are smaller than the size of the symbols.

ing several times 10^6 samples per system size.

Some data for S_2 are shown as a function of the system size L in Fig. 7. Since logarithmic scales are used, linear behavior corresponds with power-law behavior, in accordance with scaling. This figure applies to the case $K=1.0$, which is close to the KT transition. Small deviations from linearity, which are due to corrections to scaling, are visible. Plots for other values of K (not shown) display a somewhat better approximation to linear behavior.

We fitted S_2 by the finite-size-scaling formula Eq. (7) and thus obtained estimates of the fractal dimension y_h , which are listed in Table I for several values of K . For $K=1.0$, a satisfactory fit could only be obtained by discarding system sizes $L < 128$ and including system sizes up to $L=2048$.

D. Distribution n_s of the cluster size s

At the percolation transition θ_c , the requirement that the cluster size distribution scales in a covariant way yields a power law for this distribution

$$n_s(s, \theta_c) \propto s^{-\tau}, \quad (12)$$

where τ is a critical exponent equal to $1+d/y_h$, with $d=2$ dimensions and the fractal dimension y_h of the percolation clusters. For the ordinary $d=2$ percolation model, one has $y_h=91/48$ [20,21], and thus $\tau=187/91$.

An example of the data for n_s is shown in Fig. 8 as a function of cluster size s , for XY coupling $K=1.0$. This figure uses logarithmic scales. The data points are, in a wide range of cluster sizes, well approximated by straight lines, corresponding with a scale-invariant distribution according to a power law. The slopes of these lines are close to the value of τ that applies to the ordinary percolation model. The curves in Fig. 8 can obviously be brought to an approximate data collapse by introducing suitable prefactors, but it appears that there is still an appreciable finite-size dependence. Similar plots for other values of K (not shown) display practically the same behavior.

For a numerical determination of τ from the data for n_s , we use a least-squares analysis in the range of s where n_s is almost linear when plotted as in Fig. 8, thus excluding small

FIG. 8. (Color online) Cluster size distribution function as a function of cluster size s for $K=1.0$, $\theta_c=0.62210$. The curves are seen to shift to the right with increasing L .

clusters smaller than 20, and the largest clusters exceeding a size $L^{y_h}/10$. Even then, two correction terms had to be included in order to obtain satisfactory residuals, according to the fit formula

$$n_s(s, \theta_c, L) = a_0 s^{-\tau} (1 + a_1 s^{y_1} + a_2 L^{y_2}), \quad (13)$$

where the term with coefficient a_2 is significant only for $K > 1$. Satisfactory fits were obtained for all inverse temperatures. For example, for $K=0.75$, we fitted the data simultaneously for system sizes $L=32, 64, \dots, 512$ using the above formula and found $\tau=2.055(2)$, $y_1=-0.57(1)$ with χ^2 per degree of freedom almost equal to 1. For $K=2.0$, a fit of the data for system sizes $L=32, 64, \dots, 512$ yielded $\tau=2.054(2)$, $y_1=-1.0(1)$, and $y_2=-1.6(1)$, again with a satisfactory value of χ^2 as compared with the number of degrees of freedom.

We thus obtained estimates of the exponent τ that agree well with the value $\tau=187/91=2.0549\dots$ for the ordinary 2D percolation model, for several temperatures of the XY model.

IV. DISCUSSION

The numerical results presented in Sec. III for the exponents τ , y_h , and y_p , as well as for the universal probability $P^{(c)}$ and the universal ratio $Q^{(c)}$, agree accurately with the literature values applying to the two-dimensional percolation model. Thus our analysis shows the existence of a transition in the ordinary percolation universality class, independent of the XY temperature. This result is as expected in the high-temperature phase where the XY spins display strong random disorder, but may seem somewhat surprising in the critical region including the low-temperature range. The problem discussed here can be viewed as a correlated percolation problem, in which the bond probabilities correlate as the nearest-neighbor differences in the XY model. For such long-range correlated percolation Weinrib [22] formulated a generalized Harris criterion to decide if the correlations are relevant for the percolation behavior or not. The critical behavior is expected to be in the universality class of ordinary (uncorrelated) percolation if $\alpha\nu > 2$ for correlations decaying with distance as $r^{-\alpha}$ with ν the percolation correlation

length exponent $\nu=4/3$. Indeed in the case at hand $a=2$ and the correlations should thus be irrelevant. It stands in a strong contrast with a recent analysis [7] of a percolation problem defined on the basis of the two-dimensional $O(n)$ model where n is a continuously variable parameter. For the $O(2)$ model, which belongs to the same universality class as the XY model, it was found [7] that the percolation transition was driven by a bond dilution field that is only marginally relevant, i.e., $y_p=0$ at that transition, while $y_p=3/4$ for ordinary percolation. It thus appears that the character of a percolation transition on a critical substrate depends on the precise definition of the percolation problem. The percolation problem of Ref. [7] was defined within regions separated by loops as defined in the context of the $O(n)$ loop model. These loops obviously display fractal properties [6] at $O(n)$ criticality and thus affect the nature of the percolation transition. In contrast, the spatial variation in the angle between neighbor-

ing XY spins is very smooth near criticality, and does not display boundarylike structures, even at the KT transition. While it is natural that the KT transition is reflected in some way in the behavior of the percolation transition at the critical value of the conducting angle such effects are exposed by our numerical results only in terms of slow convergence in the analysis of the finite-size data.

ACKNOWLEDGMENTS

W.G. acknowledges hospitality extended to him by the Lorentz Institute. This work is supported by the Lorentz Fund, by the National Science Foundation of China (NSFC) under Grant No. 10675021, by the Program for New Century Excellent Talents in University (NCET), and by the HSCC (High Performance Scientific Computing Center) of the Beijing Normal University.

-
- [1] J. M. Kosterlitz and D. J. Thouless, *J. Phys. C* **5**, L124 (1972); **6**, 1181 (1973); J. M. Kosterlitz, *ibid.* **7**, 1046 (1974).
 - [2] B. Nienhuis, *Phys. Rev. Lett.* **49**, 1062 (1982).
 - [3] N. D. Mermin and H. Wagner, *Phys. Rev. Lett.* **17**, 1133 (1966); N. D. Mermin, *J. Math. Phys.* **8**, 1061 (1967).
 - [4] X.-F. Qian, Y. Deng, and H. W. J. Blöte, *Phys. Rev. B* **71**, 144303 (2005).
 - [5] Y. Deng, H. W. J. Blöte, and B. Nienhuis, *Phys. Rev. E* **69**, 026123 (2004).
 - [6] C.-X. Ding, Y. Deng, W.-A. Guo, X.-F. Qian, and H. W. J. Blöte, *J. Phys. A* **40**, 3305 (2007).
 - [7] C.-X. Ding, Y. Deng, W.-A. Guo, and H. W. J. Blöte, *Phys. Rev. E* **79**, 061118 (2009).
 - [8] The equivalence of the critical percolation model and the dense phase of the $O(0)$ loop model indicates a relation of this class of percolation problems with the self-avoiding loops on an Ising substrate studied by F. Seno, A. Stella, and C. Vanderzande, *Phys. Rev. Lett.* **65**, 2897 (1990).
 - [9] *Powers and Grains 2005*, edited by R. Garcia-Rojo, H. J. Herrmann, and S. McNamara (Balkema, Rotterdam, 2005).
 - [10] N. M. Jaeger, S. R. Nagel, and R. P. Behringer, *Rev. Mod. Phys.* **68**, 1259 (1996).
 - [11] S. Ostojic, E. Somfai, and B. Nienhuis, *Nature (London)* **439**, 828 (2006).
 - [12] R. H. Swendsen and J. S. Wang, *Phys. Rev. Lett.* **58**, 86 (1987).
 - [13] U. Wolff, *Phys. Rev. Lett.* **62**, 361 (1989).
 - [14] H. T. Pinson, *J. Stat. Phys.* **75**, 1167 (1994).
 - [15] For a review, see, e.g. M. P. Nightingale, in *Finite-Size Scaling and Numerical Simulation of Statistical Systems*, edited by V. Privman (World Scientific, Singapore, 1990).
 - [16] K. Binder, *Z. Phys. B* **43**, 119 (1981).
 - [17] R. M. Ziff, C. D. Lorenz, and P. Kleban, *Physica A* **266**, 17 (1999).
 - [18] M. E. J. Newman and R. M. Ziff, *Phys. Rev. Lett.* **85**, 4104 (2000).
 - [19] Y. Deng and H. W. J. Blöte, *Phys. Rev. E* **71**, 016117 (2005).
 - [20] B. Nienhuis, in *Phase Transitions and Critical Phenomena*, edited by C. Domb and J. L. Lebowitz (Academic, London, 1987), Vol. 11.
 - [21] J. L. Cardy, in *Phase Transitions and Critical Phenomena*, edited by C. Domb and J. L. Lebowitz (Academic, London, 1987), Vol. 11.
 - [22] A. Weinrib, *Phys. Rev. B* **29**, 387 (1984).